



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 8, 2024 – 07:03 AM EDT

PDB ID : 4QHU
Title : Crystal Structure of IL-17A/Fab6785 complex
Authors : Luo, J.; Gilliland, G.L.; Malia, T.; Obmolova, G.; Teplyakov, A.
Deposited on : 2014-05-29
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION






The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	5791 (2.20-2.20)
Clashscore	180529	6634 (2.20-2.20)
Ramachandran outliers	177936	6560 (2.20-2.20)
Sidechain outliers	177891	6561 (2.20-2.20)
RSRZ outliers	164620	5791 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	214	 2% 91% 7%
1	L	214	 0% 92% 6%
2	B	224	 0% 83% 9% 6%
2	H	224	 4% 82% 11% 6%
3	C	132	 4% 79% 11% 10%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	D	132	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	CL	L	303	-	-	X	-

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 9238 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fab6785 light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	L	209	1560	980	256	319	5	0	0	0
1	A	209	1559	980	256	318	5	0	0	0

- Molecule 2 is a protein called Fab6785 heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	H	211	1570	993	262	309	6	0	0	0
2	B	210	1573	997	263	307	6	0	1	0

- Molecule 3 is a protein called Interleukin-17A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	119	967	596	184	179	8	0	1	0
3	D	100	804	501	148	148	7	0	1	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	70	GLN	LYS	engineered mutation	UNP Q16552
C	132	GLN	-	expression tag	UNP Q16552
D	70	GLN	LYS	engineered mutation	UNP Q16552
D	132	GLN	-	expression tag	UNP Q16552

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	L	1	Total C O 6 3 3	0	0
4	L	1	Total C O 6 3 3	0	0
4	H	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0

- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	L	1	Total Cl 1 1	0	0

- Molecule 6 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	H	1	Total	O	S	0	0
			5	4	1		
6	B	1	Total	O	S	0	0
			5	4	1		
6	C	1	Total	O	S	0	0
			5	4	1		
6	C	1	Total	O	S	0	0
			5	4	1		
6	D	1	Total	O	S	0	0
			5	4	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	L	205	Total	O	0	0
			205	205		
7	H	200	Total	O	0	0
			200	200		
7	A	232	Total	O	0	0
			232	232		
7	B	281	Total	O	0	0
			281	281		
7	C	125	Total	O	0	0
			125	125		
7	D	106	Total	O	0	0
			106	106		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Fab6785 light chain




- Molecule 1: Fab6785 light chain




- Molecule 2: Fab6785 heavy chain




- Molecule 2: Fab6785 heavy chain




- Molecule 3: Interleukin-17A





- Molecule 3: Interleukin-17A



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	73.21Å 64.00Å 145.60Å 90.00° 95.64° 90.00°	Depositor
Resolution (Å)	48.30 – 2.20 48.30 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.3 (48.30-2.20) 99.3 (48.30-2.20)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.16 (at 2.20Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, R_{free}	0.171 , 0.216 0.174 , 0.218	Depositor DCC
R_{free} test set	3452 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	20.1	Xtrriage
Anisotropy	0.256	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 51.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	9238	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.78% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CL, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.23	0/1598	0.41	0/2184
1	L	0.23	0/1599	0.41	0/2187
2	B	0.25	0/1611	0.44	0/2195
2	H	0.23	0/1605	0.42	0/2188
3	C	0.23	0/995	0.41	0/1357
3	D	0.23	0/825	0.40	0/1123
All	All	0.23	0/8233	0.42	0/11234

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1559	0	1495	11	0
1	L	1560	0	1495	8	0
2	B	1573	0	1559	13	0
2	H	1570	0	1548	16	0
3	C	967	0	924	12	0
3	D	804	0	765	11	0
4	A	12	0	16	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	H	6	0	8	0	0
4	L	12	0	16	1	0
5	L	1	0	0	2	0
6	B	5	0	0	0	0
6	C	10	0	0	0	0
6	D	5	0	0	0	0
6	H	5	0	0	0	0
7	A	232	0	0	7	1
7	B	281	0	0	2	0
7	C	125	0	0	3	0
7	D	106	0	0	3	0
7	H	200	0	0	2	1
7	L	205	0	0	3	0
All	All	9238	0	7826	67	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:96:MET:SD	7:B:524:HOH:O	2.12	1.07
5:L:303:CL:CL	7:L:583:HOH:O	2.15	1.01
1:L:191:ARG:NH1	5:L:303:CL:CL	2.41	0.90
1:L:191:ARG:NH2	7:L:538:HOH:O	2.09	0.83
3:C:94:GLN:OE1	3:D:94:GLN:NE2	2.13	0.80
1:A:3:VAL:O	7:A:510:HOH:O	2.03	0.76
2:H:136:GLY:N	7:H:592:HOH:O	2.22	0.71
1:L:203:THR:O	7:L:508:HOH:O	2.08	0.71
2:H:59:TYR:OH	3:C:101[B]:ARG:NH2	2.25	0.70
1:A:53:ARG:HD2	7:A:460:HOH:O	1.91	0.69
2:H:119:LYS:NZ	7:H:486:HOH:O	2.26	0.68
1:L:92:PHE:O	3:C:101[B]:ARG:NH1	2.27	0.67
3:C:105:HIS:ND1	7:C:328:HOH:O	2.14	0.67
2:H:121:PRO:HB3	2:H:147:TYR:HB3	1.78	0.64
1:A:52:ASP:OD2	7:A:522:HOH:O	2.15	0.63
2:H:128:PRO:HG3	2:H:140:LEU:HB3	1.82	0.62
3:C:106:CYS:HB3	7:C:375:HOH:O	2.00	0.60
2:B:12:VAL:HG11	2:B:86:LEU:HD13	1.84	0.60
3:C:12:ASN:H	3:D:31:ARG:HH21	1.50	0.58
4:A:301:GOL:O3	3:D:55:ARG:NH1	2.36	0.58

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:64:VAL:HG13	2:B:68:PHE:HB2	1.85	0.58
2:B:121:PRO:HB3	2:B:147:TYR:HB3	1.87	0.57
1:L:31:TYR:HB3	1:L:49:ASP:HA	1.87	0.56
1:L:44:VAL:HG11	4:L:301:GOL:H12	1.88	0.56
2:B:128:PRO:HG3	2:B:140:LEU:HB3	1.87	0.55
1:A:158:LYS:NZ	7:A:592:HOH:O	2.40	0.54
1:A:36:GLN:HE22	4:A:302:GOL:H11	1.73	0.54
2:H:161:LEU:HD21	2:H:184:VAL:HG11	1.90	0.53
1:A:53:ARG:NH1	7:A:460:HOH:O	2.17	0.53
2:B:173:GLN:HG2	2:B:177:LEU:O	2.09	0.52
2:H:98:ARG:HB3	2:H:104:VAL:HG13	1.91	0.52
2:B:128:PRO:HD2	2:B:215:PRO:HA	1.91	0.52
2:B:3[A]:GLN:NE2	7:B:522:HOH:O	2.36	0.51
3:D:30:ASN:HD22	3:D:31:ARG:H	1.61	0.49
1:A:53:ARG:NE	1:A:61:PHE:O	2.40	0.49
3:C:92:ILE:HD12	3:D:92:ILE:HD12	1.95	0.48
2:H:165:VAL:HG22	2:H:184:VAL:HG22	1.93	0.48
4:A:301:GOL:H12	7:A:492:HOH:O	2.14	0.48
2:B:98:ARG:HB3	2:B:104:VAL:HG13	1.96	0.47
3:D:55:ARG:NH1	3:D:57:GLU:OE1	2.36	0.47
3:C:97:LEU:HD23	3:C:114:LYS:HG2	1.97	0.47
2:H:12:VAL:HG11	2:H:86:LEU:HD13	1.96	0.46
3:D:108:ASN:O	7:D:316:HOH:O	2.20	0.46
2:H:83:MET:HE2	2:H:86:LEU:HD21	1.97	0.45
2:B:2:VAL:HG13	2:B:27:PHE:CD1	2.52	0.45
2:B:148:PHE:HA	2:B:149:PRO:HA	1.80	0.45
3:C:94:GLN:NE2	7:C:399:HOH:O	2.47	0.44
2:H:22:CYS:HB3	2:H:79:LEU:HB3	1.99	0.44
2:B:216:LYS:HB2	2:B:216:LYS:HE3	1.77	0.44
2:H:12:VAL:O	2:H:113:VAL:HA	2.18	0.44
3:C:60:GLU:HA	3:C:101[B]:ARG:NH2	2.33	0.43
3:C:113:GLU:HA	3:D:28:ILE:HD13	2.00	0.43
2:H:121:PRO:HB2	2:H:144:VAL:HG23	2.00	0.43
1:L:7:PRO:HA	1:L:8:PRO:HD3	1.95	0.43
3:C:11:PRO:HB2	3:C:14:GLU:HG3	2.00	0.43
1:A:122:PRO:HD3	1:A:134:LEU:HD23	2.00	0.43
3:D:54:HIS:HE1	7:D:405:HOH:O	2.02	0.43
3:D:119:VAL:O	7:D:399:HOH:O	2.21	0.43
1:A:34:TRP:CD2	1:A:72:LEU:HB2	2.54	0.42
1:L:168:LYS:HG3	1:L:174:TYR:CZ	2.55	0.42
2:H:127:ALA:HA	2:H:128:PRO:HD3	1.94	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:D:106:CYS:HA	3:D:107:PRO:HD3	1.86	0.42
2:B:8:GLY:O	2:B:18:LEU:HD21	2.20	0.41
2:H:148:PHE:HA	2:H:149:PRO:HA	1.73	0.41
2:H:184:VAL:HG12	2:H:186:VAL:HG13	2.02	0.41
2:B:65:LYS:HB2	2:B:65:LYS:HE3	1.74	0.40
1:A:128:GLN:HG3	7:A:533:HOH:O	2.22	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:H:502:HOH:O	7:A:589:HOH:O[2_455]	2.01	0.19

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	207/214 (97%)	202 (98%)	5 (2%)	0	100	100
1	L	207/214 (97%)	204 (99%)	3 (1%)	0	100	100
2	B	207/224 (92%)	202 (98%)	5 (2%)	0	100	100
2	H	207/224 (92%)	204 (99%)	3 (1%)	0	100	100
3	C	118/132 (89%)	114 (97%)	4 (3%)	0	100	100
3	D	95/132 (72%)	91 (96%)	4 (4%)	0	100	100
All	All	1041/1140 (91%)	1017 (98%)	24 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	172/179 (96%)	169 (98%)	3 (2%)	56	71
1	L	173/179 (97%)	169 (98%)	4 (2%)	45	59
2	B	177/190 (93%)	170 (96%)	7 (4%)	27	35
2	H	177/190 (93%)	170 (96%)	7 (4%)	27	35
3	C	113/125 (90%)	110 (97%)	3 (3%)	40	53
3	D	92/125 (74%)	90 (98%)	2 (2%)	47	61
All	All	904/988 (92%)	878 (97%)	26 (3%)	37	50

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	148	VAL
1	L	157	VAL
1	L	191	ARG
1	L	211	THR
2	H	104	VAL
2	H	107	GLN
2	H	113	VAL
2	H	126	LEU
2	H	144	VAL
2	H	172	LEU
2	H	212	LYS
1	A	10	VAL
1	A	78	GLN
1	A	146	VAL
2	B	64	VAL
2	B	102	LEU
2	B	104	VAL
2	B	137	THR
2	B	140	LEU
2	B	166	HIS
2	B	173	GLN
3	C	24	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	C	74	LEU
3	C	87	MET
3	D	30	ASN
3	D	98	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
3	D	30	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 1 is monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GOL	L	302	-	5,5,5	0.39	0	5,5,5	0.33	0
6	SO4	B	301	-	4,4,4	0.24	0	6,6,6	0.10	0
6	SO4	C	201	-	4,4,4	0.23	0	6,6,6	0.06	0
6	SO4	C	202	-	4,4,4	0.24	0	6,6,6	0.10	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	GOL	L	301	-	5,5,5	0.36	0	5,5,5	0.25	0
4	GOL	A	302	-	5,5,5	0.46	0	5,5,5	0.20	0
4	GOL	H	302	-	5,5,5	0.38	0	5,5,5	0.37	0
6	SO4	H	301	-	4,4,4	0.24	0	6,6,6	0.07	0
4	GOL	A	301	-	5,5,5	0.38	0	5,5,5	0.31	0
6	SO4	D	201	-	4,4,4	0.20	0	6,6,6	0.09	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	L	302	-	-	2/4/4/4	-
4	GOL	L	301	-	-	2/4/4/4	-
4	GOL	A	302	-	-	2/4/4/4	-
4	GOL	H	302	-	-	2/4/4/4	-
4	GOL	A	301	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	L	301	GOL	O1-C1-C2-C3
4	L	302	GOL	O1-C1-C2-C3
4	H	302	GOL	O1-C1-C2-C3
4	A	301	GOL	O1-C1-C2-O2
4	A	302	GOL	O1-C1-C2-O2
4	A	302	GOL	O1-C1-C2-C3
4	L	301	GOL	O1-C1-C2-O2
4	A	301	GOL	O1-C1-C2-C3
4	H	302	GOL	O1-C1-C2-O2
4	L	302	GOL	O1-C1-C2-O2

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	L	301	GOL	1	0
4	A	302	GOL	1	0
4	A	301	GOL	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	209/214 (97%)	-0.35	4 (1%) 66 62	10, 20, 35, 51	0
1	L	209/214 (97%)	-0.36	2 (0%) 79 76	12, 22, 36, 64	0
2	B	210/224 (93%)	-0.61	2 (0%) 79 76	8, 15, 30, 51	1 (0%)
2	H	211/224 (94%)	-0.06	9 (4%) 40 36	10, 23, 56, 92	0
3	C	119/132 (90%)	-0.26	5 (4%) 41 37	8, 19, 37, 60	1 (0%)
3	D	100/132 (75%)	-0.02	12 (12%) 10 8	9, 19, 57, 76	1 (1%)
All	All	1058/1140 (92%)	-0.30	34 (3%) 50 47	8, 20, 46, 92	3 (0%)

All (34) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	D	106	CYS	4.7
2	B	129	SER	4.5
3	C	10	CYS	4.2
3	D	128	VAL	4.1
3	D	127	ILE	4.0
1	L	211	THR	3.7
2	H	130	SER	3.7
3	D	107	PRO	3.4
3	D	104	PRO	3.3
1	A	211	THR	2.9
3	D	31	ARG	2.8
2	H	212	LYS	2.7
3	C	127	ILE	2.7
3	C	105	HIS	2.6
3	D	30	ASN	2.6
3	D	40	SER	2.6
1	A	26	ASN	2.5
3	D	108	ASN	2.5
3	D	20	ARG	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	C	12	ASN	2.5
2	B	216	LYS	2.4
1	L	3	VAL	2.4
2	H	193	THR	2.4
2	H	129	SER	2.3
1	A	67	GLY	2.3
3	C	128	VAL	2.3
3	D	21	THR	2.3
2	H	136	GLY	2.3
3	D	103	PRO	2.2
1	A	68	ASN	2.2
2	H	188	SER	2.1
2	H	208	LYS	2.0
2	H	1	GLN	2.0
2	H	192	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q < 0.9
4	GOL	A	302	6/6	0.76	0.18	27,28,34,40	0
4	GOL	A	301	6/6	0.81	0.27	29,33,41,46	0
4	GOL	L	302	6/6	0.84	0.13	38,47,51,55	0
5	CL	L	303	1/1	0.90	0.11	41,41,41,41	0
4	GOL	H	302	6/6	0.92	0.08	21,30,37,40	0
4	GOL	L	301	6/6	0.93	0.09	25,26,28,29	0
6	SO4	B	301	5/5	0.94	0.18	35,45,48,50	0
6	SO4	H	301	5/5	0.95	0.13	33,48,60,64	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
6	SO4	C	202	5/5	0.98	0.10	24,26,28,31	0
6	SO4	C	201	5/5	0.99	0.05	19,20,22,33	0
6	SO4	D	201	5/5	0.99	0.05	19,20,24,35	0

6.5 Other polymers [i](#)

There are no such residues in this entry.