



Full wwPDB X-ray Structure Validation Report ⓘ

May 23, 2020 – 06:35 pm BST

PDB ID : 3QJM
Title : Structural flexibility of Shank PDZ domain is important for its binding to different ligands
Authors : Lee, J.H.; Park, H.; Park, S.J.; Kim, H.J.; Eom, S.H.
Deposited on : 2011-01-30
Resolution : 2.31 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.31 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5974 (2.34-2.30)
Clashscore	141614	6604 (2.34-2.30)
Ramachandran outliers	138981	6523 (2.34-2.30)
Sidechain outliers	138945	6523 (2.34-2.30)
RSRZ outliers	127900	5855 (2.34-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	115	 11% 61% 22% 15%
1	B	115	 8% 65% 18% 15%
2	C	5	 100%
2	D	5	 100%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 1690 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SH3 and multiple ankyrin repeat domains protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	98	762	488	134	136	4	0	0	0
1	B	98	767	489	137	137	4	0	1	0

- Molecule 2 is a protein called Beta-PIX.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	C	5	40	23	6	11	0	0	0
2	D	5	41	23	6	12	0	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	27	Total	O	0	0
			27	27		
3	B	49	Total	O	0	0
			49	49		
3	C	2	Total	O	0	0
			2	2		
3	D	2	Total	O	0	0
			2	2		

4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	57.62Å 57.62Å 144.04Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.37 – 2.31 26.98 – 2.31	Depositor EDS
% Data completeness (in resolution range)	89.5 (25.37-2.31) 89.3 (26.98-2.31)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	29.93 (at 2.31Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.6.1_357)	Depositor
R, R_{free}	0.235 , 0.267 0.228 , 0.215	Depositor DCC
R_{free} test set	481 reflections (4.77%)	wwPDB-VP
Wilson B-factor (Å ²)	27.0	Xtrriage
Anisotropy	0.699	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 53.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	1690	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.82% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.38	0/773	0.55	0/1040
1	B	0.41	0/780	0.60	0/1046
2	C	0.33	0/39	0.64	0/52
2	D	0.27	0/40	0.49	0/52
All	All	0.39	0/1632	0.58	0/2190

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	762	0	781	20	0
1	B	767	0	789	16	0
2	C	40	0	33	0	0
2	D	41	0	33	0	0
3	A	27	0	0	3	0
3	B	49	0	0	4	0
3	C	2	0	0	0	0
3	D	2	0	0	0	0
All	All	1690	0	1636	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:656:ASP:OD1	1:B:660:LYS:HE2	1.88	0.74
1:A:693:THR:HG22	1:A:696:PHE:H	1.55	0.70
1:B:726:ASN:ND2	1:B:750:MET:H	1.90	0.69
1:A:711:ALA:HB1	1:A:716:LEU:HD12	1.72	0.69
1:A:667:GLN:OE1	1:A:748:THR:HG23	1.94	0.66
1:A:694:PRO:HA	3:A:58:HOH:O	1.96	0.66
1:B:726:ASN:HD21	1:B:750:MET:H	1.45	0.62
1:A:705:VAL:HG23	1:A:712:TRP:HD1	1.72	0.55
1:A:667:GLN:HG2	1:A:748:THR:HG22	1.89	0.55
1:A:679:ARG:HD2	1:A:701:TYR:OH	2.06	0.54
1:B:659:ILE:CD1	1:B:756:VAL:HG22	2.39	0.53
1:A:739:VAL:O	1:A:743:ARG:HG3	2.11	0.51
1:B:657:TYR:HA	1:B:758[A]:ARG:HG2	1.93	0.50
1:B:723:ILE:CG1	1:B:754:VAL:HG23	2.41	0.50
1:A:668:LYS:O	1:A:747:ASN:HB3	2.11	0.49
1:A:705:VAL:CG2	1:A:712:TRP:HD1	2.27	0.47
1:A:693:THR:HB	3:A:76:HOH:O	2.15	0.46
1:B:704:SER:HA	1:B:718:MET:HG2	1.97	0.46
1:A:731:VAL:HB	1:B:691:THR:HG21	1.97	0.46
1:B:723:ILE:HG12	1:B:754:VAL:HG23	1.97	0.45
1:A:724:GLU:HB2	1:A:752:LYS:HB3	1.98	0.45
1:A:681:ALA:C	3:A:63:HOH:O	2.55	0.45
1:A:723:ILE:HB	1:A:752:LYS:HG3	1.99	0.45
1:A:679:ARG:HG2	1:A:680:GLY:N	2.31	0.45
1:A:723:ILE:HB	1:A:752:LYS:CG	2.47	0.44
1:B:721:PHE:O	1:B:753:VAL:HA	2.18	0.44
1:B:667:GLN:NE2	3:B:45:HOH:O	2.52	0.43
1:B:659:ILE:HD12	1:B:756:VAL:HG22	1.99	0.43
1:B:750:MET:HE3	3:B:77:HOH:O	2.21	0.41
1:B:690:PHE:CG	1:B:699:LEU:HD13	2.56	0.41
1:A:691:THR:HA	1:A:692:PRO:HD2	1.91	0.41
1:B:750:MET:CE	3:B:77:HOH:O	2.68	0.41
1:A:667:GLN:HG2	1:A:748:THR:CG2	2.51	0.41
1:B:680:GLY:HA3	1:B:700:GLN:NE2	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	94/115 (82%)	93 (99%)	0	1 (1%)	14	15
1	B	94/115 (82%)	94 (100%)	0	0	100	100
2	C	3/5 (60%)	3 (100%)	0	0	100	100
2	D	3/5 (60%)	3 (100%)	0	0	100	100
All	All	194/240 (81%)	193 (100%)	0	1 (0%)	29	35

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	688	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	81/96 (84%)	75 (93%)	6 (7%)	13	17
1	B	82/96 (85%)	75 (92%)	7 (8%)	10	12
2	C	5/5 (100%)	5 (100%)	0	100	100
2	D	5/5 (100%)	5 (100%)	0	100	100
All	All	173/202 (86%)	160 (92%)	13 (8%)	15	16

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	665	LEU
1	A	667	GLN
1	A	687	ILE
1	A	693	THR
1	A	752	LYS
1	A	758	ARG
1	B	656	ASP
1	B	678	LEU
1	B	699	LEU
1	B	706	ASP
1	B	737	GLN
1	B	758[A]	ARG
1	B	758[B]	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	700	GLN
1	A	726	ASN
1	B	700	GLN
1	B	726	ASN
1	B	740	ASN
1	B	744	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	98/115 (85%)	0.76	13 (13%) 3 4	18, 28, 47, 52	0
1	B	98/115 (85%)	0.46	9 (9%) 9 12	13, 20, 33, 63	0
2	C	5/5 (100%)	-0.11	0 100 100	22, 23, 27, 29	0
2	D	5/5 (100%)	0.86	0 100 100	23, 24, 29, 29	0
All	All	206/240 (85%)	0.60	22 (10%) 6 8	13, 24, 41, 63	0

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	670	ASP	5.2
1	A	687	ILE	4.1
1	A	671	SER	4.0
1	B	689	GLU	3.6
1	A	747	ASN	3.4
1	B	654	GLY	3.4
1	A	689	GLU	3.4
1	B	670	ASP	3.4
1	A	688	GLU	2.9
1	A	691	THR	2.8
1	B	671	SER	2.8
1	A	728	GLN	2.6
1	B	716	LEU	2.6
1	B	736	ARG	2.5
1	A	748	THR	2.4
1	B	758[A]	ARG	2.3
1	A	746	GLY	2.3
1	A	744	GLN	2.2
1	A	740	ASN	2.1
1	A	750	MET	2.1
1	B	751	VAL	2.1
1	B	669	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.