



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 16, 2024 – 10:32 am BST

PDB ID : 8QJ5  
Title : Crystal structure of the Levansucrase beta from *Pseudomonas syringae* pv. *actinidiae*  
Authors : Ferraroni, M.; Peritore, L.  
Deposited on : 2023-09-12  
Resolution : 1.63 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 3.0  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.002 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.38.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.63 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	1015 (1.64-1.64)
Clashscore	180529	1093 (1.64-1.64)
Ramachandran outliers	177936	1077 (1.64-1.64)
Sidechain outliers	177891	1077 (1.64-1.64)
RSRZ outliers	164620	1015 (1.64-1.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	432	 87% 8% 5%
1	B	432	 86% 9% 5%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	GOL	A	506	-	-	X	-

## 2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 7248 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Glutamate 5-kinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	412	3254	2072	542	626	14	0	6	0
1	B	410	3257	2075	542	627	13	0	10	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	expression tag	UNP A0A2V0R8Q9
B	0	SER	-	expression tag	UNP A0A2V0R8Q9

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	P		
2	A	1	5	4	1	0	0

*Continued on next page...*

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	P		
2	B	1	5	4	1	0	0

- Molecule 3 is polyethylene glycol (three-letter code: P4K) (formula:  $C_{30}H_{62}O_{15}$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
3	A	1	21	14	7	0	0

- Molecule 4 is ACETATE ION (three-letter code: ACT) (formula:  $C_2H_3O_2$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is THIOCYANATE ION (three-letter code: SCN) (formula:  $CNS$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
			Total	C	N			S
6	B	1	3	1	1	1	0	0

- Molecule 7 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: C<sub>6</sub>H<sub>13</sub>NO<sub>4</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
7	B	1	12	6	1	4	1	0	0

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	327	Total 327	O 327	0	0
8	B	312	Total 312	O 312	0	0




### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Glutamate 5-kinase

Chain A: 



- Molecule 1: Glutamate 5-kinase

Chain B: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	57.61Å 100.63Å 77.93Å 90.00° 96.06° 90.00°	Depositor
Resolution (Å)	29.30 – 1.63 29.30 – 1.63	Depositor EDS
% Data completeness (in resolution range)	99.7 (29.30-1.63) 99.7 (29.30-1.63)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.44 (at 1.63Å)	Xtrriage
Refinement program	REFMAC 5.8.0267	Depositor
R, $R_{free}$	0.161 , 0.190 0.160 , 0.193	Depositor DCC
$R_{free}$ test set	5431 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	19.6	Xtrriage
Anisotropy	0.421	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.43 , 50.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	7248	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	24.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.48% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MES, SCN, GOL, ACT, PO4, P4K

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.58	0/3364	0.95	3/4589 (0.1%)
1	B	0.56	1/3379 (0.0%)	0.93	4/4609 (0.1%)
All	All	0.57	1/6743 (0.0%)	0.94	7/9198 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	252	GLU	CD-OE1	5.18	1.31	1.25

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	304	ARG	CG-CD-NE	-8.98	92.95	111.80
1	B	348	ARG	CG-CD-NE	6.51	125.46	111.80
1	B	29	ARG	NE-CZ-NH2	-6.25	117.18	120.30
1	A	67	ARG	NE-CZ-NH2	6.02	123.31	120.30
1	B	117[A]	ARG	CG-CD-NE	-5.24	100.80	111.80
1	B	117[B]	ARG	CG-CD-NE	-5.24	100.80	111.80
1	A	67	ARG	NE-CZ-NH1	-5.14	117.73	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3254	0	3096	32	0
1	B	3257	0	3105	40	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
3	A	21	0	0	0	0
4	A	4	0	3	0	0
5	A	24	0	32	5	0
5	B	24	0	32	2	0
6	B	3	0	0	0	0
7	B	12	0	13	3	0
8	A	327	0	0	12	1
8	B	312	0	0	20	1
All	All	7248	0	6281	75	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (75) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:307:TYR:H	5:A:506:GOL:H11	1.22	1.01
5:A:506:GOL:H32	8:A:798:HOH:O	1.64	0.96
1:A:227:ASN:HB3	8:A:880:HOH:O	1.68	0.92
1:B:319[A]:ILE:HD11	8:B:720:HOH:O	1.71	0.88
1:A:307:TYR:H	5:A:506:GOL:C1	1.87	0.85
5:B:507:GOL:H11	8:B:785:HOH:O	1.75	0.84
1:A:307:TYR:N	5:A:506:GOL:H11	1.95	0.80
1:B:319[B]:ILE:HG23	8:B:608:HOH:O	1.81	0.79
1:A:83:LEU:HD23	1:A:118[B]:MET:CE	2.14	0.77
1:A:106:LYS:HE2	1:A:392:TYR:OH	1.86	0.76
1:B:303:GLU:HB2	1:B:319[A]:ILE:HG12	1.70	0.73
1:B:330:GLU:HG3	8:B:868:HOH:O	1.91	0.70
1:A:88:ASP:OD1	1:A:117[B]:ARG:NH2	2.25	0.70
1:B:200[B]:GLU:OE2	8:B:601:HOH:O	2.11	0.68
1:B:319[B]:ILE:CG2	8:B:608:HOH:O	2.41	0.67
1:B:321:HIS:ND1	8:B:603:HOH:O	2.27	0.67
5:B:507:GOL:C1	8:B:785:HOH:O	2.37	0.66
1:B:319[A]:ILE:HD12	1:B:369[A]:SER:HA	1.78	0.65
1:B:117[A]:ARG:CD	1:B:134:ARG:NH2	2.60	0.64
1:A:90:HIS:HD2	1:A:108:ASP:OD1	1.80	0.64
1:A:107:ARG:NE	8:A:601:HOH:O	2.06	0.64
1:B:90:HIS:HD2	1:B:108:ASP:OD1	1.81	0.63

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:321:HIS:CE1	8:B:603:HOH:O	2.51	0.63
1:B:44[A]:GLN:NE2	8:B:604:HOH:O	2.32	0.63
1:B:117[A]:ARG:HD2	1:B:134:ARG:NH2	2.15	0.62
1:B:303:GLU:HB2	1:B:319[A]:ILE:CG1	2.30	0.62
1:A:242:GLU:CG	8:A:912:HOH:O	2.49	0.59
1:A:44[B]:GLN:NE2	1:A:357:LEU:CD2	2.66	0.58
1:A:138:GLU:HG3	8:A:602:HOH:O	2.03	0.58
1:A:306:HIS:HB2	5:A:506:GOL:H12	1.87	0.57
1:B:308:ILE:HD11	1:B:372:VAL:HG12	1.87	0.57
1:A:44[B]:GLN:NE2	1:A:357:LEU:HD21	2.19	0.57
1:A:37:GLU:HG2	1:A:332:PRO:HD3	1.87	0.57
1:A:308[B]:ILE:HD12	1:A:374:PRO:HA	1.87	0.56
1:B:260:HIS:HD2	8:B:774:HOH:O	1.88	0.56
1:A:260:HIS:HE1	8:A:893:HOH:O	1.90	0.55
1:B:117[A]:ARG:HD2	1:B:134:ARG:HH22	1.71	0.55
1:A:60:ILE:C	1:A:60:ILE:HD12	2.27	0.54
1:A:260:HIS:HD2	8:A:786:HOH:O	1.91	0.53
1:B:117[A]:ARG:HE	1:B:134:ARG:NH2	2.06	0.53
1:B:134:ARG:NH2	1:B:138[B]:GLU:HG3	2.24	0.52
1:B:319[A]:ILE:CD1	1:B:369[A]:SER:HA	2.39	0.52
1:A:138:GLU:CG	8:A:602:HOH:O	2.57	0.52
1:B:144[B]:THR:CG2	8:B:745:HOH:O	2.58	0.52
1:B:144[B]:THR:HG21	8:B:745:HOH:O	2.10	0.52
1:A:308[A]:ILE:HD11	1:A:372:VAL:HG12	1.92	0.51
1:B:117[A]:ARG:NE	1:B:134:ARG:NH2	2.59	0.51
1:A:44[B]:GLN:HE22	1:A:357:LEU:CD2	2.24	0.51
1:A:83:LEU:HD23	1:A:118[B]:MET:HE2	1.93	0.50
1:B:60:ILE:HD12	1:B:60:ILE:C	2.32	0.50
1:A:107:ARG:NH2	8:A:601:HOH:O	2.39	0.50
1:A:227:ASN:CB	8:A:880:HOH:O	2.38	0.49
1:B:116:ALA:O	1:B:117[A]:ARG:HD3	2.13	0.49
1:B:155:ASN:HA	7:B:503:MES:H21	1.95	0.49
1:B:387:THR:HG21	1:B:393:ARG:HG3	1.96	0.48
1:B:335:VAL:HG22	1:B:370:HIS:HB2	1.97	0.47
1:B:200[B]:GLU:OE2	8:B:602:HOH:O	2.21	0.47
1:A:95:GLN:O	1:A:107:ARG:NH2	2.48	0.46
1:B:102:ARG:HD3	8:B:759:HOH:O	2.14	0.46
1:B:102:ARG:NH2	8:B:611:HOH:O	2.48	0.46
1:B:155:ASN:CA	7:B:503:MES:H21	2.45	0.46
1:A:134:ARG:NH1	8:A:602:HOH:O	2.49	0.45
1:B:340:GLY:HA3	8:B:713:HOH:O	2.16	0.45

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:90:HIS:HE1	8:B:873:HOH:O	1.99	0.45
1:A:117[B]:ARG:NH1	8:A:603:HOH:O	2.49	0.45
1:B:365:PHE:O	1:B:383:ASP:HB3	2.18	0.44
1:A:37:GLU:CG	1:A:332:PRO:HD3	2.49	0.42
1:B:54:MET:SD	1:B:398:GLU:HG2	2.60	0.42
1:A:365:PHE:O	1:A:383:ASP:HB3	2.20	0.41
1:B:102:ARG:CD	8:B:759:HOH:O	2.67	0.41
1:B:37:GLU:HG2	1:B:332:PRO:HD3	2.01	0.41
1:A:335:VAL:HG22	1:A:370:HIS:HB2	2.01	0.41
1:B:68:GLU:OE2	7:B:503:MES:H32	2.22	0.40
1:B:200[B]:GLU:CD	8:B:602:HOH:O	2.59	0.40
1:A:305:PRO:HA	1:A:317:PHE:O	2.22	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
8:A:881:HOH:O	8:B:620:HOH:O[2_556]	2.19	0.01

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	416/432 (96%)	400 (96%)	15 (4%)	1 (0%)	44	26
1	B	418/432 (97%)	401 (96%)	16 (4%)	1 (0%)	44	26
All	All	834/864 (96%)	801 (96%)	31 (4%)	2 (0%)	44	26

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	168	THR

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	168	THR

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	347/363 (96%)	346 (100%)	1 (0%)	91 86
1	B	348/363 (96%)	346 (99%)	2 (1%)	84 71
All	All	695/726 (96%)	692 (100%)	3 (0%)	89 83

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	215	TRP
1	B	215	TRP
1	B	391	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	90	HIS
1	A	260	HIS
1	A	414	GLN
1	B	90	HIS
1	B	95	GLN
1	B	100	ASN
1	B	260	HIS
1	B	363	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	GOL	A	507	-	5,5,5	0.09	0	5,5,5	0.44	0
5	GOL	B	507	-	5,5,5	0.15	0	5,5,5	0.59	0
2	PO4	B	501	-	4,4,4	0.95	0	6,6,6	0.72	0
5	GOL	B	506	-	5,5,5	0.09	0	5,5,5	0.44	0
5	GOL	A	506	-	5,5,5	0.11	0	5,5,5	0.32	0
6	SCN	B	502	-	1,2,2	3.25	1 (100%)	0,1,1	-	-
5	GOL	A	505	-	5,5,5	0.24	0	5,5,5	0.51	0
5	GOL	B	504	-	5,5,5	0.20	0	5,5,5	0.50	0
2	PO4	A	501	-	4,4,4	1.17	0	6,6,6	0.45	0
4	ACT	A	503	-	3,3,3	0.91	0	3,3,3	0.80	0
5	GOL	A	504	-	5,5,5	0.11	0	5,5,5	0.28	0
5	GOL	B	505	-	5,5,5	0.10	0	5,5,5	0.16	0
7	MES	B	503	-	12,12,12	0.73	0	14,16,16	0.86	0
3	P4K	A	502	-	20,20,44	0.34	0	19,19,43	0.34	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	507	-	-	3/4/4/4	-

*Continued on next page...*



*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	B	507	-	-	4/4/4/4	-
5	GOL	B	506	-	-	1/4/4/4	-
5	GOL	A	506	-	-	4/4/4/4	-
5	GOL	A	505	-	-	0/4/4/4	-
5	GOL	B	504	-	-	0/4/4/4	-
5	GOL	A	504	-	-	0/4/4/4	-
5	GOL	B	505	-	-	0/4/4/4	-
7	MES	B	503	-	-	2/6/14/14	0/1/1/1
3	P4K	A	502	-	-	5/18/18/42	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	B	502	SCN	C-N	3.25	1.26	1.15

There are no bond angle outliers.

There are no chirality outliers.

All (19) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	A	506	GOL	C1-C2-C3-O3
5	B	507	GOL	C1-C2-C3-O3
7	B	503	MES	C8-C7-N4-C3
3	A	502	P4K	O1-C1-C2-O2
5	A	506	GOL	O1-C1-C2-C3
5	A	507	GOL	C1-C2-C3-O3
5	B	507	GOL	O1-C1-C2-C3
5	A	506	GOL	O1-C1-C2-O2
5	A	506	GOL	O2-C2-C3-O3
5	A	507	GOL	O2-C2-C3-O3
3	A	502	P4K	O3-C5-C6-O4
7	B	503	MES	C8-C7-N4-C5
5	B	507	GOL	O1-C1-C2-O2
5	A	507	GOL	O1-C1-C2-O2
5	B	507	GOL	O2-C2-C3-O3
3	A	502	P4K	O4-C7-C8-O5
3	A	502	P4K	C14-C13-O7-C12
5	B	506	GOL	O1-C1-C2-O2
3	A	502	P4K	O2-C3-C4-O3

There are no ring outliers.

3 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	507	GOL	2	0
5	A	506	GOL	5	0
7	B	503	MES	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	412/432 (95%)	-0.47	2 (0%) 87 90	10, 20, 36, 60	6 (1%)
1	B	410/432 (94%)	-0.47	0 100 100	10, 21, 36, 75	10 (2%)
All	All	822/864 (95%)	-0.47	2 (0%) 92 93	10, 20, 36, 75	16 (1%)

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	99	ALA	2.2
1	A	20	ILE	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
7	MES	B	503	12/12	0.80	0.14	44,51,71,74	0
3	P4K	A	502	21/45	0.84	0.15	32,44,57,59	0
4	ACT	A	503	4/4	0.88	0.13	40,42,49,53	0

*Continued on next page...*

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
6	SCN	B	502	3/3	0.89	0.12	29,29,39,66	0
2	PO4	A	501	5/5	0.90	0.11	38,40,48,49	0
5	GOL	B	506	6/6	0.92	0.10	25,37,47,51	0
2	PO4	B	501	5/5	0.92	0.10	36,37,43,46	0
5	GOL	A	507	6/6	0.92	0.11	25,42,45,48	0
5	GOL	A	506	6/6	0.93	0.12	24,45,47,54	0
5	GOL	B	507	6/6	0.93	0.10	22,35,43,46	0
5	GOL	B	504	6/6	0.94	0.08	23,28,29,30	0
5	GOL	B	505	6/6	0.94	0.07	24,27,29,32	0
5	GOL	A	505	6/6	0.95	0.08	21,28,30,31	0
5	GOL	A	504	6/6	0.96	0.07	27,31,32,32	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.