



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 22, 2023 – 03:47 PM EDT

PDB ID : 2QK4
Title : Human glycinamide ribonucleotide synthetase
Authors : Lehtio, L.; Welin, M.; Arrowsmith, C.H.; Berglund, H.; Busam, R.; Collins, R.; Dahlgren, L.G.; Herman, M.D.; Edwards, A.; Flodin, S.; Flores, A.; Graslund, S.; Hammarstrom, M.; Hallberg, B.M.; Holmberg-Schiavone, L.; Johansson, I.; Kallas, A.; Karlberg, T.; Kotenyova, T.; Moche, M.; Nyman, T.; Persson, C.; Sagemark, J.; Stenmark, P.; Sundstrom, M.; Thorsell, A.G.; Tresaugues, L.; van den Berg, S.; Weigelt, J.; Nordlund, P.; Structural Genomics Consortium (SGC)
Deposited on : 2007-07-10
Resolution : 2.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1544 (2.48-2.44)
Clashscore	141614	1613 (2.48-2.44)
Ramachandran outliers	138981	1598 (2.48-2.44)
Sidechain outliers	138945	1598 (2.48-2.44)
RSRZ outliers	127900	1523 (2.48-2.44)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	452	
1	B	452	

CCP4 : 7.0.044 (Gargrove)
 Ideal geometry (proteins) : Engh & Huber (2001)
 Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
 Validation Pipeline (wwPDB-VP) : 2.35

2 Entry composition i

There are 6 unique types of molecules in this entry. The entry contains 6471 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Trifunctional purine biosynthetic protein adenosine-3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	420	3117	1980	529	589	19	0	0	0
1	B	424	3145	1996	535	595	19	0	0	0

There are 46 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-21	MET	-	cloning artifact	UNP P22102
A	-20	HIS	-	cloning artifact	UNP P22102
A	-19	HIS	-	cloning artifact	UNP P22102
A	-18	HIS	-	cloning artifact	UNP P22102
A	-17	HIS	-	cloning artifact	UNP P22102
A	-16	HIS	-	cloning artifact	UNP P22102
A	-15	HIS	-	cloning artifact	UNP P22102
A	-14	SER	-	cloning artifact	UNP P22102
A	-13	SER	-	cloning artifact	UNP P22102
A	-12	GLY	-	cloning artifact	UNP P22102
A	-11	VAL	-	cloning artifact	UNP P22102
A	-10	ASP	-	cloning artifact	UNP P22102
A	-9	LEU	-	cloning artifact	UNP P22102
A	-8	GLY	-	cloning artifact	UNP P22102
A	-7	THR	-	cloning artifact	UNP P22102
A	-6	GLU	-	cloning artifact	UNP P22102
A	-5	ASN	-	cloning artifact	UNP P22102
A	-4	LEU	-	cloning artifact	UNP P22102
A	-3	TYR	-	cloning artifact	UNP P22102
A	-2	PHE	-	cloning artifact	UNP P22102
A	-1	GLN	-	cloning artifact	UNP P22102
A	0	SER	-	cloning artifact	UNP P22102
A	421	ILE	VAL	variant	UNP P22102
B	-21	MET	-	cloning artifact	UNP P22102
B	-20	HIS	-	cloning artifact	UNP P22102

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-19	HIS	-	cloning artifact	UNP P22102
B	-18	HIS	-	cloning artifact	UNP P22102
B	-17	HIS	-	cloning artifact	UNP P22102
B	-16	HIS	-	cloning artifact	UNP P22102
B	-15	HIS	-	cloning artifact	UNP P22102
B	-14	SER	-	cloning artifact	UNP P22102
B	-13	SER	-	cloning artifact	UNP P22102
B	-12	GLY	-	cloning artifact	UNP P22102
B	-11	VAL	-	cloning artifact	UNP P22102
B	-10	ASP	-	cloning artifact	UNP P22102
B	-9	LEU	-	cloning artifact	UNP P22102
B	-8	GLY	-	cloning artifact	UNP P22102
B	-7	THR	-	cloning artifact	UNP P22102
B	-6	GLU	-	cloning artifact	UNP P22102
B	-5	ASN	-	cloning artifact	UNP P22102
B	-4	LEU	-	cloning artifact	UNP P22102
B	-3	TYR	-	cloning artifact	UNP P22102
B	-2	PHE	-	cloning artifact	UNP P22102
B	-1	GLN	-	cloning artifact	UNP P22102
B	0	SER	-	cloning artifact	UNP P22102
B	421	ILE	VAL	variant	UNP P22102

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Cl 1 1	0	0
2	B	1	Total Cl 1 1	0	0

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0

- Molecule 4 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: $C_{10}H_{16}N_5O_{13}P_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C N O P 31 10 5 13 3	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
4	B	1	31	10	5	13	3	0	0

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
5	A	1	6	3	3	0	0
5	A	1	6	3	3	0	0
5	B	1	6	3	3	0	0
5	B	1	6	3	3	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
6	A	49	49	49	0	0
6	B	62	62	62	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Trifunctional purine biosynthetic protein adenosine-3



- Molecule 1: Trifunctional purine biosynthetic protein adenosine-3



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	70.28Å 79.81Å 113.87Å 90.00° 104.07° 90.00°	Depositor
Resolution (Å)	19.98 – 2.45 19.95 – 2.45	Depositor EDS
% Data completeness (in resolution range)	100.0 (19.98-2.45) 99.4 (19.95-2.45)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.37 (at 2.44Å)	Xtrriage
Refinement program	REFMAC 5.3.0032	Depositor
R, R_{free}	0.171 , 0.221 0.166 , 0.214	Depositor DCC
R_{free} test set	2240 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	56.0	Xtrriage
Anisotropy	0.204	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 56.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6471	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.34% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, ATP, SO4, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.68	1/3172 (0.0%)	0.71	1/4296 (0.0%)
1	B	0.67	0/3200	0.72	0/4332
All	All	0.68	1/6372 (0.0%)	0.71	1/8628 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	41	CYS	CB-SG	-7.22	1.70	1.82

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	329	LEU	CA-CB-CG	5.20	127.27	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3117	0	3139	34	0
1	B	3145	0	3173	23	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	5	0	0	0	0
3	B	5	0	0	0	0
4	A	31	0	12	0	0
4	B	31	0	12	0	0
5	A	12	0	16	2	0
5	B	12	0	16	0	0
6	A	49	0	0	2	0
6	B	62	0	0	2	0
All	All	6471	0	6368	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:333:THR:HG21	1:B:396:LEU:HD22	1.62	0.81
1:A:104:GLU:HG3	1:A:290:ASN:ND2	1.96	0.80
1:B:104:GLU:HG3	1:B:290:ASN:ND2	2.02	0.75
1:A:3:ALA:HB1	1:A:68:GLU:HG3	1.72	0.71
1:A:385:ARG:HE	5:A:804:GOL:H2	1.56	0.70
1:B:116:ARG:HD2	1:B:265:GLU:OE2	1.94	0.67
1:A:264:GLN:O	1:A:264:GLN:HG2	1.96	0.64
1:A:196:GLU:OE2	1:A:424:ARG:NH2	2.32	0.62
1:B:75:GLU:OE1	1:B:291:CYS:HB2	2.03	0.59
1:A:172:GLN:OE1	1:A:172:GLN:HA	2.02	0.58
1:A:290:ASN:HD22	1:A:291:CYS:H	1.54	0.54
1:A:117:HIS:NE2	1:A:265:GLU:OE2	2.34	0.54
1:B:107:LYS:O	1:B:111:LYS:HG2	2.08	0.54
1:A:2:ALA:HB1	1:A:28:LYS:HD2	1.90	0.53
1:B:344:TYR:CD1	1:B:345:PRO:HA	2.43	0.52
1:A:44:LYS:NZ	5:A:803:GOL:H2	2.24	0.52
1:A:96:PRO:HG3	1:A:291:CYS:HB3	1.92	0.52
1:A:133:ALA:HB1	1:A:189:ILE:HD11	1.93	0.51
1:A:249:LYS:O	1:A:253:THR:HB	2.12	0.50
1:A:128:THR:CG2	1:A:186:THR:HG22	2.43	0.49
1:B:63:LYS:HE3	1:B:89:ALA:O	2.13	0.49
1:B:142:PHE:O	1:B:144:ALA:N	2.46	0.48
1:A:102:GLN:HA	1:A:105:SER:OG	2.14	0.48
1:B:290:ASN:HD22	1:B:291:CYS:H	1.61	0.47
1:B:42:SER:O	1:B:44:LYS:N	2.49	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:393:ARG:HD2	1:B:394:GLU:OE2	2.15	0.46
1:B:297:GLU:HG3	6:B:818:HOH:O	2.15	0.46
1:A:64:GLU:HG2	1:B:323:THR:HB	1.96	0.46
1:B:223:GLU:HG3	6:B:860:HOH:O	2.15	0.46
1:A:68:GLU:HG2	6:A:847:HOH:O	2.15	0.46
1:B:387:LEU:C	1:B:387:LEU:HD12	2.37	0.45
1:A:404:LYS:HA	1:A:407:LEU:HD12	1.98	0.44
1:A:28:LYS:HB2	1:A:28:LYS:HE2	1.65	0.44
1:A:104:GLU:HG3	1:A:290:ASN:HD21	1.80	0.44
1:B:141:ASP:OD1	1:B:141:ASP:N	2.45	0.44
1:A:9:GLY:HA3	1:A:14:GLU:HG2	2.00	0.44
1:A:357:PHE:CG	1:A:358:PRO:HD3	2.53	0.43
1:A:274:TYR:CD1	1:A:274:TYR:C	2.91	0.43
1:B:33:ALA:HA	1:B:34:PRO:HA	1.88	0.43
1:B:148:LYS:HG2	1:B:158:VAL:HG22	2.00	0.43
1:A:104:GLU:HB2	1:A:290:ASN:HA	2.01	0.43
1:B:333:THR:CG2	1:B:396:LEU:HD22	2.43	0.42
1:A:397:ILE:HG13	1:A:426:ILE:HD13	2.00	0.42
1:A:306:LYS:HB3	1:A:306:LYS:HE2	1.74	0.42
1:B:378:LYS:HA	1:B:378:LYS:HD3	1.75	0.42
1:A:393:ARG:HA	1:A:393:ARG:HD2	1.82	0.42
1:B:42:SER:O	1:B:43:GLU:C	2.56	0.42
1:B:274:TYR:CD1	1:B:274:TYR:C	2.93	0.42
1:A:75:GLU:OE2	1:A:104:GLU:OE1	2.38	0.42
1:A:344:TYR:CD1	1:A:345:PRO:HA	2.54	0.42
1:A:357:PHE:N	1:A:358:PRO:CD	2.83	0.41
1:B:206:GLY:HA2	1:B:269:TYR:O	2.20	0.41
1:A:143:PRO:HA	6:A:849:HOH:O	2.19	0.41
1:A:33:ALA:HA	1:A:34:PRO:HA	1.99	0.41
1:A:393:ARG:CZ	1:A:402:GLU:HG3	2.51	0.41
1:A:230:THR:HB	1:A:344:TYR:CE2	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	414/452 (92%)	397 (96%)	17 (4%)	0	100	100
1	B	418/452 (92%)	398 (95%)	19 (4%)	1 (0%)	47	57
All	All	832/904 (92%)	795 (96%)	36 (4%)	1 (0%)	51	64

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	143	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	326/355 (92%)	317 (97%)	9 (3%)	43	56
1	B	328/355 (92%)	319 (97%)	9 (3%)	44	57
All	All	654/710 (92%)	636 (97%)	18 (3%)	43	56

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	48	THR
1	A	147	VAL
1	A	249	LYS
1	A	252	ASP
1	A	290	ASN
1	A	292	ARG
1	A	323	THR
1	A	393	ARG
1	A	397	ILE
1	B	129	LYS
1	B	132	GLU
1	B	141	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	147	VAL
1	B	185	GLU
1	B	207	LYS
1	B	241	GLN
1	B	290	ASN
1	B	292	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	23	GLN
1	A	85	ASN
1	A	290	ASN
1	B	85	ASN
1	B	102	GLN
1	B	229	ASN
1	B	290	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 2 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	A	701	-	4,4,4	0.20	0	6,6,6	0.55	0
5	GOL	A	803	-	5,5,5	0.33	0	5,5,5	0.61	0
5	GOL	A	804	-	5,5,5	0.51	0	5,5,5	0.65	0
5	GOL	B	801	-	5,5,5	0.37	0	5,5,5	0.35	0
5	GOL	B	802	-	5,5,5	0.42	0	5,5,5	0.12	0
4	ATP	A	501	-	26,33,33	0.88	2 (7%)	31,52,52	1.26	3 (9%)
3	SO4	B	702	-	4,4,4	0.09	0	6,6,6	0.96	0
4	ATP	B	501	-	26,33,33	0.86	1 (3%)	31,52,52	1.35	2 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	804	-	-	0/4/4/4	-
5	GOL	A	803	-	-	3/4/4/4	-
5	GOL	B	801	-	-	0/4/4/4	-
5	GOL	B	802	-	-	0/4/4/4	-
4	ATP	A	501	-	-	6/18/38/38	0/3/3/3
4	ATP	B	501	-	-	4/18/38/38	0/3/3/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	501	ATP	C5-C4	2.54	1.47	1.40
4	B	501	ATP	C5-C4	2.16	1.46	1.40
4	A	501	ATP	C2-N3	2.00	1.35	1.32

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	501	ATP	N3-C2-N1	-3.59	123.07	128.68
4	A	501	ATP	N3-C2-N1	-3.21	123.67	128.68
4	B	501	ATP	C4-C5-N7	-2.96	106.31	109.40
4	A	501	ATP	PA-O3A-PB	-2.71	123.54	132.83
4	A	501	ATP	O3G-PG-O2G	2.03	115.40	107.64

There are no chirality outliers.

All (13) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	501	ATP	PB-O3B-PG-O2G
4	A	501	ATP	C5'-O5'-PA-O1A
4	A	501	ATP	C5'-O5'-PA-O2A
4	B	501	ATP	PB-O3B-PG-O2G
5	A	803	GOL	O2-C2-C3-O3
5	A	803	GOL	C1-C2-C3-O3
5	A	803	GOL	O1-C1-C2-O2
4	B	501	ATP	PG-O3B-PB-O2B
4	B	501	ATP	C4'-C5'-O5'-PA
4	A	501	ATP	PB-O3B-PG-O3G
4	A	501	ATP	C5'-O5'-PA-O3A
4	B	501	ATP	PG-O3B-PB-O1B
4	A	501	ATP	C4'-C5'-O5'-PA


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	803	GOL	1	0
5	A	804	GOL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	420/452 (92%)	0.39	39 (9%) 8 5	34, 50, 68, 97	1 (0%)
1	B	424/452 (93%)	0.50	46 (10%) 5 4	33, 49, 70, 98	1 (0%)
All	All	844/904 (93%)	0.45	85 (10%) 7 4	33, 50, 70, 98	2 (0%)

All (85) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	430	GLN	5.6
1	A	0	SER	5.0
1	B	176	GLN	4.7
1	A	387	LEU	4.2
1	B	183	ALA	4.1
1	B	71	VAL	4.0
1	B	0	SER	4.0
1	A	25	HIS	3.8
1	A	152	LEU	3.6
1	B	387	LEU	3.6
1	B	189	ILE	3.6
1	B	376	ASN	3.6
1	B	338	VAL	3.4
1	A	173	GLU	3.4
1	B	147	VAL	3.4
1	A	162	LYS	3.4
1	B	141	ASP	3.2
1	A	337	VAL	3.2
1	B	273	LEU	3.2
1	B	337	VAL	3.2
1	A	273	LEU	3.2
1	A	1	MET	3.1
1	A	70	VAL	3.1
1	B	1	MET	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	156	LYS	3.0
1	A	71	VAL	3.0
1	A	176	GLN	3.0
1	A	177	GLU	2.9
1	A	388	ALA	2.9
1	A	252	ASP	2.9
1	A	331	ASN	2.9
1	B	233	MET	2.8
1	B	386	VAL	2.8
1	A	234	GLY	2.8
1	B	421	ILE	2.8
1	A	171	VAL	2.7
1	B	258	THR	2.7
1	B	152	LEU	2.7
1	A	338	VAL	2.7
1	B	94	PHE	2.7
1	A	185	GLU	2.7
1	A	58	LEU	2.7
1	B	234	GLY	2.7
1	B	177	GLU	2.6
1	A	323	THR	2.6
1	B	314	GLN	2.6
1	A	137	ILE	2.6
1	B	388	ALA	2.6
1	A	189	ILE	2.6
1	B	178	LYS	2.6
1	A	146	VAL	2.6
1	A	103	LEU	2.6
1	B	335	LEU	2.6
1	B	70	VAL	2.5
1	B	289	PHE	2.5
1	B	336	THR	2.5
1	B	340	ALA	2.5
1	B	339	MET	2.4
1	A	330	GLU	2.4
1	B	146	VAL	2.4
1	B	254	VAL	2.4
1	B	382	HIS	2.3
1	A	245	ASP	2.3
1	A	233	MET	2.3
1	B	413	GLU	2.3
1	B	222	LEU	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	66	LYS	2.3
1	B	375	LYS	2.3
1	B	202	CYS	2.2
1	A	72	VAL	2.2
1	A	335	LEU	2.2
1	A	339	MET	2.2
1	B	389	VAL	2.2
1	B	255	LEU	2.2
1	B	153	ALA	2.2
1	A	118	GLY	2.2
1	B	269	TYR	2.1
1	A	258	THR	2.1
1	B	378	LYS	2.1
1	B	226	GLY	2.1
1	A	142	PHE	2.1
1	B	185	GLU	2.1
1	A	144	ALA	2.0
1	A	282	ASN	2.0
1	B	377	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	ATP	B	501	31/31	0.71	0.28	40,68,108,109	31
4	ATP	A	501	31/31	0.74	0.29	57,71,93,95	31
5	GOL	A	803	6/6	0.75	0.34	77,85,88,88	0
5	GOL	B	801	6/6	0.85	0.17	78,89,90,93	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	GOL	A	804	6/6	0.89	0.17	53,60,66,69	0
5	GOL	B	802	6/6	0.90	0.14	65,75,82,83	0
2	CL	B	601	1/1	0.96	0.12	57,57,57,57	0
3	SO4	A	701	5/5	0.98	0.21	54,54,66,67	0
3	SO4	B	702	5/5	0.98	0.30	55,60,65,67	0
2	CL	A	602	1/1	0.99	0.24	59,59,59,59	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.