



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 2, 2023 – 02:32 pm GMT

PDB ID : 1QNN
Title : Cambialistic superoxide dismutase from *Porphyromonas gingivalis*
Authors : Sugio, S.; Hiraoka, B.Y.; Yamakura, F.
Deposited on : 1999-10-20
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	191	 84% 16% .
1	B	191	 76% 24%
1	C	191	 86% 13% .
1	D	191	 81% 17% ..

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6597 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SUPEROXIDE DISMUTASE.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
1	A	191	Total	C	N	O	S	0	3	0
			1540	990	259	290	1			
1	B	191	Total	C	N	O	S	0	3	0
			1542	991	260	290	1			
1	C	191	Total	C	N	O	S	0	6	0
			1552	999	261	291	1			
1	D	190	Total	C	N	O		0	1	0
			1523	978	258	287				

- Molecule 2 is FE (III) ION (three-letter code: FE) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Fe	0	0
			1	1		
2	B	1	Total	Fe	0	0
			1	1		
2	C	1	Total	Fe	0	0
			1	1		
2	D	1	Total	Fe	0	0
			1	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	114	Total	O	0	0
			114	114		
3	B	100	Total	O	0	0
			100	100		
3	C	143	Total	O	0	0
			143	143		
3	D	79	Total	O	0	0
			79	79		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: SUPEROXIDE DISMUTASE





- Molecule 1: SUPEROXIDE DISMUTASE



- Molecule 1: SUPEROXIDE DISMUTASE



- Molecule 1: SUPEROXIDE DISMUTASE



4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	75.47Å 102.66Å 99.59Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 1.80 45.63 – 1.80	Depositor EDS
% Data completeness (in resolution range)	87.7 (50.00-1.80) 87.7 (45.63-1.80)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.80 (at 1.79Å)	Xtrriage
Refinement program	X-PLOR 3.851	Depositor
R, R_{free}	0.179 , 0.223 0.179 , 0.218	Depositor DCC
R_{free} test set	1927 reflections (3.04%)	wwPDB-VP
Wilson B-factor (Å ²)	21.4	Xtrriage
Anisotropy	0.317	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 50.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.018 for -h,l,k	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6597	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.65% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.37	0/1593	0.58	0/2157
1	B	0.34	0/1599	0.57	0/2165
1	C	0.37	0/1617	0.59	0/2187
1	D	0.34	0/1568	0.53	0/2124
All	All	0.35	0/6377	0.57	0/8633

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1540	0	1490	25	0
1	B	1542	0	1498	25	0
1	C	1552	0	1509	18	0
1	D	1523	0	1469	28	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	114	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	100	0	0	3	0
3	C	143	0	0	3	0
3	D	79	0	0	4	0
All	All	6597	0	5966	96	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (96) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:100:GLN:HE22	1:D:134:LEU:H	1.02	0.95
1:D:184:TRP:O	1:D:188:GLU:HG2	1.72	0.89
1:B:103:SER:OG	1:B:106:LYS:HG3	1.80	0.81
1:B:144:ASN:HD22	1:B:146:VAL:H	1.33	0.74
1:D:100:GLN:HE22	1:D:134:LEU:N	1.85	0.71
1:A:1:MET:HB2	1:A:50:ASN:HD22	1.59	0.67
1:A:1:MET:HB3	1:A:49:GLU:O	1.95	0.67
1:A:144:ASN:HD22	1:A:146:VAL:H	1.44	0.66
1:B:1:MET:HG3	3:B:2001:HOH:O	1.96	0.64
1:D:46:THR:HB	1:D:47[B]:GLU:OE2	1.99	0.63
1:B:144:ASN:ND2	1:B:146:VAL:H	1.96	0.62
1:A:124:VAL:CG1	1:A:156:PHE:HB2	2.29	0.62
1:D:145:PRO:HG2	1:D:152:PRO:HB3	1.81	0.62
1:D:100:GLN:NE2	1:D:134:LEU:H	1.87	0.62
1:A:124:VAL:HG12	1:A:156:PHE:HB2	1.82	0.61
1:D:148:LYS:HE3	3:D:2050:HOH:O	2.01	0.59
1:C:106:LYS:O	1:C:110[A]:GLU:HG3	2.03	0.59
1:C:148[B]:LYS:HE3	3:C:2109:HOH:O	2.01	0.59
1:D:144:ASN:OD1	1:D:146:VAL:HG22	2.04	0.57
1:A:61[A]:GLU:HG2	1:A:62:GLY:N	2.20	0.57
1:C:169:ASN:HD22	1:C:169:ASN:H	1.53	0.55
1:A:183:ASP:HB3	1:A:186:ILE:HD12	1.90	0.54
1:D:77:TYR:O	1:D:80:GLN:HG2	2.08	0.54
1:B:89:PRO:HB3	1:B:184:TRP:CE3	2.42	0.53
1:B:88:ALA:HB1	3:B:2062:HOH:O	2.09	0.53
1:D:89:PRO:HB3	1:D:184:TRP:CE3	2.43	0.53
1:A:89:PRO:HB3	1:A:184:TRP:CE3	2.44	0.52
1:A:144:ASN:HD22	1:A:146:VAL:HG22	1.74	0.52
1:C:144:ASN:OD1	1:C:146:VAL:HG13	2.10	0.52
1:B:104:PHE:CE2	1:B:108:LYS:HD2	2.44	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1:MET:N	3:A:2002:HOH:O	2.43	0.51
1:A:30:LYS:HE3	3:A:2030:HOH:O	2.09	0.51
1:D:124:VAL:CG1	1:D:156:PHE:HB2	2.40	0.51
1:B:124:VAL:CG1	1:B:156:PHE:HB2	2.40	0.51
1:C:82:ARG:HG2	1:C:82:ARG:HH11	1.75	0.51
1:A:104:PHE:CE2	1:A:108:LYS:HD2	2.46	0.51
1:A:6[A]:ILE:HD13	1:A:75:ASN:HA	1.94	0.50
1:A:1:MET:HB2	1:A:50:ASN:ND2	2.27	0.50
1:B:82:ARG:HB2	1:B:183:ASP:HB2	1.93	0.50
1:A:169:ASN:H	1:A:169:ASN:HD22	1.60	0.49
1:C:77:TYR:O	1:C:80:GLN:HG2	2.13	0.49
1:D:80:GLN:HE21	1:D:190:ARG:HH12	1.59	0.49
1:C:101:PHE:O	1:C:106:LYS:HD2	2.12	0.49
1:C:59[B]:LYS:NZ	1:C:59[B]:LYS:HB3	2.28	0.49
1:B:95:GLU:O	1:B:99:LYS:HG3	2.13	0.48
1:B:106:LYS:O	1:B:109:GLU:HG3	2.13	0.48
1:C:19:ILE:HG22	1:C:24:VAL:HG23	1.94	0.48
1:D:21:LYS:O	1:D:25:GLU:HG3	2.13	0.48
1:D:174:HIS:HE1	3:D:2028:HOH:O	1.95	0.48
1:C:146:VAL:HG13	1:C:152:PRO:HG3	1.97	0.47
1:A:1:MET:CG	1:A:2:THR:N	2.77	0.47
1:B:105:GLU:HG2	3:B:2068:HOH:O	2.15	0.47
1:A:54:ASN:O	1:A:58:GLN:HG3	2.15	0.47
1:B:90:LYS:HD2	1:B:91:GLY:H	1.78	0.47
1:D:2:THR:HG22	1:D:2:THR:O	2.15	0.47
1:D:184:TRP:O	1:D:188:GLU:CG	2.54	0.47
1:D:82:ARG:HB2	1:D:85:LYS:HG3	1.97	0.47
1:D:82:ARG:HG2	1:D:82:ARG:HH11	1.80	0.46
1:B:169:ASN:H	1:B:169:ASN:HD22	1.64	0.46
1:D:160:GLU:HG3	3:D:2057:HOH:O	2.16	0.46
1:C:73:ASN:HB3	1:C:123:TRP:CZ2	2.51	0.45
1:B:46:THR:O	1:B:49[B]:GLU:OE1	2.35	0.45
1:A:124:VAL:O	1:A:124:VAL:HG13	2.17	0.45
1:B:113:THR:HA	1:B:116[C]:THR:HG22	1.98	0.45
1:C:21:LYS:HE3	3:C:2022:HOH:O	2.15	0.45
1:C:106:LYS:HG3	3:C:2088:HOH:O	2.16	0.45
1:D:33:LYS:HE3	3:D:2018:HOH:O	2.15	0.45
1:C:81:PHE:O	1:C:82:ARG:HG2	2.16	0.44
1:B:46:THR:O	1:B:49[B]:GLU:HG2	2.17	0.44
1:C:169:ASN:H	1:C:169:ASN:ND2	2.15	0.44
1:A:1:MET:HE3	1:A:50:ASN:HA	1.99	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:29:GLY:O	1:B:33[B]:LYS:HE3	2.18	0.44
1:D:39:LEU:O	1:D:43:ILE:HG23	2.18	0.44
1:A:128:SER:HA	1:A:133:LYS:O	2.17	0.44
1:C:48:PHE:HB3	1:C:56:ILE:HG12	2.00	0.43
1:B:116[C]:THR:HG23	1:B:117:THR:HG23	2.00	0.43
1:B:129:ASP:OD2	1:B:133:LYS:HB3	2.19	0.43
1:D:73:ASN:HB3	1:D:123:TRP:CZ2	2.54	0.42
1:B:85:LYS:HE3	1:B:183:ASP:OD1	2.19	0.42
1:A:73:ASN:HB3	1:A:123:TRP:CZ2	2.55	0.42
1:C:122:GLY:O	1:C:158:VAL:HG22	2.20	0.42
1:D:85:LYS:HE2	1:D:183:ASP:OD1	2.20	0.42
1:A:144:ASN:ND2	1:A:146:VAL:H	2.15	0.41
1:B:26:PHE:O	1:B:30:LYS:HB2	2.20	0.41
1:B:73:ASN:HB3	1:B:123:TRP:CZ2	2.55	0.41
1:B:112:ASN:O	1:B:116[B]:THR:HG23	2.20	0.41
1:D:18:VAL:HG22	1:D:177:ASP:HB3	2.03	0.41
1:A:144:ASN:ND2	1:A:146:VAL:HG22	2.36	0.40
1:C:89:PRO:HB3	1:C:184:TRP:CE3	2.57	0.40
1:B:126:LEU:HG	1:B:153:LEU:HD12	2.04	0.40
1:D:188:GLU:HG2	1:D:188:GLU:H	1.73	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	192/191 (100%)	187 (97%)	5 (3%)	0	100	100
1	B	193/191 (101%)	180 (93%)	13 (7%)	0	100	100
1	C	195/191 (102%)	190 (97%)	5 (3%)	0	100	100
1	D	189/191 (99%)	177 (94%)	12 (6%)	0	100	100
All	All	769/764 (101%)	734 (95%)	35 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	163/160 (102%)	159 (98%)	4 (2%)	47	34
1	B	164/160 (102%)	157 (96%)	7 (4%)	29	14
1	C	166/160 (104%)	163 (98%)	3 (2%)	59	48
1	D	160/160 (100%)	155 (97%)	5 (3%)	40	25
All	All	653/640 (102%)	634 (97%)	19 (3%)	41	29

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	47	GLU
1	A	147	ARG
1	A	148	LYS
1	A	169	ASN
1	B	6	ILE
1	B	7	SER
1	B	118	LEU
1	B	128	SER
1	B	131	ASN
1	B	148	LYS
1	B	164	TYR
1	C	54	ASN
1	C	95	GLU
1	C	169	ASN
1	D	2	THR
1	D	7	SER
1	D	95	GLU
1	D	164	TYR
1	D	188	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	50	ASN
1	A	144	ASN
1	A	169	ASN
1	B	131	ASN
1	B	144	ASN
1	B	169	ASN
1	C	3	HIS
1	C	38	ASN
1	C	67	ASN
1	C	169	ASN
1	D	54	ASN
1	D	80	GLN
1	D	100	GLN
1	D	174	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	191/191 (100%)	-0.54	1 (0%) 91 89	12, 24, 39, 56	0
1	B	191/191 (100%)	-0.34	1 (0%) 91 89	14, 28, 49, 57	0
1	C	191/191 (100%)	-0.50	0 100 100	13, 22, 37, 41	0
1	D	190/191 (99%)	-0.09	4 (2%) 63 59	16, 36, 53, 63	0
All	All	763/764 (99%)	-0.37	6 (0%) 86 84	12, 27, 48, 63	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1	MET	4.9
1	A	1	MET	4.0
1	D	50	ASN	3.7
1	D	45	GLY	2.5
1	D	51	ALA	2.2
1	D	90	LYS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	FE	D	201	1/1	0.99	0.06	27,27,27,27	0
2	FE	B	201	1/1	1.00	0.06	20,20,20,20	0
2	FE	C	201	1/1	1.00	0.05	18,18,18,18	0
2	FE	A	201	1/1	1.00	0.04	20,20,20,20	0

6.5 Other polymers [i](#)

There are no such residues in this entry.