



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 10, 2024 – 10:49 PM EST

PDB ID : 2QNA
Title : Crystal structure of human Importin-beta (127-876) in complex with the IBB-domain of Snurportin1 (1-65)
Authors : Wohlwend, D.; Strasser, A.; Dickmanns, A.; Ficner, R.
Deposited on : 2007-07-18
Resolution : 2.84 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.84 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1031 (2.86-2.82)
Clashscore	141614	1078 (2.86-2.82)
Ramachandran outliers	138981	1050 (2.86-2.82)
Sidechain outliers	138945	1051 (2.86-2.82)
RSRZ outliers	127900	1019 (2.86-2.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	762	 6% 86% 11% ..
2	B	66	 3% 39% 6% 55%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 6090 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Importin subunit beta-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	744	5769	3639	957	1129	44	0	0	0

There are 13 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	126	SER	-	cloning artifact	UNP Q14974
A	876	GLY	ALA	cloning artifact	UNP Q14974
A	877	LEU	-	cloning artifact	UNP Q14974
A	878	THR	-	cloning artifact	UNP Q14974
A	879	ARG	-	cloning artifact	UNP Q14974
A	880	VAL	-	cloning artifact	UNP Q14974
A	881	ASP	-	cloning artifact	UNP Q14974
A	882	SER	-	cloning artifact	UNP Q14974
A	883	SER	-	cloning artifact	UNP Q14974
A	884	GLY	-	cloning artifact	UNP Q14974
A	885	ARG	-	cloning artifact	UNP Q14974
A	886	ILE	-	cloning artifact	UNP Q14974
A	887	VAL	-	cloning artifact	UNP Q14974

- Molecule 2 is a protein called Snurportin-1.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	30	261	158	58	45	5	0	0

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	46	Total	O	0	0
			46	46		
4	B	4	Total	O	0	0
			4	4		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Importin subunit beta-1



- Molecule 2: Snurportin-1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	64.22Å 79.07Å 208.46Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	15.00 – 2.84 14.99 – 2.84	Depositor EDS
% Data completeness (in resolution range)	100.0 (15.00-2.84) 100.0 (14.99-2.84)	Depositor EDS
R_{merge}	0.19	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.59 (at 2.86Å)	Xtrriage
Refinement program	REFMAC 5.3.0008	Depositor
R, R_{free}	0.233 , 0.282 0.230 , 0.278	Depositor DCC
R_{free} test set	1304 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	32.5	Xtrriage
Anisotropy	0.085	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 44.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	6090	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.51% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.31	0/5863	0.46	0/7957
2	B	0.34	0/262	0.51	0/346
All	All	0.31	0/6125	0.46	0/8303

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5769	0	5750	36	0
2	B	261	0	276	2	0
3	A	10	0	0	0	0
4	A	46	0	0	0	0
4	B	4	0	0	0	0
All	All	6090	0	6026	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:537:LYS:HG3	1:A:587:THR:HG22	1.65	0.78
1:A:533:MET:HB3	1:A:587:THR:HG21	1.72	0.70
1:A:478:ALA:HB2	1:A:535:ILE:HD13	1.73	0.69
1:A:265:ILE:HA	1:A:268:MET:HE3	1.76	0.68
1:A:377:ASN:HD22	1:A:379:ASP:H	1.42	0.66
1:A:419:ASP:O	1:A:425:ARG:HD2	1.97	0.64
1:A:536:VAL:HG13	1:A:547:VAL:HG13	1.84	0.60
1:A:253:GLU:HG3	1:A:255:TYR:HB3	1.88	0.56
1:A:206:LYS:HD3	1:A:207:ALA:H	1.72	0.55
1:A:848:THR:HA	1:A:851:ARG:HH21	1.73	0.53
1:A:551:THR:HG21	1:A:591:VAL:HG11	1.91	0.53
1:A:777:LYS:HG3	1:A:783:VAL:HG13	1.92	0.52
1:A:665:GLN:HE21	1:A:665:GLN:HA	1.75	0.51
1:A:151:LEU:HD13	1:A:194:ALA:HB2	1.93	0.50
1:A:559:GLN:HG3	1:A:614:MET:HG3	1.93	0.49
1:A:383:ARG:O	1:A:387:VAL:HG23	2.12	0.49
1:A:706:HIS:HD2	1:A:708:SER:H	1.61	0.47
1:A:582:SER:HB3	2:B:43:ARG:HD2	1.96	0.46
1:A:518:ASP:O	1:A:521:GLN:HG2	2.16	0.45
1:A:713:ILE:O	1:A:716:VAL:HG12	2.17	0.45
1:A:592:LEU:HD21	1:A:635:LEU:HD12	1.99	0.45
1:A:511:LEU:O	1:A:514:THR:HB	2.16	0.44
1:A:659:LYS:HG3	1:A:695:LEU:HD22	2.00	0.44
1:A:343:ASN:HB2	1:A:344:PRO:HD2	2.01	0.42
1:A:628:ALA:O	1:A:632:VAL:HG23	2.20	0.42
2:B:45:ARG:O	2:B:49:GLU:HG2	2.19	0.42
1:A:403:LYS:O	1:A:407:ILE:HG12	2.18	0.42
1:A:507:VAL:HG13	1:A:550:THR:HG21	2.00	0.42
1:A:198:LEU:O	1:A:202:LEU:HG	2.21	0.41
1:A:253:GLU:CG	1:A:255:TYR:HB3	2.51	0.41
1:A:319:LEU:HD21	1:A:363:ILE:HA	2.03	0.41
1:A:333:GLN:O	1:A:381:ARG:NH2	2.54	0.41
1:A:608:MET:HE1	1:A:632:VAL:HG13	2.03	0.41
1:A:434:ARG:HG3	1:A:472:TRP:HH2	1.86	0.41
1:A:562:LEU:HA	1:A:581:GLN:HE21	1.85	0.41
1:A:516:ARG:HA	1:A:517:PRO:HD3	1.84	0.40
1:A:665:GLN:HA	1:A:665:GLN:NE2	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	740/762 (97%)	722 (98%)	18 (2%)	0	100	100
2	B	28/66 (42%)	27 (96%)	1 (4%)	0	100	100
All	All	768/828 (93%)	749 (98%)	19 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	638/656 (97%)	607 (95%)	31 (5%)	25	47
2	B	28/60 (47%)	27 (96%)	1 (4%)	35	60
All	All	666/716 (93%)	634 (95%)	32 (5%)	25	49

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	146	MET
1	A	148	GLU
1	A	171	ASN
1	A	206	LYS
1	A	234	ARG
1	A	278	GLN
1	A	296	GLU
1	A	338	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	340	ASP
1	A	341	ASP
1	A	343	ASN
1	A	368	LEU
1	A	377	ASN
1	A	413	LEU
1	A	434	ARG
1	A	463	GLU
1	A	489	ASP
1	A	492	GLU
1	A	497	TYR
1	A	524	LEU
1	A	537	LYS
1	A	558	LEU
1	A	583	LEU
1	A	589	GLN
1	A	612	LEU
1	A	625	GLN
1	A	673	LEU
1	A	758	LEU
1	A	811	THR
1	A	821	LEU
1	A	856	ASN
2	B	48	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	139	ASN
1	A	159	GLN
1	A	208	ASN
1	A	240	ASN
1	A	285	ASN
1	A	320	GLN
1	A	328	GLN
1	A	377	ASN
1	A	445	ASN
1	A	522	ASN
1	A	559	GLN
1	A	581	GLN
1	A	665	GLN
1	A	706	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	712	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	A	1	-	4,4,4	0.14	0	6,6,6	0.04	0
3	SO4	A	2	-	4,4,4	0.14	0	6,6,6	0.08	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	744/762 (97%)	0.02	45 (6%) 21 15	18, 28, 49, 58	0
2	B	30/66 (45%)	0.44	2 (6%) 17 12	30, 32, 41, 41	1 (3%)
All	All	774/828 (93%)	0.04	47 (6%) 21 15	18, 29, 49, 58	1 (0%)

All (47) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	127	PRO	6.8
1	A	750	SER	5.8
1	A	340	ASP	5.6
1	A	164	GLU	5.3
1	A	339	ASP	5.2
1	A	489	ASP	5.0
1	A	336	ASN	4.8
1	A	578	ASN	4.7
1	A	752	TYR	4.5
1	A	781	GLU	4.4
2	B	66	ALA	3.9
1	A	165	GLN	3.8
1	A	337	ASP	3.8
1	A	779	ASP	3.8
1	A	140	PRO	3.7
1	A	255	TYR	3.7
1	A	880	VAL	3.2
1	A	879	ARG	3.1
1	A	341	ASP	3.0
1	A	126	SER	2.9
1	A	878	THR	2.9
1	A	446	ASP	2.8
1	A	271	ASP	2.8
1	A	167	GLN	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	37	SER	2.7
1	A	305	ARG	2.7
1	A	856	ASN	2.6
1	A	617	SER	2.6
1	A	162	ASP	2.6
1	A	748	ASP	2.5
1	A	619	ALA	2.4
1	A	216	HIS	2.4
1	A	780	GLN	2.4
1	A	490	ASP	2.3
1	A	620	GLY	2.3
1	A	210	ASP	2.3
1	A	577	PHE	2.2
1	A	782	ASN	2.2
1	A	229	PRO	2.2
1	A	212	GLU	2.1
1	A	254	THR	2.1
1	A	334	ASP	2.1
1	A	335	GLU	2.1
1	A	213	SER	2.1
1	A	338	ASP	2.1
1	A	161	ILE	2.0
1	A	163	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
-----	------	-------	-----	-------	------	-----	-----------------------------	-------

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	SO4	A	1	5/5	0.92	0.31	60,61,61,61	0
3	SO4	A	2	5/5	0.92	0.28	66,67,67,67	0

6.5 Other polymers [i](#)

There are no such residues in this entry.