



# Full wwPDB X-ray Structure Validation Report ⓘ

May 26, 2020 – 11:13 pm BST

PDB ID : 6QNP  
Title : CLATHRIN HEAVY CHAIN N-TERMINAL DOMAIN BOUND TO GTSE1  
LIDL MOTIF  
Authors : Porfetye, A.T.; Lin, Y.; Vetter, I.R.  
Deposited on : 2019-02-11  
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	364	
1	B	364	
1	C	364	
1	D	364	
2	H	67	
2	I	67	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	J	67	
2	K	67	

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 12084 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Clathrin heavy chain 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	361	Total 2819	C 1792	N 484	O 525	S 18	0	0	0
1	B	360	Total 2810	C 1787	N 482	O 523	S 18	0	0	0
1	C	356	Total 2784	C 1771	N 479	O 516	S 18	0	0	0
1	D	354	Total 2770	C 1763	N 476	O 513	S 18	0	0	0

- Molecule 2 is a protein called G2 and S phase-expressed protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	H	20	Total 148	C 94	N 24	O 29	S 1	0	0	0
2	I	20	Total 148	C 94	N 24	O 29	S 1	0	0	0
2	J	24	Total 179	C 118	N 26	O 34	S 1	0	0	0
2	K	24	Total 180	C 118	N 26	O 35	S 1	0	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	86	Total 86	O 86	0	0
3	B	89	Total 89	O 89	0	0
3	C	23	Total 23	O 23	0	0
3	D	35	Total 35	O 35	0	0

*Continued on next page...*


*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>	<b>ZeroOcc</b>	<b>AltConf</b>
3	H	6	Total O 6 6	0	0
3	I	2	Total O 2 2	0	0
3	J	1	Total O 1 1	0	0
3	K	4	Total O 4 4	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Clathrin heavy chain 1




- Molecule 1: Clathrin heavy chain 1




- Molecule 1: Clathrin heavy chain 1




- Molecule 1: Clathrin heavy chain 1







• Molecule 2: G2 and S phase-expressed protein 1




• Molecule 2: G2 and S phase-expressed protein 1



• Molecule 2: G2 and S phase-expressed protein 1



• Molecule 2: G2 and S phase-expressed protein 1



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	95.09Å 90.15Å 100.23Å 90.00° 110.12° 90.00°	Depositor
Resolution (Å)	49.19 – 2.70 49.19 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.0 (49.19-2.70) 99.3 (49.19-2.70)	Depositor EDS
$R_{merge}$	0.14	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.38 (at 2.69Å)	Xtrriage
Refinement program	PHENIX 1.14_3260	Depositor
R, $R_{free}$	0.241 , 0.291 0.241 , 0.292	Depositor DCC
$R_{free}$ test set	2181 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	64.1	Xtrriage
Anisotropy	0.416	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 47.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	12084	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	71.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 74.76 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.4371e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality i

### 5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.42	1/2877 (0.0%)	0.65	2/3901 (0.1%)
1	B	0.36	0/2868	0.62	1/3889 (0.0%)
1	C	0.42	0/2842	0.77	9/3854 (0.2%)
1	D	0.47	3/2828 (0.1%)	0.71	1/3835 (0.0%)
2	H	0.30	0/148	0.62	0/199
2	I	0.28	0/148	0.63	0/199
2	J	0.41	0/181	0.89	1/247 (0.4%)
2	K	0.36	0/182	0.81	0/247
All	All	0.42	4/12074 (0.0%)	0.69	14/16371 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	C	0	1
2	H	0	1
All	All	0	3

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	223	GLN	CG-CD	7.49	1.68	1.51
1	D	336	PRO	N-CD	-5.99	1.39	1.47
1	D	294	TYR	CD2-CE2	-5.85	1.30	1.39
1	D	186	VAL	CB-CG2	-5.67	1.41	1.52

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	251	PHE	CB-CG-CD2	7.57	126.10	120.80

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	J	664	LEU	CB-CG-CD2	7.45	123.67	111.00
1	C	251	PHE	CB-CG-CD1	-7.12	115.82	120.80
1	C	221	ARG	CA-CB-CG	6.74	128.23	113.40
1	C	292	CYS	CA-CB-SG	-6.62	102.08	114.00
1	A	223	GLN	CG-CD-NE2	-6.42	101.28	116.70
1	A	223	GLN	CG-CD-OE1	6.03	133.66	121.60
1	C	357	LEU	CA-CB-CG	5.91	128.90	115.30
1	C	338	ILE	CG1-CB-CG2	-5.80	98.65	111.40
1	D	222	GLY	N-CA-C	-5.75	98.74	113.10
1	C	88	LEU	CA-CB-CG	5.69	128.38	115.30
1	B	349	LEU	CA-CB-CG	5.49	127.92	115.30
1	C	342	LEU	CA-CB-CG	5.29	127.48	115.30
1	C	350	ARG	NE-CZ-NH2	5.00	122.80	120.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	45	GLY	Peptide
1	C	301	GLU	Peptide
2	H	707	SER	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2819	0	2827	35	0
1	B	2810	0	2819	15	0
1	C	2784	0	2796	34	0
1	D	2770	0	2783	29	0
2	H	148	0	157	2	0
2	I	148	0	157	1	0
2	J	179	0	185	2	0
2	K	180	0	186	2	0
3	A	86	0	0	3	0
3	B	89	0	0	1	0
3	C	23	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	D	35	0	0	0	0
3	H	6	0	0	0	0
3	I	2	0	0	0	0
3	J	1	0	0	0	0
3	K	4	0	0	0	0
All	All	12084	0	11910	114	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (114) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:173:GLN:HB2	1:D:174:GLN:OE1	1.44	1.17
1:A:223:GLN:HE21	1:A:223:GLN:HA	1.26	0.97
1:A:223:GLN:HE21	1:A:223:GLN:CA	1.77	0.96
1:A:223:GLN:HA	1:A:223:GLN:NE2	1.81	0.93
1:C:335:ILE:CG1	1:C:357:LEU:HD13	2.11	0.81
1:D:173:GLN:CB	1:D:174:GLN:OE1	2.30	0.79
1:B:44:VAL:O	1:B:47:GLN:HB2	1.89	0.72
1:D:261:PRO:HB3	1:D:275:LEU:HD11	1.78	0.65
1:C:335:ILE:HD11	1:C:357:LEU:HD13	1.78	0.64
1:B:275:LEU:HB3	1:B:283:HIS:HB2	1.80	0.63
1:D:294:TYR:HE2	1:D:296:ASN:HB2	1.62	0.63
1:D:176:ARG:NH1	1:D:221:ARG:O	2.32	0.62
1:C:14:GLN:NE2	1:C:321:LYS:O	2.34	0.60
1:B:83:LYS:HB3	1:B:88:LEU:HD23	1.83	0.60
1:D:335:ILE:HG13	1:D:336:PRO:HD3	1.82	0.60
1:C:335:ILE:CD1	1:C:357:LEU:HD13	2.30	0.60
1:C:335:ILE:HD11	1:C:357:LEU:HB2	1.85	0.57
1:C:220:VAL:HG22	1:C:222:GLY:H	1.69	0.57
1:B:286:ASP:OD1	1:B:354:ARG:NH2	2.31	0.57
1:C:338:ILE:HA	1:C:342:LEU:HD23	1.86	0.56
1:C:228:LEU:HD13	1:C:285:TYR:CD1	2.41	0.56
1:A:223:GLN:NE2	1:A:223:GLN:CA	2.51	0.55
1:A:205:LYS:NZ	1:A:210:ALA:O	2.40	0.54
1:C:318:VAL:HG22	1:C:324:VAL:HG22	1.90	0.54
1:C:303:ILE:HA	1:C:319:ASN:HA	1.90	0.54
1:D:284:LEU:O	1:D:293:ILE:N	2.40	0.53
1:C:144:ARG:NH2	1:C:148:LEU:O	2.41	0.52
1:C:335:ILE:HG12	1:C:357:LEU:HD13	1.89	0.52

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:281:TYR:HA	1:A:297:ARG:HA	1.91	0.52
1:C:176:ARG:NH1	1:C:222:GLY:O	2.43	0.52
1:C:261:PRO:HA	1:C:277:THR:HA	1.91	0.52
1:D:338:ILE:HD11	1:D:344:ASN:HB3	1.92	0.52
1:C:157:ARG:NH2	1:C:264:MET:O	2.38	0.51
1:D:279:TYR:HB3	1:D:297:ARG:HH21	1.75	0.51
1:B:256:ALA:HB1	1:B:259:ASP:HB2	1.92	0.50
1:A:89:GLN:HB2	2:H:689:LEU:HD12	1.94	0.50
1:A:261:PRO:HA	1:A:277:THR:HA	1.94	0.50
1:C:319:ASN:HD21	1:C:323:GLN:HB2	1.76	0.50
1:C:80:ILE:HB	1:C:93:ILE:HD11	1.94	0.50
1:A:28:SER:O	1:A:41:ARG:NH1	2.45	0.50
1:D:261:PRO:HA	1:D:277:THR:HA	1.93	0.50
1:D:98:LYS:NZ	2:K:666:ASP:OD1	2.45	0.49
1:A:223:GLN:C	1:A:223:GLN:HE21	2.14	0.49
1:A:13:LEU:HD11	1:A:58:PRO:HB3	1.94	0.49
1:C:103:THR:O	2:J:662:GLN:NE2	2.42	0.48
1:C:88:LEU:HD11	1:C:108:VAL:HG21	1.94	0.48
1:D:298:ILE:HD13	1:D:325:LEU:HD11	1.96	0.48
1:D:122:VAL:HG21	1:D:167:LEU:HD11	1.94	0.48
1:A:32:MET:HG3	1:A:38:ILE:HG12	1.95	0.48
1:B:65:PRO:HD2	2:I:691:ASP:HB3	1.96	0.48
1:C:205:LYS:NZ	1:C:210:ALA:O	2.47	0.48
1:C:200:SER:OG	1:C:201:PHE:N	2.47	0.47
1:D:276:ILE:HA	1:D:281:TYR:O	2.14	0.47
1:C:221:ARG:NH2	1:C:257:GLN:O	2.48	0.47
1:A:51:VAL:HG22	1:A:63:ARG:HG2	1.95	0.47
1:A:275:LEU:HB3	1:A:283:HIS:HB2	1.97	0.46
1:A:221:ARG:NH1	1:A:259:ASP:O	2.40	0.46
1:B:52:ILE:HB	1:B:62:ILE:HG13	1.98	0.46
1:A:83:LYS:HB2	1:A:88:LEU:HD23	1.98	0.46
1:B:263:ALA:HB3	1:B:276:ILE:HB	1.97	0.46
1:D:352:ALA:HA	1:D:357:LEU:HB2	1.98	0.46
1:A:14:GLN:NE2	3:A:421:HOH:O	2.49	0.46
1:A:72:ILE:HG13	1:A:81:ALA:HB3	1.97	0.46
1:C:352:ALA:HA	1:C:357:LEU:HD12	1.99	0.45
1:D:232:GLU:HB2	1:D:243:PHE:HB3	1.99	0.45
1:A:356:ASN:ND2	3:A:414:HOH:O	2.46	0.45
1:D:318:VAL:HG22	1:D:324:VAL:HG22	1.99	0.45
1:A:276:ILE:HA	1:A:281:TYR:O	2.17	0.45
1:A:319:ASN:HD21	1:A:323:GLN:HB3	1.83	0.44

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:172:ALA:HA	1:C:177:VAL:HA	2.00	0.44
1:D:299:SER:OG	1:D:301:GLU:O	2.34	0.44
1:A:293:ILE:HA	1:A:342:LEU:HD11	2.00	0.44
1:A:245:LYS:HE2	1:A:245:LYS:HB2	1.80	0.44
1:A:36:LYS:NZ	3:A:422:HOH:O	2.50	0.44
1:B:336:PRO:HA	1:B:339:THR:HG22	1.99	0.44
1:D:303:ILE:HG12	1:D:319:ASN:HB3	2.00	0.44
1:C:335:ILE:HG13	1:C:357:LEU:HD13	1.97	0.44
1:D:288:GLU:OE1	1:D:354:ARG:NH1	2.51	0.44
1:C:350:ARG:O	1:C:354:ARG:HB2	2.18	0.43
1:D:184:TYR:CE1	1:D:186:VAL:HG12	2.52	0.43
1:A:16:GLN:HE22	1:A:22:PRO:HD3	1.82	0.43
1:A:184:TYR:CE1	1:A:186:VAL:HG12	2.54	0.43
1:C:228:LEU:HD13	1:C:285:TYR:CE1	2.54	0.43
1:C:100:LYS:NZ	1:C:134:GLY:O	2.52	0.43
1:D:153:ILE:HA	1:D:169:GLY:HA2	2.01	0.43
1:C:299:SER:OG	1:C:301:GLU:O	2.37	0.43
1:A:318:VAL:HG22	1:A:324:VAL:HG22	2.01	0.43
1:A:83:LYS:CB	1:A:88:LEU:HD23	2.49	0.43
1:C:196:GLY:HA2	1:C:220:VAL:HB	2.01	0.43
1:D:335:ILE:HA	1:D:338:ILE:HG22	2.01	0.43
1:A:84:ALA:O	1:A:87:THR:OG1	2.34	0.42
1:D:72:ILE:HG13	1:D:81:ALA:HB3	2.01	0.42
1:D:246:LYS:HE2	1:D:289:THR:HA	2.00	0.42
1:A:96:LYS:HA	2:H:692:LEU:HD13	2.01	0.42
1:A:338:ILE:HA	1:A:342:LEU:HB2	2.01	0.42
1:B:8:ARG:NH1	1:B:330:GLU:OE2	2.50	0.42
1:D:318:VAL:HA	1:D:323:GLN:O	2.19	0.42
1:A:157:ARG:NH2	1:A:264:MET:O	2.47	0.42
1:B:227:LYS:NZ	3:B:417:HOH:O	2.49	0.41
1:A:276:ILE:HG12	1:A:282:ILE:HD12	2.02	0.41
1:B:121:LEU:HB2	1:B:128:TYR:HB2	2.02	0.41
1:C:226:GLY:HA3	1:C:250:VAL:HB	2.01	0.41
1:A:53:ILE:HD13	1:A:61:PRO:HB3	2.02	0.41
1:A:280:GLY:O	1:A:298:ILE:N	2.54	0.41
1:D:283:HIS:HB3	1:D:285:TYR:HE2	1.84	0.41
1:C:62:ILE:HD13	1:C:78:LYS:HZ1	1.86	0.41
1:B:236:PRO:HA	1:B:237:PRO:HD3	1.96	0.40
1:B:25:ILE:HG23	1:B:30:LEU:HD13	2.03	0.40
1:D:64:ARG:HB2	2:K:670:ILE:HD11	2.03	0.40
1:B:244:PRO:O	1:B:246:LYS:NZ	2.46	0.40

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:166:LEU:HD11	1:C:181:MET:HE3	2.03	0.40
1:C:231:ILE:HG23	1:C:245:LYS:HG2	2.02	0.40
1:D:256:ALA:HB1	1:D:259:ASP:HB2	2.04	0.40
2:J:662:GLN:HA	2:J:663:PRO:HD3	1.92	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	359/364 (99%)	349 (97%)	10 (3%)	0	100	100
1	B	358/364 (98%)	342 (96%)	16 (4%)	0	100	100
1	C	354/364 (97%)	341 (96%)	13 (4%)	0	100	100
1	D	352/364 (97%)	339 (96%)	13 (4%)	0	100	100
2	H	16/67 (24%)	16 (100%)	0	0	100	100
2	I	16/67 (24%)	16 (100%)	0	0	100	100
2	J	20/67 (30%)	19 (95%)	1 (5%)	0	100	100
2	K	20/67 (30%)	19 (95%)	1 (5%)	0	100	100
All	All	1495/1724 (87%)	1441 (96%)	54 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	309/311 (99%)	301 (97%)	8 (3%)	46	75
1	B	308/311 (99%)	302 (98%)	6 (2%)	57	82
1	C	306/311 (98%)	297 (97%)	9 (3%)	42	71
1	D	305/311 (98%)	296 (97%)	9 (3%)	41	70
2	H	19/59 (32%)	19 (100%)	0	100	100
2	I	19/59 (32%)	18 (95%)	1 (5%)	22	48
2	J	22/59 (37%)	21 (96%)	1 (4%)	27	55
2	K	23/59 (39%)	23 (100%)	0	100	100
All	All	1311/1480 (89%)	1277 (97%)	34 (3%)	46	75

All (34) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	43	LYS
1	A	106	ASP
1	A	111	TRP
1	A	174	GLN
1	A	223	GLN
1	A	245	LYS
1	A	296	ASN
1	A	325	LEU
1	B	70	SER
1	B	111	TRP
1	B	176	ARG
1	B	185	SER
1	B	326	SER
1	B	349	LEU
1	C	111	TRP
1	C	133	GLU
1	C	155	ASN
1	C	221	ARG
1	C	284	LEU
1	C	299	SER
1	C	326	SER
1	C	350	ARG
1	C	357	LEU
1	D	14	GLN
1	D	107	ASP

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	D	111	TRP
1	D	171	SER
1	D	173	GLN
1	D	258	ASN
1	D	294	TYR
1	D	299	SER
1	D	325	LEU
2	I	717	SER
2	J	664	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	223	GLN
1	C	3	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.



## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	361/364 (99%)	0.16	0 <span style="border: 1px solid blue; padding: 2px;">100</span> <span style="border: 1px solid blue; padding: 2px;">100</span>	39, 61, 83, 109	0
1	B	360/364 (98%)	0.19	6 (1%) <span style="border: 1px solid blue; padding: 2px;">70</span> <span style="border: 1px solid blue; padding: 2px;">72</span>	39, 59, 86, 107	0
1	C	356/364 (97%)	0.50	19 (5%) <span style="border: 1px solid red; padding: 2px;">26</span> <span style="border: 1px solid red; padding: 2px;">25</span>	52, 83, 105, 119	0
1	D	354/364 (97%)	0.44	18 (5%) <span style="border: 1px solid red; padding: 2px;">28</span> <span style="border: 1px solid red; padding: 2px;">26</span>	48, 76, 107, 119	0
2	H	20/67 (29%)	0.41	0 <span style="border: 1px solid blue; padding: 2px;">100</span> <span style="border: 1px solid blue; padding: 2px;">100</span>	58, 77, 103, 109	0
2	I	20/67 (29%)	0.29	0 <span style="border: 1px solid blue; padding: 2px;">100</span> <span style="border: 1px solid blue; padding: 2px;">100</span>	56, 77, 99, 101	0
2	J	24/67 (35%)	1.67	9 (37%) <span style="border: 1px solid red; padding: 2px;">0</span> <span style="border: 1px solid red; padding: 2px;">0</span>	70, 91, 106, 107	0
2	K	24/67 (35%)	1.00	4 (16%) <span style="border: 1px solid red; padding: 2px;">1</span> <span style="border: 1px solid red; padding: 2px;">1</span>	65, 83, 103, 114	0
All	All	1519/1724 (88%)	0.35	56 (3%) <span style="border: 1px solid red; padding: 2px;">41</span> <span style="border: 1px solid red; padding: 2px;">41</span>	39, 69, 103, 119	0

All (56) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	44	VAL	5.8
2	J	708	PRO	5.2
1	B	47	GLN	4.4
2	J	707	SER	3.7
1	B	45	GLY	3.7
1	C	260	PHE	3.6
2	J	661	SER	3.6
1	C	7	ILE	3.5
1	D	225	GLY	3.5
2	J	713	LEU	3.2
1	D	262	VAL	3.2
1	C	5	LEU	3.1
1	C	44	VAL	3.1
1	D	6	PRO	3.1
2	J	711	GLY	3.0
1	C	152	GLN	3.0

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
2	K	716	LEU	3.0
1	B	238	THR	2.9
2	J	665	ILE	2.8
1	D	15	LEU	2.7
2	J	710	VAL	2.7
2	K	709	VAL	2.7
1	D	282	ILE	2.7
1	D	7	ILE	2.6
1	C	284	LEU	2.5
1	D	172	ALA	2.5
2	J	664	LEU	2.5
1	C	171	SER	2.5
1	C	13	LEU	2.5
2	J	667	LEU	2.4
1	D	298	ILE	2.4
1	D	177	VAL	2.4
1	C	218	PHE	2.3
2	K	710	VAL	2.3
1	C	172	ALA	2.3
1	C	298	ILE	2.2
1	C	108	VAL	2.2
1	D	324	VAL	2.2
1	D	349	LEU	2.2
1	C	178	VAL	2.2
1	D	341	VAL	2.2
1	C	137	GLN	2.2
1	C	148	LEU	2.2
1	B	224	ALA	2.2
1	D	224	ALA	2.2
1	C	282	ILE	2.1
1	D	5	LEU	2.1
2	K	711	GLY	2.1
1	D	340	ASN	2.1
1	D	67	SER	2.1
1	C	322	GLY	2.1
1	D	284	LEU	2.0
1	C	4	ILE	2.0
1	B	294	TYR	2.0
1	D	343	GLN	2.0
1	C	107	ASP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.