



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 9, 2024 – 03:25 pm GMT

PDB ID : 8QN3  
Title : OPR3 wildtype in complex with NADH4  
Authors : Bijelic, A.; Macheroux, P.; Keschbaumer, B.  
Deposited on : 2023-09-25  
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2340 (1.76-1.76)
Clashscore	141614	2466 (1.76-1.76)
Ramachandran outliers	138981	2437 (1.76-1.76)
Sidechain outliers	138945	2437 (1.76-1.76)
RSRZ outliers	127900	2298 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	402	 6% 82% 9% 9%
1	B	402	 26% 76% 12% 12%

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5821 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 12-oxophytodienoate reductase 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	364	2797	1779	497	511	10	0	0	0
1	B	355	2617	1669	461	478	9	0	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	HIS	-	expression tag	UNP Q9FEW9
A	-4	HIS	-	expression tag	UNP Q9FEW9
A	-3	HIS	-	expression tag	UNP Q9FEW9
A	-2	HIS	-	expression tag	UNP Q9FEW9
A	-1	HIS	-	expression tag	UNP Q9FEW9
A	0	HIS	-	expression tag	UNP Q9FEW9
B	-5	HIS	-	expression tag	UNP Q9FEW9
B	-4	HIS	-	expression tag	UNP Q9FEW9
B	-3	HIS	-	expression tag	UNP Q9FEW9
B	-2	HIS	-	expression tag	UNP Q9FEW9
B	-1	HIS	-	expression tag	UNP Q9FEW9
B	0	HIS	-	expression tag	UNP Q9FEW9

- Molecule 2 is FLAVIN MONONUCLEOTIDE (three-letter code: FMN) (formula: C<sub>17</sub>H<sub>21</sub>N<sub>4</sub>O<sub>9</sub>P).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	31	17	4	9	1	0	0
2	B	1	31	17	4	9	1	0	0

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O S		
3	A	1	5	4 1	0	0
3	A	1	5	4 1	0	0

- Molecule 4 is 1,4,5,6-Tetrahydronicotinamide adenine dinucleotide (three-letter code: WI6) (formula:  $C_{21}H_{31}N_7O_{14}P_2$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf		
			Total	C	N	O			P	
4	A	1	Total	26	11	2	11	2	0	0
4	B	1	Total	26	11	2	11	2	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
5	A	264	Total	264	0	0
5	B	19	Total	19	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 12-oxophytodienoate reductase 3



- Molecule 1: 12-oxophytodienoate reductase 3



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	57.74Å 89.46Å 81.02Å 90.00° 107.40° 90.00°	Depositor
Resolution (Å)	46.91 – 1.75 46.91 – 1.75	Depositor EDS
% Data completeness (in resolution range)	99.9 (46.91-1.75) 99.9 (46.91-1.75)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.82 (at 1.75Å)	Xtrriage
Refinement program	PHENIX dev_4761, PHENIX dev_4761	Depositor
R, $R_{free}$	0.200 , 0.225 0.198 , 0.222	Depositor DCC
$R_{free}$ test set	3954 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	34.4	Xtrriage
Anisotropy	0.502	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 49.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	5821	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	51.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.68% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, FMN, WI6

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.38	4/2862 (0.1%)	0.58	1/3893 (0.0%)
1	B	0.26	0/2681	0.50	0/3663
All	All	0.33	4/5543 (0.1%)	0.55	1/7556 (0.0%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	84	GLU	C-N	-5.36	1.21	1.34
1	A	156	GLY	C-N	-5.29	1.21	1.34
1	A	83	LYS	C-N	-5.25	1.22	1.34
1	A	17	LYS	C-N	-5.16	1.22	1.34

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	304	LEU	CA-CB-CG	5.26	127.39	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2797	0	2749	18	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	2617	0	2471	35	0
2	A	31	0	19	3	0
2	B	31	0	19	4	0
3	A	10	0	0	0	0
4	A	26	0	0	0	0
4	B	26	0	0	0	0
5	A	264	0	0	0	0
5	B	19	0	0	0	0
All	All	5821	0	5258	54	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:32:MET:HE2	1:B:342:GLY:HA2	1.65	0.78
1:B:303:ARG:HE	1:B:303:ARG:H	1.34	0.73
1:A:280:THR:HG22	1:A:319:SER:HB3	1.77	0.64
1:B:324:ARG:HE	1:B:328:ILE:HD11	1.62	0.64
2:B:401:FMN:O5'	2:B:401:FMN:O3'	2.05	0.62
1:A:253:LEU:HD13	1:A:304:LEU:HD22	1.82	0.61
1:A:32:MET:HA	2:A:401:FMN:N5	2.19	0.57
1:B:321:GLY:N	2:B:401:FMN:O2P	2.28	0.56
1:B:370:TYR:CZ	2:B:401:FMN:HM72	2.42	0.55
1:B:32:MET:CE	1:B:342:GLY:HA2	2.36	0.54
1:B:27:VAL:HB	1:B:352:VAL:HG22	1.89	0.53
1:B:197:LYS:HD3	1:B:249:ASP:HB2	1.90	0.53
1:B:186:GLY:HA2	1:B:192:ILE:HB	1.90	0.53
1:B:188:HIS:NE2	1:B:280:THR:HG21	2.24	0.53
1:B:324:ARG:O	1:B:328:ILE:HG13	2.09	0.53
1:B:49:TYR:HB2	1:B:378:TYR:CE2	2.45	0.52
1:A:165:GLU:O	1:A:169:ARG:HG3	2.11	0.51
1:B:354:ARG:NH1	1:B:362:ASN:OD1	2.45	0.50
1:A:185:HIS:HD1	1:A:187:ALA:H	1.60	0.49
1:A:197:LYS:HD3	1:A:249:ASP:HB2	1.94	0.48
1:A:253:LEU:HD11	1:A:307:THR:HG21	1.94	0.48
1:B:133:ILE:HG13	1:B:245:LEU:HD22	1.95	0.48
1:B:133:ILE:HG21	1:B:137:TRP:HB2	1.95	0.48
1:B:303:ARG:H	1:B:303:ARG:NE	2.07	0.48
1:B:112:ARG:HB3	1:B:127:SER:HB2	1.96	0.47

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:32:MET:HB3	2:A:401:FMN:C6	2.45	0.47
1:B:253:LEU:HA	1:B:304:LEU:HD13	1.96	0.46
1:A:216:LYS:O	1:A:220:GLN:HG3	2.16	0.46
1:B:205:ASP:HB2	1:B:206:GLU:OE1	2.16	0.46
1:B:71:SER:HB2	1:B:109:HIS:HA	1.98	0.45
1:B:354:ARG:HB3	1:B:359:ALA:HB3	1.99	0.45
1:B:237:ARG:HA	1:B:278:HIS:O	2.16	0.45
1:B:112:ARG:NH1	1:B:131:LYS:O	2.45	0.45
1:B:206:GLU:OE1	1:B:206:GLU:N	2.31	0.44
1:A:305:MET:HE3	1:A:305:MET:HB2	1.64	0.43
1:A:60:LEU:O	1:A:103:ILE:HA	2.19	0.43
1:B:348:ASN:HD21	1:B:362:ASN:N	2.17	0.42
1:B:62:THR:HG23	1:B:104:PHE:O	2.19	0.42
1:B:34:ARG:HA	1:B:34:ARG:HD2	1.80	0.42
1:B:62:THR:HG22	1:B:103:ILE:HD11	2.02	0.42
1:B:370:TYR:CE2	2:B:401:FMN:HM72	2.55	0.42
1:A:324:ARG:HG3	1:A:355:ILE:HG23	2.02	0.42
1:B:112:ARG:NH2	1:B:200:ILE:HG12	2.34	0.42
1:A:87:ARG:NE	1:A:88:GLU:OE2	2.53	0.41
1:A:370:TYR:CZ	2:A:401:FMN:HM72	2.55	0.41
1:B:32:MET:H	1:B:63:GLU:HA	1.85	0.41
1:B:251:ASN:ND2	1:B:254:SER:HB2	2.35	0.41
1:B:350:ASP:O	1:B:354:ARG:HG3	2.21	0.41
1:A:129:THR:OG1	1:A:131:LYS:HD3	2.21	0.41
1:B:20:LYS:HB3	1:B:20:LYS:HE3	1.80	0.41
1:A:80:ILE:HD12	1:A:89:TRP:CD1	2.56	0.41
1:A:237:ARG:HA	1:A:278:HIS:O	2.21	0.41
1:A:210:SER:O	1:A:214:ARG:HG3	2.21	0.40
1:B:350:ASP:OD1	1:B:350:ASP:N	2.47	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	360/402 (90%)	349 (97%)	11 (3%)	0	100	100
1	B	351/402 (87%)	335 (95%)	16 (5%)	0	100	100
All	All	711/804 (88%)	684 (96%)	27 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	289/328 (88%)	287 (99%)	2 (1%)	84	75
1	B	254/328 (77%)	245 (96%)	9 (4%)	36	13
All	All	543/656 (83%)	532 (98%)	11 (2%)	55	34

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	309	ARG
1	A	345	PHE
1	B	14	SER
1	B	20	LYS
1	B	77	VAL
1	B	185	HIS
1	B	203	ARG
1	B	303	ARG
1	B	309	ARG
1	B	345	PHE
1	B	362	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	FMN	A	401	-	33,33,33	1.03	2 (6%)	48,50,50	1.31	7 (14%)
3	SO4	A	402	-	4,4,4	0.59	0	6,6,6	0.07	0
2	FMN	B	401	-	33,33,33	1.07	2 (6%)	48,50,50	1.28	7 (14%)
3	SO4	A	404	-	4,4,4	0.60	0	6,6,6	0.04	0
4	WI6	B	402	-	24,27,48	0.91	1 (4%)	33,41,73	1.26	3 (9%)
4	WI6	A	403	-	24,27,48	0.91	1 (4%)	33,41,73	1.45	5 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FMN	A	401	-	-	1/18/18/18	0/3/3/3
4	WI6	B	402	-	-	5/20/46/72	0/2/2/5
2	FMN	B	401	-	-	7/18/18/18	0/3/3/3
4	WI6	A	403	-	-	4/20/46/72	0/2/2/5

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	FMN	C4A-N5	3.86	1.38	1.30
2	B	401	FMN	C4A-N5	3.86	1.38	1.30
4	A	403	WI6	PA-O5'	2.62	1.64	1.54
4	B	402	WI6	PA-O5'	2.59	1.64	1.54
2	B	401	FMN	C10-N1	2.52	1.38	1.33
2	A	401	FMN	C10-N1	2.17	1.37	1.33

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	403	WI6	C3N-C7N-N7N	-4.22	112.29	116.85
2	A	401	FMN	C5'-C4'-C3'	-3.83	104.81	112.20
4	A	403	WI6	C3N-C4N-C5N	-3.25	117.63	123.09
4	B	402	WI6	C3N-C4N-C5N	-3.11	117.86	123.09
2	B	401	FMN	C4-N3-C2	-2.99	120.12	125.64
2	A	401	FMN	C4A-C10-N10	2.91	120.73	116.48
2	A	401	FMN	C4-N3-C2	-2.76	120.55	125.64
4	A	403	WI6	O1-PA-O2P	2.72	121.31	110.68
4	B	402	WI6	O1-PA-O2P	2.70	121.24	110.68
2	B	401	FMN	C9A-C5A-N5	-2.68	119.52	122.43
2	B	401	FMN	C10-C4A-N5	-2.61	119.33	124.86
2	B	401	FMN	C4A-C4-N3	2.57	119.73	113.19
4	B	402	WI6	C3N-C7N-N7N	-2.48	114.18	116.85
2	B	401	FMN	O4-C4-C4A	-2.45	120.10	126.60
2	A	401	FMN	C4A-C4-N3	2.43	119.36	113.19
2	A	401	FMN	C10-C4A-N5	-2.39	119.78	124.86
4	A	403	WI6	O7N-C7N-C3N	2.30	123.98	120.92
2	B	401	FMN	C4A-C10-N10	2.24	119.76	116.48
2	A	401	FMN	C4A-C10-N1	-2.13	119.80	124.73
2	A	401	FMN	O4-C4-C4A	-2.11	121.00	126.60
2	B	401	FMN	C4A-C10-N1	-2.03	120.02	124.73
4	A	403	WI6	O5'-PA-O3	-2.01	97.88	104.64

There are no chirality outliers.

All (17) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	401	FMN	C1'-C2'-C3'-O3'
2	B	401	FMN	C1'-C2'-C3'-C4'
2	B	401	FMN	O2'-C2'-C3'-O3'
2	B	401	FMN	O2'-C2'-C3'-C4'
2	B	401	FMN	C5'-O5'-P-O1P

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Res	Type	Atoms
4	A	403	WI6	C2N-C3N-C7N-O7N
4	A	403	WI6	C2N-C3N-C7N-N7N
4	A	403	WI6	O4'A-C1'A-N1N-C2N
4	B	402	WI6	C4N-C3N-C7N-O7N
4	B	402	WI6	O4'A-C4'A-C5'A-O5'A
4	B	402	WI6	C3'A-C4'A-C5'A-O5'A
4	B	402	WI6	C4N-C3N-C7N-N7N
2	A	401	FMN	C4'-C5'-O5'-P
4	A	403	WI6	C2'A-C1'A-N1N-C2N
4	B	402	WI6	PA-O3-PN-O1N
2	B	401	FMN	C5'-O5'-P-O3P
2	B	401	FMN	N10-C1'-C2'-O2'


There are no ring outliers.

2 monomers are involved in 7 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	FMN	3	0
2	B	401	FMN	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	364/402 (90%)	0.46	23 (6%) 20 25	24, 35, 49, 76	0
1	B	355/402 (88%)	1.60	105 (29%) 0 0	43, 70, 86, 98	0
All	All	719/804 (89%)	1.03	128 (17%) 1 2	24, 48, 85, 98	0

All (128) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	357	LEU	13.0
1	B	328	ILE	10.1
1	B	360	PRO	8.6
1	B	302	ALA	8.1
1	A	287	TYR	7.5
1	B	145	THR	7.4
1	B	341	TYR	7.0
1	B	352	VAL	6.6
1	B	355	ILE	6.6
1	B	215	CYS	6.4
1	B	376	VAL	5.9
1	B	345	PHE	5.6
1	B	356	LYS	5.6
1	B	27	VAL	5.5
1	B	353	MET	5.5
1	B	98	ALA	5.5
1	B	349	PRO	5.4
1	B	327	GLY	5.2
1	B	382	PRO	5.2
1	A	290	THR	5.0
1	A	35	CYS	5.0
1	B	364	TYR	4.9
1	B	348	ASN	4.9
1	B	331	VAL	4.9

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	B	350	ASP	4.8
1	B	362	ASN	4.8
1	B	381	TYR	4.8
1	B	351	LEU	4.8
1	B	334	GLY	4.7
1	B	359	ALA	4.7
1	B	324	ARG	4.6
1	B	379	THR	4.6
1	B	243	ASP	4.5
1	B	383	PHE	4.5
1	B	363	LYS	4.4
1	B	380	ASP	4.1
1	B	342	GLY	4.1
1	B	13	PHE	4.1
1	B	45	ALA	4.0
1	B	332	ALA	4.0
1	B	54	ALA	4.0
1	B	49	TYR	4.0
1	B	361	LEU	3.9
1	B	378	TYR	3.9
1	B	326	LEU	3.8
1	B	77	VAL	3.7
1	B	323	THR	3.7
1	B	375	VAL	3.6
1	A	286	ALA	3.6
1	B	344	LEU	3.6
1	B	280	THR	3.5
1	B	305	MET	3.5
1	B	155	ILE	3.4
1	B	208	GLY	3.3
1	B	358	ASN	3.3
1	B	95	VAL	3.3
1	B	253	LEU	3.3
1	B	339	VAL	3.3
1	B	336	ALA	3.3
1	B	94	ASP	3.2
1	B	259	VAL	3.2
1	B	369	PHE	3.2
1	B	347	SER	3.2
1	B	354	ARG	3.1
1	B	124	ALA	3.1
1	B	321	GLY	3.1

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	B	75	PRO	3.1
1	A	182	ILE	3.0
1	B	252	PRO	3.0
1	A	242	ILE	3.0
1	B	121	ALA	3.0
1	B	76	HIS	2.9
1	B	198	ASP	2.9
1	B	209	GLY	2.9
1	B	325	GLU	2.9
1	B	374	PRO	2.8
1	B	11	PRO	2.8
1	B	141	MET	2.8
1	B	28	VAL	2.8
1	A	62	THR	2.7
1	B	108	TRP	2.7
1	B	207	TYR	2.7
1	B	115	HIS	2.7
1	B	56	ALA	2.6
1	B	371	THR	2.6
1	B	24	SER	2.6
1	B	249	ASP	2.6
1	B	250	SER	2.6
1	B	137	TRP	2.6
1	B	310	ASN	2.5
1	B	204	THR	2.5
1	B	146	HIS	2.5
1	A	288	GLY	2.5
1	B	38	LEU	2.5
1	A	229	ILE	2.5
1	B	240	PRO	2.5
1	B	367	LYS	2.5
1	B	202	ASP	2.5
1	B	158	TYR	2.4
1	A	289	GLN	2.4
1	B	343	ARG	2.4
1	B	143	ASP	2.4
1	A	305	MET	2.4
1	B	25	HIS	2.4
1	B	312	TYR	2.4
1	B	377	GLY	2.4
1	A	89	TRP	2.3
1	B	241	ALA	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	311	ALA	2.3
1	A	105	CYS	2.3
1	A	302	ALA	2.3
1	A	107	LEU	2.3
1	B	196	LEU	2.3
1	A	41	ILE	2.3
1	B	133	ILE	2.2
1	B	206	GLU	2.2
1	A	86	VAL	2.2
1	B	211	LEU	2.2
1	B	212	ALA	2.2
1	A	234	VAL	2.2
1	A	263	LEU	2.2
1	B	113	ALA	2.1
1	A	235	GLY	2.1
1	A	252	PRO	2.1
1	A	303	ARG	2.1
1	A	67	ILE	2.1
1	B	41	ILE	2.1
1	B	308	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	WI6	B	402	26/44	0.74	0.23	67,82,119,133	0
3	SO4	A	404	5/5	0.78	0.21	46,52,55,57	5
4	WI6	A	403	26/44	0.84	0.15	23,43,85,139	0

*Continued on next page...*

*Continued from previous page...*




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	FMN	B	401	31/31	0.87	0.18	63,73,84,86	0
2	FMN	A	401	31/31	0.94	0.12	24,27,31,32	0
3	SO4	A	402	5/5	0.97	0.12	50,51,54,56	5

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around WI6 A 403:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around FMN B 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







## 6.5 Other polymers [i](#)

There are no such residues in this entry.