



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 29, 2024 – 09:11 PM EDT

PDB ID : 4QPM
Title : Structure of Bub1 kinase domain
Authors : Lin, Z.H.; Jia, L.Y.; Tomchick, D.R.; Luo, X.L.; Yu, H.T.
Deposited on : 2014-06-24
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	5791 (2.20-2.20)
Clashscore	180529	6634 (2.20-2.20)
Ramachandran outliers	177936	6560 (2.20-2.20)
Sidechain outliers	177891	6561 (2.20-2.20)
RSRZ outliers	164620	5791 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	356	
1	B	356	

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 11035 atoms, of which 5390 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Mitotic checkpoint serine/threonine-protein kinase BUB1.

Mol	Chain	Residues	Atoms								ZeroOcc	AltConf	Trace
			Total	C	H	N	O	P	S				
1	A	330	5376	1749	2693	445	466	1	22	0	0	0	
1	B	329	5348	1744	2675	440	466	1	22	0	0	0	

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	730	GLY	-	expression tag	UNP O43683
A	731	ALA	-	expression tag	UNP O43683
A	732	MET	-	expression tag	UNP O43683
A	733	ASP	-	expression tag	UNP O43683
A	734	PRO	-	expression tag	UNP O43683
A	735	GLU	-	expression tag	UNP O43683
A	736	PHE	-	expression tag	UNP O43683
A	737	GLY	-	expression tag	UNP O43683
A	738	ARG	-	expression tag	UNP O43683
A	739	PRO	-	expression tag	UNP O43683
B	730	GLY	-	expression tag	UNP O43683
B	731	ALA	-	expression tag	UNP O43683
B	732	MET	-	expression tag	UNP O43683
B	733	ASP	-	expression tag	UNP O43683
B	734	PRO	-	expression tag	UNP O43683
B	735	GLU	-	expression tag	UNP O43683
B	736	PHE	-	expression tag	UNP O43683
B	737	GLY	-	expression tag	UNP O43683
B	738	ARG	-	expression tag	UNP O43683
B	739	PRO	-	expression tag	UNP O43683

- Molecule 2 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	H	N	O			P
2	A	1	38	10	11	5	10	2	0	0
2	B	1	38	10	11	5	10	2	0	0

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
3	A	2	2	2	0	0
3	B	2	2	2	0	0

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Cl		
4	A	1	1	1	0	0
4	B	2	2	2	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
5	A	121	121	121	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	107	Total 107	O 107	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Mitotic checkpoint serine/threonine-protein kinase BUB1



- Molecule 1: Mitotic checkpoint serine/threonine-protein kinase BUB1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	90.01Å 47.18Å 93.15Å 90.00° 107.40° 90.00°	Depositor
Resolution (Å)	36.88 – 2.20 36.88 – 2.20	Depositor EDS
% Data completeness (in resolution range)	98.1 (36.88-2.20) 98.1 (36.88-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.77 (at 2.20Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.8.3_1479)	Depositor
R, R_{free}	0.229 , 0.269 0.235 , 0.273	Depositor DCC
R_{free} test set	1885 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	32.3	Xtrriage
Anisotropy	0.359	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 43.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.125 for l,-k,h	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	11035	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.54% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ADP, CL, SEP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.26	0/2743	0.37	0/3698
1	B	0.22	0/2733	0.37	0/3687
All	All	0.24	0/5476	0.37	0/7385

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2683	2693	2704	3	0
1	B	2673	2675	2688	2	0
2	A	27	11	12	0	0
2	B	27	11	12	0	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
4	A	1	0	0	0	0
4	B	2	0	0	1	0
5	A	121	0	0	1	0
5	B	107	0	0	2	0
All	All	5645	5390	5416	6	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (6) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1055:LYS:NZ	1:B:1059:GLN:OE1	2.28	0.65
4:B:1505:CL:CL	5:B:1655:HOH:O	2.50	0.65
1:A:939:SER:N	5:A:1668:HOH:O	2.30	0.65
1:B:972:GLN:NE2	5:B:1669:HOH:O	2.37	0.57
1:A:1055:LYS:NZ	1:A:1059:GLN:OE1	2.52	0.43
1:A:1040:ASP:CG	1:A:1043:HIS:HD1	2.23	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	323/356 (91%)	312 (97%)	11 (3%)	0	100	100
1	B	322/356 (90%)	307 (95%)	15 (5%)	0	100	100
All	All	645/712 (91%)	619 (96%)	26 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	294/315 (93%)	293 (100%)	1 (0%)	91	96
1	B	293/315 (93%)	291 (99%)	2 (1%)	81	90
All	All	587/630 (93%)	584 (100%)	3 (0%)	86	93

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	949	GLN
1	B	827	ASN
1	B	1063	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
1	SEP	A	969	1	8,9,10	1.62	1 (12%)	7,12,14	1.50	1 (14%)
1	SEP	B	969	1	8,9,10	1.62	1 (12%)	7,12,14	1.76	1 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	SEP	A	969	1	-	1/6/8/10	-
1	SEP	B	969	1	-	4/6/8/10	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	969	SEP	P-O1P	3.53	1.61	1.50
1	A	969	SEP	P-O1P	3.52	1.61	1.50

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	969	SEP	OG-CB-CA	4.12	112.16	108.14
1	A	969	SEP	OG-CB-CA	3.39	111.44	108.14

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	B	969	SEP	N-CA-CB-OG
1	B	969	SEP	CB-OG-P-O3P
1	B	969	SEP	CB-OG-P-O2P
1	B	969	SEP	CB-OG-P-O1P
1	A	969	SEP	N-CA-CB-OG

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 7 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	ADP	A	1501	3	24,29,29	0.91	1 (4%)	29,45,45	1.20	2 (6%)
2	ADP	B	1501	3	24,29,29	0.95	2 (8%)	29,45,45	1.19	2 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ADP	A	1501	3	-	2/12/32/32	0/3/3/3
2	ADP	B	1501	3	-	2/12/32/32	0/3/3/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1501	ADP	O4'-C1'	2.26	1.43	1.40
2	A	1501	ADP	O4'-C1'	2.15	1.43	1.40
2	B	1501	ADP	PA-O3A	2.06	1.61	1.59

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1501	ADP	N3-C2-N1	-4.31	122.83	128.67
2	B	1501	ADP	N3-C2-N1	-4.22	122.95	128.67
2	B	1501	ADP	C4-C5-N7	-2.42	106.78	109.34
2	A	1501	ADP	C4-C5-N7	-2.27	106.94	109.34

There are no chirality outliers.


All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	1501	ADP	PA-O3A-PB-O2B
2	B	1501	ADP	PA-O3A-PB-O2B
2	A	1501	ADP	PB-O3A-PA-O2A
2	B	1501	ADP	PB-O3A-PA-O2A

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	329/356 (92%)	0.97	41 (12%) 9 7	24, 43, 87, 151	0
1	B	328/356 (92%)	1.28	71 (21%) 3 2	24, 46, 102, 147	0
All	All	657/712 (92%)	1.12	112 (17%) 5 4	24, 44, 93, 151	0

All (112) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	739	PRO	6.2
1	B	929	PHE	6.0
1	B	970	GLY	5.4
1	A	1018	LEU	4.9
1	B	931	GLU	4.6
1	A	967	GLU	4.6
1	B	937	ASP	4.4
1	A	979	ASN	4.3
1	A	941	GLY	4.3
1	B	981	PRO	4.2
1	B	1003	TYR	4.1
1	B	971	PHE	4.1
1	B	1018	LEU	4.1
1	A	966	CYS	4.0
1	B	967	GLU	4.0
1	B	871	TYR	4.0
1	B	786	LEU	3.9
1	B	927	ASN	3.8
1	B	1017	GLY	3.8
1	A	931	GLU	3.8
1	B	851	LYS	3.8
1	A	815	LYS	3.8
1	A	1020	ARG	3.7
1	B	817	LYS	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	981	PRO	3.7
1	A	1083	SER	3.7
1	B	816	GLN	3.6
1	A	893	LEU	3.6
1	B	777	LYS	3.6
1	B	930	LEU	3.5
1	A	1017	GLY	3.5
1	B	1008	ASN	3.4
1	B	867	GLU	3.4
1	B	968	THR	3.4
1	A	939	SER	3.4
1	A	1003	TYR	3.4
1	A	1070	ARG	3.4
1	B	791	HIS	3.4
1	A	1059	GLN	3.3
1	B	974	VAL	3.3
1	A	791	HIS	3.2
1	B	1066	ILE	3.2
1	A	739	PRO	3.2
1	B	780	PHE	3.2
1	A	806	GLN	3.2
1	B	966	CYS	3.2
1	A	940	ALA	3.2
1	B	1011	GLY	3.2
1	B	938	LEU	3.1
1	B	941	GLY	3.0
1	B	1020	ARG	3.0
1	B	1063	THR	2.9
1	A	991	VAL	2.9
1	B	806	GLN	2.8
1	A	780	PHE	2.8
1	B	1081	LYS	2.8
1	B	979	ASN	2.8
1	A	1066	ILE	2.7
1	B	784	SER	2.7
1	B	893	LEU	2.7
1	B	775	LYS	2.7
1	B	869	TYR	2.7
1	A	970	GLY	2.7
1	B	973	CYS	2.7
1	A	995	VAL	2.7
1	B	963	THR	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	971	PHE	2.6
1	B	790	HIS	2.6
1	B	1080	CYS	2.6
1	B	782	LEU	2.5
1	B	1073	LEU	2.5
1	A	968	THR	2.5
1	B	1043	HIS	2.5
1	B	824	LYS	2.5
1	B	965	LYS	2.5
1	B	779	GLU	2.4
1	B	872	GLY	2.4
1	A	782	LEU	2.4
1	A	1060	GLN	2.4
1	B	1042	HIS	2.4
1	B	1079	GLU	2.4
1	B	749	LYS	2.4
1	B	788	TYR	2.4
1	A	1082	ARG	2.4
1	B	982	TRP	2.3
1	B	980	LYS	2.3
1	A	788	TYR	2.3
1	B	949	GLN	2.3
1	B	1012	GLU	2.3
1	B	1010	GLY	2.3
1	B	940	ALA	2.3
1	A	798	PHE	2.3
1	A	929	PHE	2.3
1	A	949	GLN	2.3
1	B	924	ILE	2.3
1	B	740	ASN	2.2
1	B	910	ASP	2.2
1	B	921	ASP	2.2
1	B	1052	GLN	2.2
1	A	851	LYS	2.2
1	A	1041	CYS	2.2
1	B	928	GLY	2.2
1	B	1050	LEU	2.2
1	A	784	SER	2.1
1	A	1012	GLU	2.1
1	A	1010	GLY	2.1
1	A	924	ILE	2.1
1	B	1038	ILE	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	785	LYS	2.0
1	B	1009	GLU	2.0
1	B	988	TYR	2.0
1	B	785	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	SEP	B	969	10/11	0.49	0.18	100,119,145,150	0
1	SEP	A	969	10/11	0.60	0.17	85,107,124,125	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.


6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	CL	B	1504	1/1	0.67	0.23	85,85,85,85	0
4	CL	B	1505	1/1	0.79	0.32	84,84,84,84	0
4	CL	A	1504	1/1	0.82	0.26	75,75,75,75	0
3	MG	A	1503	1/1	0.91	0.07	26,26,26,26	0
2	ADP	A	1501	27/27	0.93	0.09	20,29,36,37	0
2	ADP	B	1501	27/27	0.94	0.08	21,28,40,40	0
3	MG	B	1503	1/1	0.95	0.06	25,25,25,25	0
3	MG	B	1502	1/1	0.95	0.06	28,28,28,28	0
3	MG	A	1502	1/1	0.96	0.04	21,21,21,21	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different

orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.