



Full wwPDB X-ray Structure Validation Report

May 22, 2020 – 08:42 am BST

PDB ID : 4QTH
Title : Crystal structure of anti-uPAR Fab 8B12
Authors : Zhao, B.; Yuan, C.; Luo, Z.; Huang, M.
Deposited on : 2014-07-08
Resolution : 2.17 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.17 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6864 (2.20-2.16)
Clashscore	141614	7689 (2.20-2.16)
Ramachandran outliers	138981	7564 (2.20-2.16)
Sidechain outliers	138945	7564 (2.20-2.16)
RSRZ outliers	127900	6738 (2.20-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	224	 23% 83% 11%
1	H	224	 11% 72% 25%
2	B	214	 25% 78% 20%
2	L	214	 8% 83% 15%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6833 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called anti-uPAR antibody, heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	H	224	1686	1062	276	338	10	0	0	0
1	A	215	1621	1027	262	323	9	0	0	0

- Molecule 2 is a protein called anti-uPAR antibody, light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	L	214	1658	1024	280	347	7	0	0	0
2	B	211	1635	1012	276	341	6	0	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	H	64	Total 64	O 64	0	0
3	L	63	Total 63	O 63	0	0
3	A	38	Total 38	O 38	0	0
3	B	68	Total 68	O 68	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: anti-uPAR antibody, heavy chain




- Molecule 1: anti-uPAR antibody, heavy chain




- Molecule 2: anti-uPAR antibody, light chain



- Molecule 2: anti-uPAR antibody, light chain





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	80.08Å 72.92Å 85.55Å 90.00° 105.37° 90.00°	Depositor
Resolution (Å)	41.31 – 2.17 41.20 – 2.17	Depositor EDS
% Data completeness (in resolution range)	99.5 (41.31-2.17) 99.6 (41.20-2.17)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.71 (at 2.18Å)	Xtrriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.229 , 0.273 0.227 , 0.270	Depositor DCC
R_{free} test set	2547 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	43.4	Xtrriage
Anisotropy	0.074	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 49.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6833	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.64% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.59	0/1664	0.68	0/2275
1	H	0.66	2/1730 (0.1%)	0.76	0/2366
2	B	0.63	1/1668 (0.1%)	0.73	1/2261 (0.0%)
2	L	0.59	1/1691 (0.1%)	0.71	0/2292
All	All	0.62	4/6753 (0.1%)	0.72	1/9194 (0.0%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	35	TRP	CD2-CE2	5.73	1.48	1.41
1	H	197	TRP	CD2-CE2	5.53	1.48	1.41
1	H	33	TRP	CD2-CE2	5.28	1.47	1.41
2	L	35	TRP	CD2-CE2	5.20	1.47	1.41

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	89	LEU	CB-CG-CD2	6.48	122.02	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1621	0	1572	13	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	H	1686	0	1630	35	0
2	B	1635	0	1569	22	0
2	L	1658	0	1585	21	0
3	A	38	0	0	0	0
3	B	68	0	0	2	0
3	H	64	0	0	0	0
3	L	63	0	0	1	0
All	All	6833	0	6356	87	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (87) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:218:LYS:HD3	1:H:218:LYS:H	1.29	0.96
2:B:85:ASP:OD1	2:B:103:LYS:HD3	1.69	0.93
2:L:145:ASN:HB2	2:L:197:THR:HG22	1.55	0.89
2:L:190:ASN:ND2	2:L:210:ASN:OD1	2.10	0.85
2:L:195:GLU:HG2	2:L:206:VAL:HG22	1.61	0.83
2:L:85:ASP:OD1	2:L:103:LYS:HD3	1.79	0.82
2:B:66:ARG:HD2	2:B:68:GLY:O	1.82	0.80
1:H:198:PRO:HB3	1:H:221:PRO:HG3	1.73	0.70
1:H:224:CYS:HA	2:L:214:CYS:HB3	1.74	0.67
2:B:145:ASN:HB2	2:B:197:THR:HG22	1.78	0.66
1:H:152:LYS:HE2	1:H:180:GLN:NE2	2.13	0.64
2:B:69:SER:HB2	3:B:337:HOH:O	1.97	0.63
1:H:224:CYS:HA	2:L:214:CYS:CB	2.29	0.63
2:L:145:ASN:CB	2:L:197:THR:HG22	2.28	0.63
1:H:224:CYS:CA	2:L:214:CYS:HB3	2.27	0.63
2:L:143:ASP:HB2	3:L:355:HOH:O	1.98	0.63
1:H:32:TYR:OH	1:H:107:LEU:HG	2.00	0.61
1:H:163:TRP:CZ3	1:H:204:CYS:HB3	2.34	0.61
2:L:149:LYS:HB2	2:L:193:THR:HB	1.83	0.60
1:H:147:LEU:HD22	1:H:219:ILE:HG21	1.84	0.59
1:H:218:LYS:CD	1:H:218:LYS:H	2.10	0.59
1:H:106:ASN:H	1:H:106:ASN:HD22	1.50	0.59
2:L:78:LEU:HD13	2:L:83:PHE:CZ	2.39	0.58
1:H:198:PRO:HB3	1:H:221:PRO:CG	2.33	0.58
1:A:106:ASN:HD22	1:A:106:ASN:H	1.52	0.57
1:H:152:LYS:HE2	1:H:180:GLN:HE22	1.70	0.57

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:187:GLU:O	2:B:211:ARG:NH1	2.37	0.56
1:A:62:GLU:HA	1:A:65:LYS:HD3	1.89	0.55
1:A:38:ARG:HB2	1:A:48:ILE:HD11	1.88	0.54
2:B:163:TRP:CD1	2:B:175:MET:HG3	2.43	0.54
1:H:162:THR:OG1	1:H:205:ASN:HB2	2.09	0.53
2:L:151:ASP:OD2	2:L:189:HIS:HD2	1.91	0.53
1:H:135:PRO:HD3	1:H:147:LEU:HD23	1.89	0.53
2:B:149:LYS:HB2	2:B:193:THR:HB	1.90	0.53
1:H:36:TRP:C	1:H:37:ILE:HD12	2.31	0.50
2:B:105:GLU:HB3	3:B:317:HOH:O	2.11	0.50
1:A:196:THR:O	1:A:200:GLU:HG2	2.12	0.49
2:B:193:THR:HG23	2:B:206:VAL:HG13	1.93	0.49
2:L:61:ARG:CZ	2:L:79:GLU:HG3	2.42	0.49
2:B:61:ARG:CZ	2:B:79:GLU:HG3	2.43	0.49
1:H:100:PHE:HB2	1:H:107:LEU:HB3	1.94	0.49
1:H:186:LEU:C	1:H:186:LEU:HD12	2.34	0.48
1:A:53:PRO:HB3	1:A:72:ALA:CB	2.42	0.48
2:L:151:ASP:OD2	2:L:189:HIS:CD2	2.66	0.48
2:L:186:TYR:CZ	2:L:211:ARG:HD2	2.48	0.48
1:H:199:SER:HB3	1:H:200:GLU:HG2	1.96	0.48
2:B:49:TYR:O	2:B:50:ALA:C	2.52	0.47
2:B:89:LEU:HG	2:B:98:PHE:CE2	2.50	0.47
1:H:33:TRP:HB2	1:H:99:ASP:HB3	1.97	0.47
1:A:24:ALA:HB1	1:A:27:TYR:CE1	2.50	0.47
1:H:75:SER:OG	2:B:199:LYS:HD2	2.15	0.47
1:H:208:HIS:HB3	1:H:213:THR:HB	1.97	0.47
1:A:73:ASP:OD1	1:A:76:SER:OG	2.33	0.46
1:A:100:PHE:HB2	1:A:107:LEU:HB3	1.97	0.46
2:B:94:SER:HA	2:B:95:PRO:C	2.36	0.46
2:B:150:ILE:HD11	2:B:179:LEU:HD21	1.97	0.46
1:H:11:LEU:HD21	1:H:155:PHE:HZ	1.81	0.46
1:A:30:THR:HA	1:A:53:PRO:O	2.16	0.46
2:B:115:VAL:HA	2:B:135:PHE:O	2.15	0.45
2:L:91:TYR:HA	2:L:96:TYR:CD1	2.51	0.45
2:B:150:ILE:CG2	2:B:189:HIS:HB3	2.47	0.44
1:H:58:THR:O	1:H:59:ASN:OD1	2.34	0.44
1:A:53:PRO:HA	1:A:54:GLY:HA2	1.79	0.44
1:H:223:ASP:O	1:H:224:CYS:HB2	2.17	0.44
1:H:38:ARG:HB2	1:H:48:ILE:HD11	2.00	0.44
2:L:115:VAL:HG12	2:L:207:LYS:HG3	1.99	0.44
2:L:33:LEU:HD22	2:L:71:TYR:HB3	2.00	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:122:SER:HA	2:B:125:LEU:HD12	2.00	0.43
2:B:91:TYR:HA	2:B:96:TYR:CD1	2.53	0.43
2:B:138:ASN:ND2	2:B:170:ASP:OD2	2.51	0.43
1:H:131:TYR:HA	1:H:132:PRO:HD3	1.79	0.43
2:B:113:PRO:HG3	2:B:144:ILE:HD11	2.01	0.42
1:A:68:ALA:HA	1:A:82:GLN:O	2.20	0.42
1:H:223:ASP:O	1:H:224:CYS:CB	2.67	0.42
1:A:32:TYR:OH	1:A:107:LEU:HG	2.20	0.41
1:H:193:PRO:O	1:H:196:THR:HB	2.21	0.41
2:B:31:SER:O	2:B:50:ALA:HA	2.21	0.41
1:H:218:LYS:HD3	1:H:218:LYS:N	2.12	0.41
2:L:192:TYR:O	2:L:208:SER:HA	2.20	0.41
1:H:11:LEU:HD21	1:H:155:PHE:CZ	2.55	0.41
1:H:53:PRO:HB3	1:H:72:ALA:HB1	2.02	0.41
1:A:180:GLN:HB3	1:A:181:SER:H	1.66	0.40
1:H:20:LEU:HD22	1:H:116:THR:HG21	2.02	0.40
1:H:136:GLY:HA2	1:H:222:ARG:HD3	2.04	0.40
2:L:1:ASP:HB3	2:L:95:PRO:HG2	2.04	0.40
1:H:144:MET:HE1	1:H:193:PRO:HD3	2.03	0.40
2:L:193:THR:HA	2:L:207:LYS:O	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	211/224 (94%)	198 (94%)	13 (6%)	0	100	100
1	H	222/224 (99%)	210 (95%)	12 (5%)	0	100	100
2	B	209/214 (98%)	201 (96%)	8 (4%)	0	100	100
2	L	212/214 (99%)	207 (98%)	5 (2%)	0	100	100
All	All	854/876 (98%)	816 (96%)	38 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	185/192 (96%)	173 (94%)	12 (6%)	17	17
1	H	192/192 (100%)	171 (89%)	21 (11%)	6	5
2	B	189/192 (98%)	181 (96%)	8 (4%)	30	35
2	L	192/192 (100%)	182 (95%)	10 (5%)	23	26
All	All	758/768 (99%)	707 (93%)	51 (7%)	16	16

All (51) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	1	GLN
1	H	12	MET
1	H	28	THR
1	H	40	ARG
1	H	57	SER
1	H	65	LYS
1	H	77	ASN
1	H	88	THR
1	H	93	ILE
1	H	106	ASN
1	H	107	LEU
1	H	111	VAL
1	H	137	SER
1	H	140	GLN
1	H	143	SER
1	H	167	SER
1	H	180	GLN
1	H	195	SER
1	H	199	SER
1	H	217	LYS
1	H	218	LYS
2	L	1	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	L	33	LEU
2	L	53	SER
2	L	55	ASP
2	L	66	ARG
2	L	89	LEU
2	L	126	THR
2	L	142	LYS
2	L	190	ASN
2	L	197	THR
1	A	1	GLN
1	A	3	GLN
1	A	13	LYS
1	A	40	ARG
1	A	57	SER
1	A	65	LYS
1	A	74	THR
1	A	76	SER
1	A	93	ILE
1	A	106	ASN
1	A	107	LEU
1	A	180	GLN
2	B	33	LEU
2	B	55	ASP
2	B	89	LEU
2	B	93	THR
2	B	153	SER
2	B	197	THR
2	B	202	THR
2	B	209	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	H	3	GLN
1	H	5	GLN
1	H	106	ASN
1	H	180	GLN
2	L	3	GLN
2	L	156	GLN
2	L	189	HIS
1	A	1	GLN
1	A	5	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	77	ASN
1	A	106	ASN
2	B	3	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	215/224 (95%)	1.35	51 (23%) 0 0	24, 59, 99, 124	16 (7%)
1	H	224/224 (100%)	0.77	25 (11%) 5 5	24, 42, 74, 119	16 (7%)
2	B	211/214 (98%)	1.44	53 (25%) 0 0	20, 46, 108, 119	14 (6%)
2	L	214/214 (100%)	0.71	18 (8%) 11 11	23, 44, 66, 98	14 (6%)
All	All	864/876 (98%)	1.06	147 (17%) 1 1	20, 47, 97, 124	60 (6%)

All (147) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	129	GLY	7.5
1	H	44	GLY	7.0
2	L	208	SER	6.8
1	H	149	CYS	6.7
2	L	134	CYS	6.4
1	A	149	CYS	6.3
2	L	104	LEU	6.1
2	B	104	LEU	6.0
1	A	42	GLY	5.8
2	B	208	SER	5.8
1	H	42	GLY	5.8
2	B	163	TRP	5.8
2	L	163	TRP	5.6
2	L	194	CYS	5.6
2	B	201	SER	5.4
1	A	199	SER	5.3
2	B	187	GLU	5.2
1	H	145	VAL	5.1
1	A	200	GLU	5.1
1	A	201	THR	5.1
1	A	163	TRP	5.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	144	MET	5.1
1	A	219	ILE	5.0
1	A	203	THR	5.0
2	B	128	GLY	4.9
2	B	122	SER	4.9
1	A	211	SER	4.9
2	B	130	ALA	4.8
1	A	220	VAL	4.7
1	H	196	THR	4.5
2	B	10	SER	4.5
2	B	191	SER	4.4
2	B	175	MET	4.4
1	A	44	GLY	4.4
2	B	159	VAL	4.4
2	B	183	LYS	4.2
2	B	181	LEU	4.2
1	A	88	THR	4.2
2	B	210	ASN	4.2
2	B	189	HIS	4.2
1	A	193	PRO	4.2
2	B	152	GLY	4.1
2	B	184	ASP	3.9
2	B	155	ARG	3.9
2	B	185	GLU	3.9
2	B	149	LYS	3.8
2	B	188	ARG	3.8
2	L	175	MET	3.8
1	H	141	THR	3.7
1	A	197	TRP	3.7
2	B	156	GLN	3.7
2	B	20	SER	3.6
2	B	186	TYR	3.6
1	H	17	SER	3.6
2	B	118	PHE	3.5
2	B	148	TRP	3.5
2	B	209	PHE	3.5
2	L	69	SER	3.4
1	A	187	SER	3.3
2	B	202	THR	3.3
2	L	10	SER	3.2
2	B	127	SER	3.2
1	H	41	PRO	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	73	ASP	3.2
2	B	120	PRO	3.1
2	B	121	SER	3.1
1	A	204	CYS	3.1
2	B	112	ALA	3.1
1	A	183	LEU	3.1
1	A	192	VAL	3.1
2	L	214	CYS	3.0
1	H	43	HIS	3.0
2	B	143	ASP	3.0
1	H	59	ASN	3.0
1	H	140	GLN	3.0
1	A	75	SER	3.0
2	B	190	ASN	3.0
2	L	20	SER	3.0
1	A	16	ALA	2.9
1	H	171	GLY	2.9
1	H	142	ASN	2.9
1	A	170	SER	2.9
1	H	204	CYS	2.9
1	A	217	LYS	2.9
2	B	157	ASN	2.9
1	A	132	PRO	2.8
2	B	117	ILE	2.8
2	B	31	SER	2.8
1	A	148	GLY	2.8
1	A	17	SER	2.8
1	A	43	HIS	2.8
1	A	180	GLN	2.7
2	B	125	LEU	2.7
1	H	224	CYS	2.7
1	A	154	TYR	2.6
2	B	203	SER	2.6
1	H	136	GLY	2.6
2	B	111	ALA	2.6
1	A	41	PRO	2.6
2	L	143	ASP	2.6
2	L	127	SER	2.5
2	B	194	CYS	2.5
1	A	145	VAL	2.5
1	A	76	SER	2.5
1	H	193	PRO	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	195	GLU	2.5
2	L	129	GLY	2.5
1	A	167	SER	2.5
1	H	88	THR	2.4
1	A	164	ASN	2.4
1	A	208	HIS	2.4
1	H	167	SER	2.4
1	H	199	SER	2.4
1	A	13	LYS	2.4
1	A	202	VAL	2.4
2	B	205	ILE	2.4
2	L	213	GLU	2.3
1	H	75	SER	2.3
1	H	139	ALA	2.3
1	A	127	PRO	2.3
1	A	184	TYR	2.3
2	B	123	GLU	2.3
1	H	74	THR	2.3
1	A	11	LEU	2.2
2	B	134	CYS	2.2
2	B	200	THR	2.2
1	A	45	LEU	2.2
2	B	146	VAL	2.2
1	H	170	SER	2.2
2	L	94	SER	2.2
1	A	190	VAL	2.2
2	B	206	VAL	2.2
1	A	191	THR	2.1
2	B	17	GLU	2.1
2	B	150	ILE	2.1
1	A	62	GLU	2.1
2	L	106	ILE	2.1
2	L	202	THR	2.1
2	L	68	GLY	2.1
1	H	187	SER	2.1
1	A	196	THR	2.1
1	A	182	ASP	2.1
2	B	151	ASP	2.1
1	A	221	PRO	2.0
1	A	74	THR	2.0
1	A	116	THR	2.0
1	A	1	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.