



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 14, 2018 – 01:36 pm GMT

PDB ID : 3QWQ
Title : Crystal structure of the extracellular domain of the epidermal growth factor receptor in complex with an adnectin
Authors : Sheriff, S.
Deposited on : 2011-02-28
Resolution : 2.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : trunk31020
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk31020


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	4013 (2.80-2.72)
Clashscore	122126	1029 (2.78-2.74)
Ramachandran outliers	120053	1013 (2.78-2.74)
Sidechain outliers	120020	1013 (2.78-2.74)
RSRZ outliers	108989	3920 (2.80-2.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	648	 4% 81% 12% • 5%
2	B	114	 75% 11% • 12%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NAG	A	1420	-	-	-	X

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 5610 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Epidermal growth factor receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	613	4612	2844	815	893	60	0	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	619	HIS	-	EXPRESSION TAG	UNP P00533
A	620	HIS	-	EXPRESSION TAG	UNP P00533
A	621	HIS	-	EXPRESSION TAG	UNP P00533
A	622	HIS	-	EXPRESSION TAG	UNP P00533
A	623	HIS	-	EXPRESSION TAG	UNP P00533
A	624	HIS	-	EXPRESSION TAG	UNP P00533

- Molecule 2 is a protein called ADNECTIN.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	100	747	475	124	148	0	0	0

- Molecule 3 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	14	8	1	5	0	0
3	A	1	14	8	1	5	0	0
3	A	1	14	8	1	5	0	0
3	A	1	14	8	1	5	0	0
3	A	1	14	8	1	5	0	0
3	A	1	14	8	1	5	0	0
3	A	1	14	8	1	5	0	0
3	A	1	14	8	1	5	0	0
3	A	1	14	8	1	5	0	0
3	A	1	14	8	1	5	0	0

- Molecule 4 is BETA-D-MANNOSE (three-letter code: BMA) (formula: $C_6H_{12}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			11	6	5		
4	A	1	Total	C	O	0	0
			11	6	5		
4	A	1	Total	C	O	0	0
			11	6	5		

- Molecule 5 is ALPHA-D-MANNOSE (three-letter code: MAN) (formula: C₆H₁₂O₆).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			11	6	5		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 11 6 5	0	0
5	A	1	Total C O 11 6 5	0	0
5	A	1	Total C O 11 6 5	0	0

- Molecule 6 is ALPHA-L-FUCOSE (three-letter code: FUC) (formula: C₆H₁₂O₅).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 10 6 4	0	0
6	A	1	Total C O 10 6 4	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	12	Total O 12 12	0	0
7	B	2	Total O 2 2	0	0


3 Residue-property plots i

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Epidermal growth factor receptor



- Molecule 2: ADNECTIN



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	68.00Å 72.10Å 262.00Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.47 – 2.75 49.47 – 2.75	Depositor EDS
% Data completeness (in resolution range)	99.4 (49.47-2.75) 99.5 (49.47-2.75)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.04 (at 2.77Å)	Xtrriage
Refinement program	BUSTER-TNT BUSTER 2.13.0, BUSTER 2.13.0	Depositor
R, R_{free}	0.202 , 0.246 0.210 , 0.250	Depositor DCC
R_{free} test set	1058 reflections (3.09%)	wwPDB-VP
Wilson B-factor (Å ²)	58.6	Xtrriage
Anisotropy	0.212	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 59.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	5610	wwPDB-VP
Average B, all atoms (Å ²)	64.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.09% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FUC, BMA, NAG, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.53	0/4702	0.80	1/6383 (0.0%)
2	B	0.54	0/770	0.84	0/1062
All	All	0.53	0/5472	0.80	1/7445 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	159	HIS	C-N-CA	6.14	137.05	121.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4612	0	4321	33	0
2	B	747	0	684	3	0
3	A	140	0	121	0	0
4	A	33	0	26	0	0
5	A	44	0	40	0	0
6	A	20	0	20	0	0
7	A	12	0	0	0	0
7	B	2	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	5610	0	5212	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:87:MET:SD	1:A:87:MET:CE	2.04	1.46
1:A:521:GLU:HB2	1:A:522:PRO:HD3	1.49	0.94
1:A:327:ILE:HD11	1:A:345:LEU:HG	1.54	0.89
1:A:146:SER:HA	1:A:149:LEU:HD23	1.56	0.86
1:A:144:VAL:HG11	1:A:152:MET:HE3	1.65	0.78
1:A:48:ARG:H	1:A:71:THR:HG22	1.59	0.67
1:A:341:ILE:HG21	1:A:345:LEU:HD13	1.78	0.65
1:A:521:GLU:HB2	1:A:522:PRO:CD	2.25	0.65
1:A:48:ARG:N	1:A:71:THR:HG22	2.16	0.61
1:A:141:ARG:H	1:A:141:ARG:HE	1.49	0.60
1:A:141:ARG:HG3	1:A:189:ILE:HD12	1.83	0.58
1:A:516:ASN:CG	1:A:520:GLY:HA3	2.25	0.56
1:A:99:SER:HA	1:A:128:ASN:O	2.07	0.54
1:A:82:ILE:HG21	1:A:226:VAL:HG11	1.90	0.54
1:A:407:LYS:HE3	1:A:435:GLY:HA2	1.89	0.53
1:A:75:ILE:HG13	1:A:108:LEU:HD13	1.92	0.52
1:A:190:ILE:O	1:A:202:LYS:HG2	2.09	0.51
1:A:327:ILE:HD12	1:A:347:ILE:HG13	1.95	0.48
1:A:505:VAL:HG23	1:A:512:VAL:HG23	1.97	0.47
2:B:76:THR:O	2:B:83:TYR:HA	2.16	0.46
2:B:35:THR:HG22	2:B:47:GLU:HB3	1.99	0.45
1:A:424:LEU:O	1:A:492:TRP:HB3	2.16	0.45
1:A:68:ALA:HA	1:A:98:LEU:O	2.17	0.44
1:A:438:ILE:HD12	1:A:465:LYS:HD3	1.98	0.44
1:A:114:ARG:HA	1:A:176:TRP:CD1	2.52	0.43
1:A:198:ARG:HD2	1:A:214:ALA:O	2.19	0.43
1:A:378:THR:HG23	1:A:405:ARG:HE	1.82	0.43
1:A:200:ARG:NH1	1:A:207:CYS:H	2.17	0.43
1:A:81:GLN:HA	1:A:115:ASN:O	2.19	0.42
1:A:527:GLU:HB3	1:A:532:ILE:HD13	2.01	0.42
1:A:200:ARG:HD2	1:A:206:ASP:HA	2.02	0.41
1:A:284:VAL:HG22	1:A:285:ARG:HE	1.84	0.41
1:A:539:LEU:HD22	1:A:540:PRO:HD2	2.03	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:38:GLU:HG2	2:B:40:GLY:H	1.86	0.40
1:A:117:GLN:HG3	1:A:214:ALA:HB1	2.02	0.40
1:A:418:SER:HA	1:A:441:GLY:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	611/648 (94%)	559 (92%)	46 (8%)	6 (1%)	17	31
2	B	98/114 (86%)	91 (93%)	4 (4%)	3 (3%)	4	7
All	All	709/762 (93%)	650 (92%)	50 (7%)	9 (1%)	13	24

All (9) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	306	GLU
1	A	336	LYS
2	B	4	VAL
1	A	159	HIS
2	B	40	GLY
1	A	160	LEU
1	A	601	THR
2	B	42	ASN
1	A	521	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	508/559 (91%)	466 (92%)	42 (8%)	12	23
2	B	77/99 (78%)	71 (92%)	6 (8%)	14	25
All	All	585/658 (89%)	537 (92%)	48 (8%)	12	23

All (48) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	LYS
1	A	21	GLU
1	A	71	THR
1	A	72	VAL
1	A	92	SER
1	A	98	LEU
1	A	114	ARG
1	A	117	GLN
1	A	120	LEU
1	A	125	ARG
1	A	141	ARG
1	A	146	SER
1	A	147	ASP
1	A	157	GLN
1	A	158	ASN
1	A	200	ARG
1	A	202	LYS
1	A	233	GLU
1	A	245	LEU
1	A	266	THR
1	A	284	VAL
1	A	285	ARG
1	A	291	SER
1	A	294	MET
1	A	309	CYS
1	A	318	ILE
1	A	353	ARG
1	A	406	THR
1	A	408	GLN
1	A	427	ARG
1	A	439	ILE
1	A	445	LEU
1	A	446	CYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	449	ASN
1	A	489	GLU
1	A	523	ARG
1	A	539	LEU
1	A	544	ASN
1	A	545	ILE
1	A	550	ARG
1	A	576	MET
1	A	612	CYS
2	B	6	ARG
2	B	19	LEU
2	B	49	THR
2	B	54	VAL
2	B	71	THR
2	B	93	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	49	ASN
1	A	121	HIS
1	A	128	ASN
1	A	394	HIS
1	A	480	GLN
1	A	560	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry

19 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	A	1032	1,3,6	14,14,15	0.26	0	17,19,21	0.79	0
3	NAG	A	1033	3,4	14,14,15	0.38	0	17,19,21	0.75	0
4	BMA	A	1034	3,5	11,11,12	0.32	0	15,15,17	1.96	1 (6%)
5	MAN	A	1035	4	11,11,12	0.43	0	15,15,17	0.89	1 (6%)
5	MAN	A	1036	4	11,11,12	0.32	0	15,15,17	1.12	1 (6%)
6	FUC	A	1037	3	9,10,11	0.45	0	13,14,16	1.04	1 (7%)
3	NAG	A	1328	1,3	14,14,15	0.28	0	17,19,21	0.89	2 (11%)
3	NAG	A	1329	3,4	14,14,15	0.22	0	17,19,21	2.40	3 (17%)
4	BMA	A	1330	3	11,11,12	0.47	0	15,15,17	1.10	1 (6%)
3	NAG	A	1337	1,3	14,14,15	0.32	0	17,19,21	1.83	2 (11%)
3	NAG	A	1338	3	14,14,15	0.33	0	17,19,21	1.56	2 (11%)
3	NAG	A	1389	1	14,14,15	0.35	0	17,19,21	2.27	5 (29%)
3	NAG	A	1420	1	14,14,15	0.33	0	17,19,21	2.61	2 (11%)
3	NAG	A	1504	1,3,6	14,14,15	0.26	0	17,19,21	1.19	1 (5%)
3	NAG	A	1505	3,4	14,14,15	0.24	0	17,19,21	0.76	1 (5%)
4	BMA	A	1506	3,5	11,11,12	0.45	0	15,15,17	2.10	2 (13%)
5	MAN	A	1507	4	11,11,12	0.50	0	15,15,17	1.53	1 (6%)
5	MAN	A	1508	4	11,11,12	0.41	0	15,15,17	1.40	1 (6%)
6	FUC	A	1509	3	9,10,11	0.41	0	13,14,16	0.88	1 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	1032	1,3,6	-	0/6/23/26	0/1/1/1
3	NAG	A	1033	3,4	-	0/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	BMA	A	1034	3,5	-	0/2/19/22	1/1/1/1
5	MAN	A	1035	4	-	0/2/19/22	0/1/1/1
5	MAN	A	1036	4	-	0/2/19/22	0/1/1/1
6	FUC	A	1037	3	-	0/0/17/20	0/1/1/1
3	NAG	A	1328	1,3	-	0/6/23/26	0/1/1/1
3	NAG	A	1329	3,4	-	0/6/23/26	0/1/1/1
4	BMA	A	1330	3	-	0/2/19/22	0/1/1/1
3	NAG	A	1337	1,3	-	0/6/23/26	0/1/1/1
3	NAG	A	1338	3	-	0/6/23/26	0/1/1/1
3	NAG	A	1389	1	-	0/6/23/26	0/1/1/1
3	NAG	A	1420	1	-	0/6/23/26	0/1/1/1
3	NAG	A	1504	1,3,6	-	0/6/23/26	0/1/1/1
3	NAG	A	1505	3,4	-	0/6/23/26	0/1/1/1
4	BMA	A	1506	3,5	-	0/2/19/22	1/1/1/1
5	MAN	A	1507	4	-	0/2/19/22	0/1/1/1
5	MAN	A	1508	4	-	0/2/19/22	0/1/1/1
6	FUC	A	1509	3	-	0/0/17/20	0/1/1/1

There are no bond length outliers.

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1329	NAG	O5-C1-C2	-4.67	105.07	111.52
3	A	1328	NAG	O5-C1-C2	-2.16	108.54	111.52
3	A	1389	NAG	C2-N2-C7	2.02	125.89	122.94
3	A	1338	NAG	C3-C4-C5	2.03	113.88	110.24
3	A	1389	NAG	C3-C4-C5	2.09	113.98	110.24
5	A	1035	MAN	C3-C4-C5	2.22	114.22	110.24
3	A	1505	NAG	C1-O5-C5	2.36	115.43	112.19
3	A	1328	NAG	C1-C2-N2	2.39	114.58	110.49
6	A	1509	FUC	C1-O5-C5	2.57	118.07	112.39
3	A	1389	NAG	O5-C1-C2	3.12	115.83	111.52
6	A	1037	FUC	C1-O5-C5	3.18	119.42	112.39
5	A	1036	MAN	C1-O5-C5	3.38	116.84	112.19
4	A	1330	BMA	C1-O5-C5	3.46	116.94	112.19
4	A	1506	BMA	O3-C3-C2	3.69	116.89	110.04
3	A	1337	NAG	O5-C1-C2	3.71	116.64	111.52
3	A	1329	NAG	C1-O5-C5	4.11	117.84	112.19
3	A	1504	NAG	C1-O5-C5	4.11	117.85	112.19
5	A	1508	MAN	C1-O5-C5	4.36	118.19	112.19
3	A	1338	NAG	C1-O5-C5	5.07	119.16	112.19
5	A	1507	MAN	C1-O5-C5	5.16	119.28	112.19

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1389	NAG	C1-O5-C5	5.49	119.74	112.19
3	A	1337	NAG	C1-O5-C5	5.49	119.74	112.19
3	A	1389	NAG	C1-C2-N2	5.88	120.54	110.49
4	A	1506	BMA	C1-O5-C5	6.77	121.51	112.19
4	A	1034	BMA	C1-O5-C5	6.83	121.58	112.19
3	A	1420	NAG	C1-O5-C5	7.10	121.96	112.19
3	A	1329	NAG	C1-C2-N2	7.26	122.90	110.49
3	A	1420	NAG	O5-C1-C2	7.30	121.61	111.52

There are no chirality outliers.

There are no torsion outliers.

All (2) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1034	BMA	C1-C2-C3-C4-C5-O5
4	A	1506	BMA	C1-C2-C3-C4-C5-O5

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	613/648 (94%)	0.44	28 (4%) 32 27	32, 65, 95, 141	0
2	B	100/114 (87%)	0.27	0 100 100	32, 45, 74, 107	0
All	All	713/762 (93%)	0.42	28 (3%) 39 34	32, 63, 94, 141	0

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	614	THR	5.7
1	A	160	LEU	4.0
1	A	336	LYS	3.7
1	A	610	GLU	3.6
1	A	202	LYS	3.3
1	A	194	GLN	3.0
1	A	289	ALA	3.0
1	A	532	ILE	2.8
1	A	526	VAL	2.7
1	A	321	PHE	2.7
1	A	521	GLU	2.6
1	A	332	ILE	2.6
1	A	605	THR	2.5
1	A	528	ASN	2.5
1	A	573	ALA	2.4
1	A	609	LEU	2.4
1	A	459	THR	2.3
1	A	221	GLU	2.3
1	A	485	LEU	2.3
1	A	591	HIS	2.2
1	A	357	PHE	2.2
1	A	505	VAL	2.2
1	A	543	MET	2.2
1	A	329	ALA	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	613	PRO	2.1
1	A	512	VAL	2.0
1	A	390	ARG	2.0
1	A	582	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	BMA	A	1330	11/12	0.69	0.19	102,106,108,108	0
3	NAG	A	1420	14/15	0.74	0.53	115,123,127,128	0
3	NAG	A	1338	14/15	0.74	0.39	121,126,128,129	0
5	MAN	A	1035	11/12	0.81	0.23	102,106,108,109	0
3	NAG	A	1389	14/15	0.82	0.28	138,140,142,144	0
5	MAN	A	1036	11/12	0.83	0.28	104,106,108,108	0
3	NAG	A	1337	14/15	0.83	0.39	124,129,130,130	0
5	MAN	A	1508	11/12	0.85	0.40	112,116,119,119	0
5	MAN	A	1507	11/12	0.85	0.19	105,106,109,109	0
6	FUC	A	1509	10/11	0.90	0.28	100,102,102,103	0
4	BMA	A	1034	11/12	0.91	0.22	97,99,102,106	0
3	NAG	A	1033	14/15	0.93	0.25	81,89,94,97	0
3	NAG	A	1329	14/15	0.94	0.24	82,89,96,96	0
4	BMA	A	1506	11/12	0.94	0.14	93,96,102,107	0
3	NAG	A	1032	14/15	0.94	0.17	31,53,70,70	0
3	NAG	A	1504	14/15	0.95	0.19	74,84,93,97	0
6	FUC	A	1037	10/11	0.96	0.15	66,71,75,76	0
3	NAG	A	1328	14/15	0.97	0.20	65,72,76,78	0
3	NAG	A	1505	14/15	0.97	0.24	76,80,83,87	0

6.5 Other polymers [i](#)

There are no such residues in this entry.