



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 7, 2018 – 03:51 am GMT

PDB ID : 3R08
Title : Crystal structure of mouse cd3epsilon in complex with antibody 2C11 Fab
Authors : Shore, D.A.; Zhu, X.; Wilson, I.A.
Deposited on : 2011-03-07
Resolution : 4.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : trunk30967
Percentile statistics : (not set)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30967


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 4.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	1039 (4.54-3.66)
Clashscore	122126	1052 (4.50-3.70)
Ramachandran outliers	120053	1006 (4.50-3.70)
Sidechain outliers	120020	1005 (4.52-3.68)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	L	213	
2	H	216	
3	E	82	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3944 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Mouse anti-mouse CD3epsilon antibody 2C11 light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	L	213	1664	1041	278	340	5	0	0	0

- Molecule 2 is a protein called Mouse anti-mouse CD3epsilon antibody 2C11 heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	H	216	1626	1029	268	320	9	0	0	0

- Molecule 3 is a protein called T-cell surface glycoprotein CD3 epsilon chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	E	82	654	410	105	137	2	0	0	0


There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	80	SER	-	EXPRESSION TAG	UNP P22646
E	81	GLU	-	EXPRESSION TAG	UNP P22646
E	82	TYR	-	EXPRESSION TAG	UNP P22646


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Mouse anti-mouse CD3epsilon antibody 2C11 light chain



- Molecule 2: Mouse anti-mouse CD3epsilon antibody 2C11 heavy chain



- Molecule 3: T-cell surface glycoprotein CD3 epsilon chain



4 Data and refinement statistics

Property	Value	Source
Space group	I 41 3 2	Depositor
Cell constants a, b, c, α , β , γ	263.18Å 263.18Å 263.18Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	33.15 – 4.10 43.86 – 4.10	Depositor EDS
% Data completeness (in resolution range)	(Not available) (33.15-4.10) 96.9 (43.86-4.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.56 (at 4.13Å)	Xtrriage
Refinement program	BUSTER 2.8.0	Depositor
R, R_{free}	0.222 , 0.277 0.242 , 0.293	Depositor DCC
R_{free} test set	584 reflections (4.82%)	wwPDB-VP
Wilson B-factor (Å ²)	158.3	Xtrriage
Anisotropy	0.000	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 185.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.36$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.86	EDS
Total number of atoms	3944	wwPDB-VP
Average B, all atoms (Å ²)	180.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.64% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	L	0.54	0/1701	0.85	1/2309 (0.0%)
2	H	0.51	0/1666	0.83	0/2274
3	E	0.64	1/667 (0.1%)	0.98	0/904
All	All	0.54	1/4034 (0.0%)	0.87	1/5487 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	82	TYR	CB-CG	-6.09	1.42	1.51

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	L	40	PRO	N-CA-C	5.51	126.43	112.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	1664	0	1608	52	0
2	H	1626	0	1593	41	0
3	E	654	0	612	13	0
All	All	3944	0	3813	102	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (102) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:156:ILE:HG13	2:H:209:ASN:HB2	1.62	0.82
2:H:151:PRO:HD2	2:H:213:PRO:HG3	1.66	0.77
2:H:119:PRO:HB3	2:H:147:TYR:HB3	1.67	0.75
2:H:133:SER:HB2	2:H:136:ASN:HA	1.70	0.74
3:E:9:LYS:HB3	3:E:20:THR:HB	1.72	0.72
1:L:34:ASN:HD22	1:L:49:TYR:HA	1.56	0.71
2:H:171:VAL:HG22	2:H:191:VAL:HG23	1.71	0.71
1:L:115:VAL:HG21	1:L:196:VAL:HG21	1.74	0.68
2:H:206:VAL:H	2:H:222:LYS:HE3	1.59	0.68
1:L:90:GLN:HE21	1:L:97:THR:H	1.47	0.62
1:L:189:HIS:O	1:L:211:ARG:NH1	2.32	0.62
1:L:4:MET:HB2	1:L:99:GLY:HA2	1.82	0.62
1:L:120:PRO:HD3	1:L:132:LEU:HG	1.80	0.61
1:L:193:THR:HG23	1:L:208:THR:HG23	1.81	0.61
1:L:52:ASN:H	1:L:52:ASN:HD22	1.48	0.60
2:H:68:THR:HG23	2:H:81:GLN:HB3	1.84	0.60
2:H:5:VAL:HG23	2:H:23:GLU:HB2	1.84	0.60
2:H:87:THR:HG23	2:H:110:THR:HA	1.84	0.59
1:L:83:ILE:HG12	1:L:106:ILE:HB	1.86	0.58
1:L:66:GLY:HA3	1:L:71:SER:HA	1.85	0.58
1:L:15:LEU:HD11	1:L:106:ILE:HD11	1.86	0.57
1:L:36:TYR:OH	1:L:89:GLN:NE2	2.38	0.57
1:L:195:GLU:HA	1:L:206:VAL:HG12	1.86	0.56
3:E:7:GLU:HA	3:E:22:PRO:HB3	1.86	0.56
1:L:115:VAL:CG2	1:L:196:VAL:HG21	2.34	0.56
2:H:66:ARG:HD3	2:H:82(B):ILE:O	2.04	0.56
1:L:2:ILE:HG22	1:L:3:GLN:N	2.20	0.56
1:L:21:ILE:HG12	1:L:102:THR:HG21	1.89	0.55
1:L:95:PRO:HG3	2:H:60:ALA:HA	1.89	0.55
2:H:12:VAL:HG11	2:H:82(C):LEU:HD13	1.89	0.54
2:H:212:HIS:HE1	2:H:214:ALA:HB3	1.71	0.54
2:H:29:PHE:HB3	2:H:76:ASN:HD22	1.72	0.54
3:E:29:LEU:HD11	3:E:62:CYS:SG	2.47	0.53
1:L:37:GLN:HB2	1:L:47:LEU:HD21	1.89	0.53
1:L:6:GLN:HB2	1:L:100:PRO:HD2	1.88	0.53
1:L:34:ASN:ND2	1:L:49:TYR:HA	2.23	0.53
3:E:75:LEU:O	3:E:76:LYS:HG2	2.09	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:137:ASN:HD21	2:H:174:PHE:HZ	1.57	0.53
1:L:136:VAL:HG23	1:L:175:LEU:HB3	1.90	0.52
1:L:42:LYS:HE3	1:L:42:LYS:H	1.74	0.52
1:L:146:VAL:HG12	1:L:196:VAL:HA	1.91	0.52
2:H:133:SER:CB	2:H:136:ASN:HA	2.40	0.52
2:H:66:ARG:HG3	2:H:67:PHE:CD1	2.46	0.52
2:H:31:GLY:HA2	2:H:52(A):SER:HB2	1.92	0.51
2:H:66:ARG:HG3	2:H:67:PHE:HD1	1.74	0.51
2:H:212:HIS:CE1	2:H:214:ALA:HB3	2.46	0.51
1:L:6:GLN:NE2	1:L:102:THR:H	2.09	0.50
2:H:11:LEU:HG	2:H:12:VAL:H	1.77	0.50
1:L:19:VAL:HG11	1:L:104:LEU:HD11	1.93	0.50
1:L:2:ILE:HG22	1:L:3:GLN:H	1.75	0.50
2:H:75:LYS:HB2	2:H:77:LEU:HD12	1.94	0.48
3:E:53:GLU:HG2	3:E:76:LYS:HB2	1.95	0.48
1:L:35:TRP:CZ3	1:L:88:CYS:HB3	2.49	0.48
1:L:5:THR:HB	1:L:24:GLN:HB3	1.95	0.47
1:L:37:GLN:HB2	1:L:47:LEU:HD11	1.97	0.47
1:L:150:VAL:HG22	1:L:192:TYR:HD1	1.80	0.46
2:H:157:TRP:CH2	2:H:208:CYS:HB2	2.51	0.46
3:E:67:SER:C	3:E:69:LYS:H	2.19	0.46
2:H:120:SER:HB3	2:H:122:TYR:CZ	2.51	0.45
2:H:82(A):ASN:HB2	2:H:82(B):ILE:HD12	1.98	0.45
2:H:142:CYS:HB2	2:H:157:TRP:CZ2	2.52	0.45
1:L:43:ALA:HA	1:L:44:PRO:HD3	1.88	0.45
1:L:48:ILE:HG22	1:L:53:LYS:O	2.16	0.45
3:E:21:CYS:HB2	3:E:31:TRP:CZ2	2.52	0.45
1:L:48:ILE:HD12	1:L:64:GLY:N	2.32	0.45
1:L:131:THR:HG23	1:L:180:SER:HA	1.99	0.44
1:L:150:VAL:HG22	1:L:192:TYR:CD1	2.52	0.44
3:E:21:CYS:SG	3:E:22:PRO:HD2	2.58	0.44
1:L:49:TYR:HE1	1:L:55:ALA:HA	1.82	0.44
2:H:4:LEU:HD23	2:H:24:ALA:HA	1.99	0.44
2:H:59:TYR:HE1	2:H:69:VAL:HG12	1.81	0.44
3:E:12:ILE:HG13	3:E:17:VAL:HG13	2.00	0.44
2:H:120:SER:HB2	2:H:145:LYS:HB3	1.99	0.44
1:L:209:LEU:HG	2:H:128:CYS:HB2	2.00	0.44
2:H:121:VAL:HB	2:H:219:VAL:HG11	1.99	0.43
1:L:150:VAL:HG13	1:L:192:TYR:CE1	2.53	0.43
1:L:196:VAL:HB	1:L:205:ILE:O	2.19	0.43
2:H:200:TRP:CD1	2:H:202:PRO:HA	2.54	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:140:LEU:HD22	2:H:223:ILE:HD13	2.01	0.43
3:E:46:LEU:HD23	3:E:48:LEU:HD11	1.99	0.43
2:H:156:ILE:HD12	2:H:163:SER:HA	2.01	0.42
1:L:117:ILE:HD13	1:L:194:CYS:HB2	2.02	0.42
1:L:29:ILE:HD12	1:L:31:ASN:H	1.84	0.42
3:E:57:SER:OG	3:E:76:LYS:HB3	2.20	0.42
2:H:133:SER:HB2	2:H:136:ASN:CA	2.44	0.41
2:H:200:TRP:CG	2:H:202:PRO:HA	2.55	0.41
1:L:61:ARG:HB2	1:L:76:SER:HB3	2.02	0.41
3:E:4:GLU:HB3	3:E:5:ASN:H	1.69	0.41
1:L:2:ILE:CG2	1:L:3:GLN:N	2.84	0.41
1:L:125:LEU:O	1:L:183:LYS:HD2	2.21	0.41
1:L:52:ASN:H	1:L:52:ASN:ND2	2.15	0.41
2:H:157:TRP:CZ3	2:H:208:CYS:HB2	2.55	0.41
1:L:160:LEU:HD21	2:H:179:HIS:HB2	2.01	0.41
2:H:137:THR:HA	2:H:194:PRO:HA	2.03	0.41
1:L:191:LEU:HD12	1:L:210:ASN:HB2	2.02	0.41
3:E:31:TRP:HB3	3:E:38:LEU:HD12	2.03	0.40
1:L:193:THR:HG22	1:L:206:VAL:HB	2.04	0.40
2:H:199:THR:HA	2:H:204:GLN:NE2	2.37	0.40
1:L:48:ILE:HD11	1:L:73:PHE:HE1	1.87	0.40
2:H:101:ASN:HD22	2:H:101:ASN:HA	1.52	0.40
1:L:207:LYS:HA	1:L:207:LYS:HD2	2.01	0.40
1:L:2:ILE:CG2	1:L:3:GLN:H	2.34	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	L	211/213 (99%)	179 (85%)	22 (10%)	10 (5%)	2 27

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	H	214/216 (99%)	180 (84%)	27 (13%)	7 (3%)	4	35
3	E	80/82 (98%)	61 (76%)	11 (14%)	8 (10%)	0	11
All	All	505/511 (99%)	420 (83%)	60 (12%)	25 (5%)	2	26

All (25) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	L	18	ARG
1	L	40	PRO
1	L	43	ALA
1	L	51	THR
2	H	132	THR
2	H	136	ASN
3	E	26	ASP
1	L	60	SER
1	L	80	SER
1	L	211	ARG
2	H	146	GLY
1	L	26	SER
1	L	32	TYR
1	L	100	PRO
2	H	28	THR
3	E	6	ILE
3	E	77	ALA
2	H	131	THR
3	E	34	ASN
3	E	66	ALA
2	H	213	PRO
3	E	4	GLU
3	E	23	LEU
3	E	40	GLN
2	H	8	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	190/190 (100%)	153 (80%)	37 (20%)	1	10
2	H	184/184 (100%)	154 (84%)	30 (16%)	2	17
3	E	73/76 (96%)	61 (84%)	12 (16%)	2	17
All	All	447/450 (99%)	368 (82%)	79 (18%)	2	14

All (79) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	3	GLN
1	L	7	SER
1	L	9	SER
1	L	14	SER
1	L	17	ASP
1	L	20	THR
1	L	33	LEU
1	L	38	GLN
1	L	39	LYS
1	L	42	LYS
1	L	48	ILE
1	L	51	THR
1	L	52	ASN
1	L	72	SER
1	L	76	SER
1	L	80	SER
1	L	89	GLN
1	L	93	ASN
1	L	96	TRP
1	L	112	LYS
1	L	114	THR
1	L	115	VAL
1	L	129	SER
1	L	132	LEU
1	L	133	VAL
1	L	136	VAL
1	L	142	LYS
1	L	145	ASN
1	L	149	LYS
1	L	154	GLU
1	L	176	SER
1	L	178	THR
1	L	191	LEU
1	L	193	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	L	198	HIS
1	L	208	THR
1	L	209	LEU
2	H	3	GLN
2	H	5	VAL
2	H	11	LEU
2	H	12	VAL
2	H	18	LEU
2	H	45	LEU
2	H	54	SER
2	H	79	PHE
2	H	85	GLU
2	H	92	CYS
2	H	98	ASP
2	H	101	ASN
2	H	107	THR
2	H	108	MET
2	H	110	THR
2	H	134	THR
2	H	137	THR
2	H	139	THR
2	H	143	LEU
2	H	167	THR
2	H	168	SER
2	H	177	VAL
2	H	188	SER
2	H	192	THR
2	H	193	VAL
2	H	196	SER
2	H	203	SER
2	H	209	ASN
2	H	222	LYS
2	H	223	ILE
3	E	1	ASP
3	E	5	ASN
3	E	11	SER
3	E	19	LEU
3	E	21	CYS
3	E	25	SER
3	E	26	ASP
3	E	34	ASN
3	E	37	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	E	53	GLU
3	E	54	VAL
3	E	75	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	L	6	GLN
1	L	52	ASN
1	L	89	GLN
1	L	138	ASN
2	H	76	ASN
2	H	101	ASN
2	H	105	GLN
2	H	204	GLN
3	E	70	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers

Unable to reproduce the depositors R factor - this section is therefore empty.