



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 2, 2021 – 10:48 AM EST

PDB ID : 5R0Y
Title : PanDDA analysis group deposition – Auto-refined data of Aar2/RNaseH for ground state model 12, DMSO-free
Authors : Wollenhaupt, J.; Metz, A.; Barthel, T.; Lima, G.M.A.; Heine, A.; Mueller, U.; Klebe, G.; Weiss, M.S.
Deposited on : 2020-02-12
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.17.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.17.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2340 (1.76-1.76)
Clashscore	141614	2466 (1.76-1.76)
Ramachandran outliers	138981	2437 (1.76-1.76)
Sidechain outliers	138945	2437 (1.76-1.76)
RSRZ outliers	127900	2298 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	258	 27% 82% 9% 8%
2	B	308	 25% 85% 12% 8%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4681 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pre-mRNA-splicing factor 8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	237	2002	1283	335	372	12	0	12	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1833	GLY	-	expression tag	UNP P33334
A	1834	ALA	-	expression tag	UNP P33334
A	1835	MET	-	expression tag	UNP P33334

- Molecule 2 is a protein called A1 cistron-splicing factor AAR2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	300	2580	1654	421	485	20	0	9	0

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-3	GLY	-	expression tag	UNP P32357
B	-2	ALA	-	expression tag	UNP P32357
B	-1	MET	-	expression tag	UNP P32357
B	0	ALA	-	expression tag	UNP P32357
B	166	SER	LEU	conflict	UNP P32357
B	167	SER	LYS	conflict	UNP P32357
B	170	SER	LEU	conflict	UNP P32357
B	?	-	GLN	deletion	UNP P32357
B	?	-	LYS	deletion	UNP P32357
B	?	-	ALA	deletion	UNP P32357
B	?	-	GLY	deletion	UNP P32357
B	?	-	SER	deletion	UNP P32357
B	?	-	LYS	deletion	UNP P32357

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	?	-	MET	deletion	UNP P32357
B	?	-	GLU	deletion	UNP P32357
B	?	-	ALA	deletion	UNP P32357
B	?	-	LYS	deletion	UNP P32357
B	?	-	ASN	deletion	UNP P32357
B	?	-	GLU	deletion	UNP P32357
B	?	-	ASP	deletion	UNP P32357


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	61	Total O 61 61	0	0
3	B	38	Total O 38 38	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Pre-mRNA-splicing factor 8



- Molecule 2: A1 cistron-splicing factor AAR2



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	88.37Å 82.11Å 92.30Å 90.00° 107.71° 90.00°	Depositor
Resolution (Å)	23.29 – 1.75 44.68 – 1.75	Depositor EDS
% Data completeness (in resolution range)	99.6 (23.29-1.75) 99.7 (44.68-1.75)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.97 (at 1.75Å)	Xtrriage
Refinement program	REFMAC 5.8.0238	Depositor
R, R_{free}	0.286 , 0.342 0.293 , 0.348	Depositor DCC
R_{free} test set	2097 reflections (3.31%)	wwPDB-VP
Wilson B-factor (Å ²)	42.4	Xtrriage
Anisotropy	0.273	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 55.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4681	wwPDB-VP
Average B, all atoms (Å ²)	81.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.72% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.72	0/2049	0.95	6/2775 (0.2%)
2	B	0.67	0/2651	0.77	1/3581 (0.0%)
All	All	0.69	0/4700	0.86	7/6356 (0.1%)

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1859	ARG	NE-CZ-NH1	-8.71	115.94	120.30
1	A	1908	LEU	CB-CG-CD1	-8.04	97.34	111.00
1	A	1908	LEU	CB-CG-CD2	6.98	122.86	111.00
1	A	1859	ARG	NE-CZ-NH2	6.91	123.76	120.30
1	A	1944	LEU	CB-CG-CD2	-6.24	100.39	111.00
2	B	141	ASP	CB-CG-OD1	6.12	123.81	118.30
1	A	2011	LEU	CB-CG-CD1	-6.08	100.66	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2002	0	2029	12	0
2	B	2580	0	2450	25	0
3	A	61	0	0	0	0
3	B	38	0	0	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	4681	0	4479	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:2062:GLU:O	1:A:2066:LYS:HG2	1.87	0.74
2:B:1:MET:N	3:B:401:HOH:O	2.21	0.72
2:B:258:LYS:HD2	2:B:258:LYS:H	1.62	0.64
2:B:230[B]:ASN:ND2	2:B:239:SER:OG	2.39	0.56
1:A:1848:ILE:H	1:A:1931[A]:LYS:HZ2	1.52	0.56
2:B:287:ARG:O	2:B:291:ILE:HD13	2.07	0.55
2:B:16:GLY:HA3	2:B:45:HIS:CE1	2.44	0.53
2:B:1:MET:HB3	2:B:35:ASP:HA	1.91	0.52
2:B:242:GLN:O	2:B:246:MET:HG3	2.12	0.50
1:A:2062:GLU:HB3	1:A:2066:LYS:HE3	1.94	0.49
2:B:214:PHE:O	2:B:215:LYS:HB2	2.13	0.48
2:B:251:CYS:O	2:B:296:SER:HB2	2.13	0.48
2:B:70:GLN:HB3	2:B:81:MET:HE1	1.96	0.48
1:A:1913:THR:HB	1:A:1940:MET:HE1	1.96	0.47
1:A:1946:VAL:O	1:A:1949:LEU:HG	2.15	0.47
2:B:6:PHE:CD1	2:B:32:GLY:HA2	2.49	0.47
2:B:6:PHE:CZ	2:B:44[A]:ILE:HD11	2.49	0.47
2:B:54[B]:MET:N	3:B:402:HOH:O	2.41	0.46
1:A:1962:ARG:O	1:A:2013:ARG:NH1	2.44	0.46
2:B:243:TRP:O	2:B:247:ILE:HG13	2.16	0.46
2:B:6:PHE:HZ	2:B:44[A]:ILE:HD11	1.80	0.45
2:B:114:TRP:CE2	2:B:118:THR:HG21	2.54	0.43
2:B:229:LEU:HD23	2:B:229:LEU:HA	1.89	0.43
1:A:2066:LYS:HB2	1:A:2067:TYR:CD1	2.53	0.43
2:B:114:TRP:CZ2	2:B:118:THR:HG21	2.55	0.42
2:B:244:HIS:CD2	2:B:285:ASN:HB3	2.54	0.42
2:B:188:ALA:HA	2:B:204:TYR:CD1	2.54	0.42
2:B:42:HIS:HE1	3:B:429:HOH:O	2.02	0.42
2:B:53:SER:HA	3:B:402:HOH:O	2.20	0.41
2:B:77:LEU:HD21	2:B:79:LYS:HE3	2.03	0.41
1:A:1942:ASP:HB2	1:A:1943:PRO:HD3	2.02	0.41
1:A:1944:LEU:HD12	1:A:1944:LEU:HA	1.90	0.41
1:A:2050:THR:HG22	1:A:2054:GLN:NE2	2.36	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:134:GLU:OE1	2:B:134:GLU:N	2.47	0.40
1:A:1843:LEU:HA	1:A:1849:LYS:HD2	2.04	0.40
1:A:1917:VAL:O	1:A:1920:LEU:HB3	2.21	0.40
2:B:15:ILE:HD12	2:B:24:VAL:HG21	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	248/258 (96%)	244 (98%)	4 (2%)	0	100	100
2	B	306/308 (99%)	293 (96%)	11 (4%)	2 (1%)	22	8
All	All	554/566 (98%)	537 (97%)	15 (3%)	2 (0%)	47	17

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	54[A]	MET
2	B	54[B]	MET

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	226/233 (97%)	218 (96%)	8 (4%)	36	13
2	B	287/284 (101%)	285 (99%)	2 (1%)	84	75
All	All	513/517 (99%)	503 (98%)	10 (2%)	67	37

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1838	SER
1	A	1903	LYS
1	A	1931[A]	LYS
1	A	1931[B]	LYS
1	A	1962	ARG
1	A	1979[A]	MET
1	A	1979[B]	MET
1	A	1979[C]	MET
2	B	77	LEU
2	B	84	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1907	GLN
1	A	2038	HIS
2	B	47	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	237/258 (91%)	1.55	70 (29%) 0 0	41, 71, 119, 192	0
2	B	300/308 (97%)	1.58	76 (25%) 0 0	40, 74, 152, 288	0
All	All	537/566 (94%)	1.57	146 (27%) 0 0	40, 72, 141, 288	0

All (146) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	108	ILE	16.2
1	A	1878	CYS	9.7
2	B	109	ASP	9.0
2	B	53	SER	8.7
2	B	1	MET	8.3
2	B	52	SER	6.9
2	B	87	ALA	6.1
2	B	99	ARG	5.8
2	B	174	HIS	5.7
2	B	96	PHE	5.6
1	A	1905	LEU	5.6
1	A	2048	TRP	5.2
1	A	1833	GLY	5.0
2	B	181	ILE	5.0
1	A	1963	LEU	5.0
2	B	313	TYR	4.9
1	A	2027	LEU	4.8
2	B	54[A]	MET	4.8
1	A	1956	ILE	4.7
1	A	1991	ILE	4.6
2	B	277	GLU	4.6
1	A	2034	ILE	4.4
1	A	1933	ILE	4.4
2	B	89	PHE	4.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	175	SER	4.2
2	B	193	HIS	4.1
2	B	171	ASP	4.0
1	A	1962	ARG	3.9
1	A	1838	SER	3.9
2	B	208	VAL	3.8
2	B	172	PRO	3.8
2	B	278	GLN	3.8
1	A	2063	TYR	3.8
2	B	110	GLU	3.8
2	B	107	LYS	3.8
1	A	1927	GLU	3.7
1	A	1866	PHE	3.7
1	A	2060	LEU	3.7
1	A	1834	ALA	3.7
1	A	1917	VAL	3.7
2	B	281	ASP	3.6
2	B	101	MET	3.5
2	B	316	LEU	3.5
1	A	2066	LYS	3.5
2	B	73	PRO	3.5
1	A	1891	LEU	3.5
2	B	279	TYR	3.4
2	B	150	ASN	3.4
1	A	1882	LEU	3.4
1	A	1954	ILE	3.3
1	A	1865	THR	3.3
2	B	20	TYR	3.3
1	A	1934	ILE	3.3
1	A	1940	MET	3.3
2	B	148	GLN	3.3
1	A	1930	PRO	3.2
1	A	2040	TRP	3.2
1	A	1863	HIS	3.2
1	A	1862	VAL	3.2
2	B	77	LEU	3.2
1	A	1944	LEU	3.1
1	A	2058	LEU	3.1
2	B	22	PHE	3.1
1	A	1935	VAL	3.1
2	B	228	PHE	3.0
1	A	2024[A]	MET	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	275	LEU	3.0
1	A	2003	THR	3.0
1	A	1888	HIS	2.9
2	B	291	ILE	2.9
1	A	2028	SER	2.9
1	A	2017[A]	THR	2.9
2	B	69	ILE	2.9
2	B	84	ARG	2.8
1	A	1988	LEU	2.8
2	B	210	LEU	2.8
2	B	317	LEU	2.8
2	B	28	GLN	2.8
1	A	1989	PHE	2.8
2	B	106	PRO	2.8
1	A	1971	ILE	2.7
1	A	2039	LEU	2.7
2	B	100	GLN	2.7
2	B	103	VAL	2.7
1	A	1998	ARG	2.7
2	B	94	HIS	2.6
2	B	40	HIS	2.6
1	A	2002	TYR	2.6
1	A	2008	LEU	2.6
1	A	2046	GLU	2.6
1	A	2049	ILE	2.6
2	B	147	VAL	2.6
1	A	2026	LEU	2.6
1	A	2050	THR	2.5
2	B	170	SER	2.5
2	B	189	ILE	2.5
1	A	1946	VAL	2.5
2	B	152	LEU	2.5
2	B	26	GLU	2.5
1	A	1861	THR	2.5
1	A	1996	LEU	2.5
2	B	86	GLY	2.5
1	A	1953	ASN	2.5
2	B	146	THR	2.5
2	B	92	ILE	2.5
1	A	1836	ASN	2.4
1	A	1884	PRO	2.4
1	A	1840	TYR	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	129	ILE	2.4
2	B	247	ILE	2.4
2	B	30	PHE	2.4
2	B	205	LEU	2.4
2	B	295	SER	2.4
2	B	254	ALA	2.4
2	B	97	LYS	2.4
2	B	249	LEU	2.4
1	A	2044	THR	2.4
1	A	1839	ASN	2.3
1	A	1910	LYS	2.3
2	B	213	ILE	2.3
1	A	1835	MET	2.3
2	B	133	ASP	2.3
1	A	1870	VAL	2.3
2	B	6	PHE	2.3
2	B	294	TYR	2.3
1	A	1850	LEU	2.2
2	B	124	ASP	2.2
1	A	1980	LYS	2.2
1	A	2007	ARG	2.2
1	A	1979[A]	MET	2.2
1	A	2012	LEU	2.2
2	B	136	GLN	2.2
1	A	2018	ASN	2.2
1	A	1871	ALA	2.2
1	A	2015	LEU	2.2
2	B	91	ASN	2.1
2	B	182	ASN	2.1
1	A	1879	ILE	2.1
2	B	284	LEU	2.1
2	B	31	HIS	2.1
1	A	2059	ILE	2.1
2	B	36	ILE	2.1
2	B	226	PHE	2.1
2	B	132	LYS	2.0
2	B	212	GLY	2.0
2	B	198	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.