

wwPDB X-ray Structure Validation Summary Report (i)

Sep 2, 2023 – 11:22 AM EDT

PDB ID	:	3R1B
Title	:	Open crystal structure of cytochrome P450 2B4 covalently bound to the
		mechanism-based inactivator tert-butylphenylacetylene
Authors	:	Gay, S.C.; Zhang, H.; Stout, C.D.; Hollenberg, P.F.; Halpert, J.R.
Deposited on	:	2011-03-09
Resolution	:	3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.35
buster-report	:	1.1.7(2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY\;DIFFRACTION$

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R _{free}	130704	2092 (3.00-3.00)
Clashscore	141614	2416 (3.00-3.00)
Ramachandran outliers	138981	2333 (3.00-3.00)
Sidechain outliers	138945	2336 (3.00-3.00)
RSRZ outliers	127900	1990 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality	of chain	
1	Δ	476	.% •	240/	60/
1	11	410	2%	54%	0% •
1	В	476	53%	38%	7% •
1	С	476	5% 49%	39%	8% •
1	D	476	42%	46%	9% •
2	Е	2	10	00%	



Mol	Chain	Length	Quality of chain
2	F	2	100%
2	G	2	100%
2	Н	2	100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GLC	F	1	-	-	Х	-
2	FRU	F	2	-	-	Х	-
2	GLC	Н	1	-	-	Х	-
2	FRU	Н	2	-	-	Х	-
3	HEM	D	500	-	-	Х	-
4	TB2	А	501	-	-	Х	-
4	TB2	С	501	-	-	Х	-
4	TB2	D	501	-	-	Х	-
5	CM5	А	608	-	-	-	Х



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 14499 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace				
1	1 1	465	Total	С	Ν	0	\mathbf{S}	2	0	0				
	A	405	3583	2294	619	659	11	2						
1	1 B	Р	P	В	В	465	Total	С	Ν	0	S	0	0	0
1		405	3589	2308	615	656	10	2	0	0				
1	1 0	C 456	Total	С	Ν	0	S	1	0	0				
	450	3406	2171	589	638	8	1	0						
1 D	464	Total	С	Ν	0	S	0	0	0					
	404	3445	2199	598	640	8	0		0					

• Molecule 1 is a protein called Cytochrome P450 2B4.

There are 132 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	21	ALA	GLU	engineered mutation	UNP P00178
А	?	-	PHE	deletion	UNP P00178
А	?	-	SER	deletion	UNP P00178
А	?	-	LEU	deletion	UNP P00178
А	?	-	LEU	deletion	UNP P00178
А	?	-	LEU	deletion	UNP P00178
А	?	-	LEU	deletion	UNP P00178
А	?	-	LEU	deletion	UNP P00178
А	?	-	ALA	deletion	UNP P00178
А	?	-	PHE	deletion	UNP P00178
А	?	-	LEU	deletion	UNP P00178
А	?	-	ALA	deletion	UNP P00178
А	?	-	GLY	deletion	UNP P00178
А	?	-	LEU	deletion	UNP P00178
A	?	-	LEU	deletion	UNP P00178
А	?	-	LEU	deletion	UNP P00178
А	?	-	LEU	deletion	UNP P00178
A	?	-	LEU	deletion	UNP P00178
A	?	-	PHE	deletion	UNP P00178
А	?	-	ARG	deletion	UNP P00178
А	22	LYS	GLY	engineered mutation	UNP P00178



Chain	Residue	Modelled	Actual	Comment	Reference
А	23	LYS	HIS	engineered mutation	UNP P00178
А	24	THR	PRO	engineered mutation	UNP P00178
А	25	SER	LYS	engineered mutation	UNP P00178
А	26	SER	ALA	engineered mutation	UNP P00178
А	27	LYS	HIS	engineered mutation	UNP P00178
А	29	LYS	ARG	engineered mutation	UNP P00178
А	221	SER	PRO	conflict	UNP P00178
А	226	TYR	HIS	engineered mutation	UNP P00178
А	492	HIS	-	expression tag	UNP P00178
А	493	HIS	-	expression tag	UNP P00178
А	494	HIS	-	expression tag	UNP P00178
А	495	HIS	-	expression tag	UNP P00178
В	21	ALA	GLU	engineered mutation	UNP P00178
В	?	-	PHE	deletion	UNP P00178
В	?	-	SER	deletion	UNP P00178
В	?	-	LEU	deletion	UNP P00178
В	?	-	LEU	deletion	UNP P00178
В	?	-	LEU	deletion	UNP P00178
В	?	-	LEU	deletion	UNP P00178
В	?	-	LEU	deletion	UNP P00178
В	?	-	ALA	deletion	UNP P00178
В	?	-	PHE	deletion	UNP P00178
В	?	-	LEU	deletion	UNP P00178
В	?	-	ALA	deletion	UNP P00178
В	?	-	GLY	deletion	UNP P00178
В	?	_	LEU	deletion	UNP P00178
В	?	_	LEU	deletion	UNP P00178
В	?	-	LEU	deletion	UNP P00178
В	?	_	LEU	deletion	UNP P00178
В	?	_	LEU	deletion	UNP P00178
В	?	-	PHE	deletion	UNP P00178
В	?	-	ARG	deletion	UNP P00178
B	22	LYS	GLY	engineered mutation	UNP P00178
В	23	LYS	HIS	engineered mutation	UNP P00178
В	24	THR	PRO	engineered mutation	UNP P00178
B	25	SER	LYS	engineered mutation	UNP P00178
В	26	SER	ALA	engineered mutation	UNP P00178
B	27	LYS	HIS	engineered mutation	UNP P00178
B	29	LYS	ARG	engineered mutation	UNP P00178
B	221	SER	PRO	conflict	UNP P00178
B	226	TYR	HIS	engineered mutation	UNP P00178
В	492	HIS	_	expression tag	UNP P00178



3R1	В

Chain	Residue	Modelled	Actual	Comment	Reference
B	493	HIS	-	expression tag	UNP P00178
B	494	HIS		expression tag	UNP P00178
B	495	HIS		expression tag	UNP P00178
C	21		GLU	engineered mutation	UNP P00178
	?	-	PHE	deletion	UNP P00178
C	· ?		SEB	deletion	UNP P00178
C	· · · · · · · · · · · · · · · · · · ·		LEU	deletion	UNP P00178
C	· ?		LEU	deletion	UNP P00178
C	· ?		LEU	deletion	UNP P00178
C	· ?		LEU	deletion	UNP P00178
C	· ?		LEU	deletion	UNP P00178
C	· ?		ALA	deletion	UNP P00178
C	?		PHE	deletion	UNP P00178
C	?		LEU	deletion	UNP P00178
C	· ?		ALA	deletion	UNP P00178
<u> </u>	?		GLY	deletion	UNP P00178
C	· ?		LEU	deletion	UNP P00178
C	?		LEU	deletion	UNP P00178
C	?		LEU	deletion	UNP P00178
C	?		LEU	deletion	UNP P00178
C	?	_	LEU	deletion	UNP P00178
C	?	_	PHE	deletion	UNP P00178
C	?	_	ARG	deletion	UNP P00178
C	22	LYS	GLY	engineered mutation	UNP P00178
C	23	LYS	HIS	engineered mutation	UNP P00178
С	24	THR	PRO	engineered mutation	UNP P00178
С	25	SER	LYS	engineered mutation	UNP P00178
С	26	SER	ALA	engineered mutation	UNP P00178
С	27	LYS	HIS	engineered mutation	UNP P00178
С	29	LYS	ARG	engineered mutation	UNP P00178
С	221	SER	PRO	conflict	UNP P00178
С	226	TYR	HIS	engineered mutation	UNP P00178
С	492	HIS	-	expression tag	UNP P00178
С	493	HIS	-	expression tag	UNP P00178
С	494	HIS	-	expression tag	UNP P00178
С	495	HIS	-	expression tag	UNP P00178
D	21	ALA	GLU	engineered mutation	UNP P00178
D	?	-	PHE	deletion	UNP P00178
D	?	-	SER	deletion	UNP P00178
D	?	-	LEU	deletion	UNP P00178
D	?	-	LEU	deletion	UNP P00178
D	?	-	LEU	deletion	UNP P00178



Chain	Residue	Modelled	Actual	Comment	Reference
D	?	-	LEU	deletion	UNP P00178
D	?	-	LEU	deletion	UNP P00178
D	?	-	ALA	deletion	UNP P00178
D	?	-	PHE	deletion	UNP P00178
D	?	-	LEU	deletion	UNP P00178
D	?	-	ALA	deletion	UNP P00178
D	?	-	GLY	deletion	UNP P00178
D	?	-	LEU	deletion	UNP P00178
D	?	-	LEU	deletion	UNP P00178
D	?	-	LEU	deletion	UNP P00178
D	?	-	LEU	deletion	UNP P00178
D	?	-	LEU	deletion	UNP P00178
D	?	-	PHE	deletion	UNP P00178
D	?	-	ARG	deletion	UNP P00178
D	22	LYS	GLY	engineered mutation	UNP P00178
D	23	LYS	HIS	engineered mutation	UNP P00178
D	24	THR	PRO	engineered mutation	UNP P00178
D	25	SER	LYS	engineered mutation	UNP P00178
D	26	SER	ALA	engineered mutation	UNP P00178
D	27	LYS	HIS	engineered mutation	UNP P00178
D	29	LYS	ARG	engineered mutation	UNP P00178
D	221	SER	PRO	conflict	UNP P00178
D	226	TYR	HIS	engineered mutation	UNP P00178
D	492	HIS	-	expression tag	UNP P00178
D	493	HIS	-	expression tag	UNP P00178
D	494	HIS	-	expression tag	UNP P00178
D	495	HIS	-	expression tag	UNP P00178

• Molecule 2 is an oligosaccharide called beta-D-fructofuranose-(2-1)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	Trace
2	Е	2	Total (23 1	C O 12 11	0	0	0
2	F	2	Total (23 1	C O 12 11	0	0	0
2	G	2	Total (23 1	C O 12 11	0	0	0



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	Trace	
2	Н	2	Total 23	C 12	0 11	0	0	0

• Molecule 3 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
2	Λ	1	Total	С	Fe	Ν	0	0	0
5	Л	1	43	34	1	4	4	0	0
2	В	1	Total	С	Fe	Ν	0	0	0
5	D	1	43	34	1	4	4	0	0
2	C	C 1	Total	С	Fe	Ν	Ο	0	0
5	U	1	43	34	1	4	4	0	0
3 D	Л	1	Total	С	Fe	Ν	Ο	0	0
	D		43	34	1	4	4		U

• Molecule 4 is (4-tert-butylphenyl)acetaldehyde (three-letter code: TB2) (formula: $C_{12}H_{16}O$).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	1	Total C O 13 12 1	0	0
4	В	1	Total C O 13 12 1	0	0
4	С	1	Total C O 13 12 1	0	0
4	D	1	Total C O 13 12 1	0	0

• Molecule 5 is 5-CYCLOHEXYL-1-PENTYL-BETA-D-MALTOSIDE (three-letter code: CM5) (formula: $C_{23}H_{42}O_{11}$).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf						
5	Δ	1	Total C O	0	0						
5	Л	1	24 18 6	0	0						
5	Δ	1	Total C O	0	0						
0	11	1	34 23 11	0	0						
5	В	1	Total C O	0	0						
5	D	I	34 23 11	0	0						
5	В	1	Total C O	0	0						
5	D	I	34 23 11	0	0						
5	Л	1	Total C O	0	0						
б	D	D	D	D	D	D	D	1	34 23 11	0	U



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Cytochrome P450 2B4



1385 1386

7388 7388 73991 7395 7395 7395 7395 7395 7395 7395 7300 7300 7310 7300 7310 7300 7310 7300 7311 7300 7311 7300 7311 7312 7311 7312 7311 7312 7312 7313 7313 7313 7313 7326 7313 7335 7313 7335 7313 7335 7313 7335 7313 7335</







V324 V325 V326 V325 T328 F333 F333 F334 F335 F335 F336 F335 F336 F335 F336 F335 F344 F355 F365	
[392] [392] [395] [395] [396] [396] [396] [396] [396] [396] [396] [396] [396] [396] [396] [396] [396] [396] [396] [396] [396] [396] [396] [396] [396] [396] [396] [396] [396] [396] [396] [396] [397] [392] [398] [393] [393] [393] [393] [393] [393] [393] [393] [393] [393] [393] [393] [393] [393] [393] [393] [393] [394] [394] [394] [394] [394] [394] [394] [394] [394] [394] [394] <td></td>	
P462 P464 P465 P465 P465 P471 P472 P473 P474 P474 P475 P475 P475 P476 P478 P486 P486 P486 P487 P486 P486 P487 P486 P486 P487 P487 P486 P487 P487 P487 P486 P487 P487 P486 P486 P487 P487 P486 P486 P487 P487	
• Molecule 2: beta-D-fructofuranose-(2-1)-alpha-D-glucopyranose	
Chain E: 100%	
GLCI FRUZ	
• Molecule 2: beta-D-fructofuranose-(2-1)-alpha-D-glucopyranose	
Chain F: 100%	
GLC1 FRU2	
\bullet Molecule 2: beta-D-fructofuranose-(2-1)-alpha-D-glucopyranose	
Chain G: 100%	
GLC1	
• Molecule 2: beta-D-fructofuranose-(2-1)-alpha-D-glucopyranose	
Chain H: 100%	
FRU2	



4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants	193.92Å 153.69Å 129.66Å	Depositor
a, b, c, α , β , γ	90.00° 122.22° 90.00°	Depositor
Bosolution(A)	53.43 - 3.00	Depositor
Resolution (A)	53.44 - 3.00	EDS
% Data completeness	94.8 (53.43-3.00)	Depositor
(in resolution range)	99.3 (53.44-3.00)	EDS
R_{merge}	0.09	Depositor
R_{sym}	0.09	Depositor
$< I/\sigma(I) > 1$	$1.82 (at 3.01 \text{\AA})$	Xtriage
Refinement program	PHENIX 1.6.4_486	Depositor
P. P.	0.242 , 0.288	Depositor
n, n_{free}	0.241 , 0.281	DCC
R_{free} test set	3236 reflections $(5.06%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	92.5	Xtriage
Anisotropy	0.236	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.31,93.8	EDS
L-test for twinning ²	$ < L >=0.47, < L^2>=0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	14499	wwPDB-VP
Average B, all atoms $(Å^2)$	100.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.28% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: TB2, FRU, GLC, CM5, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bo	nd lengths	B	ond angles
IVIOI	Unain	RMSZ # Z > 5		RMSZ	# Z > 5
1	А	0.31	1/3672~(0.0%)	0.56	6/4995~(0.1%)
1	В	0.35	0/3679	0.69	14/5003~(0.3%)
1	С	0.31	0/3484	0.67	5/4749~(0.1%)
1	D	0.32	0/3527	0.70	9/4814~(0.2%)
All	All	0.32	1/14362~(0.0%)	0.66	34/19561~(0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	А	161	LYS	CA-CB	-5.92	1.41	1.53

The worst 5 of 34 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	В	37	LEU	N-CA-C	-5.98	94.85	111.00
1	D	328	ILE	CB-CA-C	-5.98	99.64	111.60
1	D	378	ARG	N-CA-C	5.97	127.12	111.00
1	В	335	HIS	N-CA-C	5.95	127.07	111.00
1	С	255	THR	N-CA-CB	-5.93	99.03	110.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	3583	0	3434	226	0
1	В	3589	0	3460	209	0
1	С	3406	0	3187	300	0
1	D	3445	0	3216	382	0
2	Е	23	0	21	5	0
2	F	23	0	21	9	0
2	G	23	0	21	4	0
2	Н	23	0	21	11	0
3	А	43	0	30	17	0
3	В	43	0	30	14	0
3	С	43	0	30	12	0
3	D	43	0	30	25	0
4	А	13	0	15	8	0
4	В	13	0	15	5	0
4	С	13	0	15	6	0
4	D	13	0	15	14	0
5	А	58	0	73	5	0
5	В	68	0	84	5	0
5	D	34	0	42	3	0
All	All	14499	0	13760	11114	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 39.

The worst 5 of 1114 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:30:LEU:HD12	1:D:31:PRO:CD	1.18	1.61
1:A:298:ALA:HB1	4:A:501:TB2:C8	1.39	1.52
1:D:30:LEU:CD1	1:D:31:PRO:HD2	1.39	1.51
1:D:298:ALA:HB2	4:D:501:TB2:C7	1.52	1.37
1:C:233:GLN:NE2	1:C:236:ARG:HD3	1.43	1.34

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	463/476~(97%)	440 (95%)	18 (4%)	5(1%)	14 50
1	В	463/476~(97%)	426~(92%)	36~(8%)	1 (0%)	47 82
1	С	452/476~(95%)	423~(94%)	27~(6%)	2~(0%)	34 72
1	D	462/476~(97%)	415 (90%)	43~(9%)	4 (1%)	17 55
All	All	1840/1904~(97%)	1704 (93%)	124 (7%)	12(1%)	22 60

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

5 of 12 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	А	39	VAL
1	А	45	GLN
1	D	430	SER
1	А	136	GLY
1	В	36	PRO

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric Outliers		Percentiles		
1	А	375/421~(89%)	338~(90%)	37 (10%)	8 30		
1	В	377/421~(90%)	336~(89%)	41 (11%)	6 25		
1	С	343/421~(82%)	299~(87%)	44 (13%)	4 19		
1	D	345/421~(82%)	301 (87%)	44 (13%)	4 19		
All	All	1440/1684~(86%)	1274 (88%)	166 (12%)	5 24		

5 of 166 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	С	437	LEU
1	D	253	ARG
1	С	483	SER



Continued from previous page...

Mol	Chain	Res	Type
1	D	107	ILE
1	D	317	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 15 such sidechains are listed below:

Mol	Chain	Res	Type
1	С	287	ASN
1	D	285	HIS
1	D	42	ASN
1	D	479	ASN
1	D	237	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

8 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Turne	Chain	Dec	Tink	Bo	ond leng	Bond angles			
MOI	туре	Unam	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GLC	Е	1	2	11,11,12	1.24	1 (9%)	$15,\!15,\!17$	2.55	8 (53%)
2	FRU	Е	2	2	11,12,12	1.37	3 (27%)	10,18,18	1.78	4 (40%)
2	GLC	F	1	2	11,11,12	1.22	2 (18%)	15,15,17	2.57	7 (46%)
2	FRU	F	2	2	11,12,12	1.28	2 (18%)	10,18,18	1.70	3 (30%)
2	GLC	G	1	2	11,11,12	1.27	2 (18%)	$15,\!15,\!17$	2.80	7 (46%)



Mal	Type Chain Reg		Tiple	Bo	ond leng	$_{\rm sths}$	Bond angles			
IVIOI	туре	Unain	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	FRU	G	2	2	11,12,12	1.29	1 (9%)	10,18,18	1.73	3 (30%)
2	GLC	Н	1	2	11,11,12	1.23	1 (9%)	15,15,17	2.46	7 (46%)
2	FRU	Н	2	2	11,12,12	1.28	1 (9%)	10,18,18	1.73	3 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	Е	1	2	-	0/2/19/22	0/1/1/1
2	FRU	Е	2	2	-	1/5/24/24	0/1/1/1
2	GLC	F	1	2	-	0/2/19/22	0/1/1/1
2	FRU	F	2	2	-	1/5/24/24	0/1/1/1
2	GLC	G	1	2	-	2/2/19/22	0/1/1/1
2	FRU	G	2	2	-	4/5/24/24	0/1/1/1
2	GLC	Н	1	2	-	0/2/19/22	0/1/1/1
2	FRU	Н	2	2	-	2/5/24/24	0/1/1/1

The worst 5 of 13 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	$\mathrm{Ideal}(\mathrm{\AA})$
2	G	2	FRU	O3-C3	-2.31	1.38	1.42
2	Е	2	FRU	O3-C3	-2.28	1.38	1.42
2	Н	2	FRU	O3-C3	-2.20	1.38	1.42
2	Е	1	GLC	O2-C2	-2.16	1.38	1.43
2	F	2	FRU	O3-C3	-2.13	1.38	1.42

The worst 5 of 42 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	G	1	GLC	O5-C5-C6	6.13	116.81	107.20
2	F	1	GLC	O5-C5-C6	5.79	116.28	107.20
2	Н	1	GLC	O5-C5-C6	5.57	115.94	107.20
2	Е	1	GLC	O5-C5-C6	5.30	115.52	107.20
2	G	1	GLC	O3-C3-C2	4.03	117.72	109.99

There are no chirality outliers.

5 of 10 torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
2	G	2	FRU	O1-C1-C2-O2
2	G	2	FRU	C4-C5-C6-O6
2	G	2	FRU	O5-C5-C6-O6
2	G	1	GLC	O5-C5-C6-O6
2	G	1	GLC	C4-C5-C6-O6

There are no ring outliers.

8 monomers are involved in 29 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	G	2	FRU	2	0
2	Н	1	GLC	11	0
2	Е	1	GLC	5	0
2	G	1	GLC	3	0
2	F	2	FRU	9	0
2	Н	2	FRU	8	0
2	Е	2	FRU	5	0
2	F	1	GLC	8	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.











5.6 Ligand geometry (i)

13 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Bos	Link	Bo	ond leng	ths	B	ond ang	les
WIOI	Type	Ullalli	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	TB2	А	501	1	13,13,13	0.60	0	18,18,18	0.95	1(5%)
4	TB2	D	501	1	13,13,13	0.69	0	18,18,18	2.00	1 (5%)
3	HEM	D	500	1	41,50,50	1.97	6 (14%)	45,82,82	1.59	6 (13%)
4	TB2	С	501	1	13,13,13	0.60	0	18,18,18	0.97	1 (5%)
5	CM5	В	607	-	36,36,36	0.41	0	49,49,49	0.79	1 (2%)
5	CM5	D	610	-	36,36,36	0.39	0	49,49,49	0.75	1 (2%)
5	CM5	А	605	-	25,25,36	0.44	0	32,32,49	0.81	1 (3%)
3	HEM	В	500	1	41,50,50	1.94	6 (14%)	45,82,82	1.75	7 (15%)
3	HEM	С	500	1	41,50,50	1.97	6 (14%)	45,82,82	1.70	6 (13%)
5	CM5	А	608	-	36,36,36	0.44	0	49,49,49	1.19	4 (8%)
3	HEM	А	500	1	41,50,50	1.98	6 (14%)	45,82,82	1.82	8 (17%)
4	TB2	В	501	1	13,13,13	0.60	0	18,18,18	0.88	1 (5%)
5	CM5	В	606	-	36,36,36	0.41	0	49,49,49	0.98	3 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	TB2	А	501	1	-	0/9/9/9	0/1/1/1
4	TB2	D	501	1	-	0/9/9/9	0/1/1/1
3	HEM	D	500	1	-	1/12/54/54	-
4	TB2	С	501	1	-	6/9/9/9	0/1/1/1
5	CM5	В	607	-	-	9/17/65/65	0/3/3/3
5	CM5	D	610	-	-	8/17/65/65	0/3/3/3
5	CM5	А	605	-	-	7/13/41/65	0/2/2/3



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	HEM	В	500	1	-	6/12/54/54	-
3	HEM	С	500	1	-	2/12/54/54	-
5	CM5	А	608	-	-	10/17/65/65	0/3/3/3
3	HEM	А	500	1	-	7/12/54/54	-
4	TB2	В	501	1	-	6/9/9/9	0/1/1/1
5	CM5	В	606	-	-	9/17/65/65	0/3/3/3

The worst 5 of 24 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	$\mathrm{Ideal}(\mathrm{\AA})$
3	С	500	HEM	C3D-C2D	8.13	1.54	1.36
3	D	500	HEM	C3D-C2D	8.02	1.53	1.36
3	В	500	HEM	C3D-C2D	7.92	1.53	1.36
3	А	500	HEM	C3D-C2D	7.89	1.53	1.36
3	А	500	HEM	C3C-C2C	-4.90	1.33	1.40

The worst 5 of 41 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms		$Observed(^{o})$	$Ideal(^{o})$
4	D	501	TB2	O1-C1-C2	-7.78	104.91	126.64
3	А	500	HEM	C4D-ND-C1D	6.35	111.63	105.07
3	С	500	HEM	C4D-ND-C1D	6.33	111.61	105.07
3	В	500	HEM	C4D-ND-C1D	6.17	111.45	105.07
3	D	500	HEM	C4D-ND-C1D	6.03	111.30	105.07

There are no chirality outliers.

5 of 71 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	А	500	HEM	C1A-C2A-CAA-CBA
3	А	500	HEM	C3A-C2A-CAA-CBA
5	А	605	CM5	O14-C13-O12-C1
5	А	608	CM5	C18-C13-O12-C1
5	А	608	CM5	O14-C13-O12-C1

There are no ring outliers.

12 monomers are involved in 114 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	А	501	TB2	8	0



Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	501	TB2	14	0
3	D	500	HEM	25	0
4	С	501	TB2	6	0
5	В	607	CM5	3	0
5	D	610	CM5	3	0
5	А	605	CM5	5	0
3	В	500	HEM	14	0
3	С	500	HEM	12	0
3	А	500	HEM	17	0
4	B	501	TB2	5	0
5	В	606	CM5	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



























Torsions



Rings











5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	465/476~(97%)	-0.21	4 (0%) 84 63	49, 79, 130, 196	2~(0%)
1	В	465/476~(97%)	-0.13	8 (1%) 70 41	54, 88, 160, 184	2 (0%)
1	С	456/476~(95%)	0.07	26 (5%) 23 8	56, 97, 181, 211	1 (0%)
1	D	464/476~(97%)	0.06	15 (3%) 47 20	60, 127, 166, 206	0
All	All	1850/1904 (97%)	-0.05	53 (2%) 51 23	49, 95, 166, 211	5 (0%)

The worst 5 of 53 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	50	GLY	10.7
1	А	38	PRO	6.6
1	С	39	VAL	6.3
1	А	39	VAL	6.0
1	С	43	LEU	5.2

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathrm{\AA}^2)$	Q<0.9
2	FRU	G	2	12/12	0.80	0.23	91,108,111,120	0
2	FRU	Е	2	12/12	0.86	0.18	66,87,93,93	0



Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(Å^2)$	Q<0.9
2	GLC	Е	1	11/12	0.86	0.15	54,76,86,98	0
2	GLC	Н	1	11/12	0.88	0.25	140,144,153,155	0
2	GLC	G	1	11/12	0.90	0.17	82,92,101,102	0
2	GLC	F	1	11/12	0.92	0.14	78,90,98,100	0
2	FRU	Н	2	12/12	0.92	0.11	141,148,151,154	0
2	FRU	F	2	12/12	0.93	0.15	85,89,94,95	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.













6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathrm{\AA}^2)$	Q<0.9
5	CM5	А	608	34/34	0.76	0.43	$61,\!125,\!148,\!155$	0
5	CM5	В	607	34/34	0.77	0.32	59,122,134,136	0
5	CM5	D	610	34/34	0.79	0.34	89,144,167,172	0
4	TB2	D	501	13/13	0.86	0.51	98,106,114,116	0
5	CM5	В	606	34/34	0.88	0.24	$68,\!114,\!133,\!137$	0
5	CM5	А	605	24/34	0.89	0.30	65,96,109,115	0
3	HEM	D	500	43/43	0.93	0.29	94,103,118,127	0
4	TB2	В	501	13/13	0.95	0.30	53,65,74,76	0
4	TB2	С	501	13/13	0.96	0.33	71,79,85,88	0
3	HEM	С	500	43/43	0.97	0.27	66, 83, 95, 108	0
4	TB2	А	501	13/13	0.97	0.29	$50,\!61,\!66,\!68$	0
3	HEM	В	500	43/43	0.98	0.28	46,60,72,85	0
3	HEM	А	500	43/43	0.98	0.27	40,54,64,71	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

























6.5 Other polymers (i)

There are no such residues in this entry.

