



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 17, 2021 – 02:05 AM EDT

PDB ID : 1R22
Title : Crystal structure of the cyanobacterial metallothionein repressor SmtB (C14S/C61S/C121S mutant) in the Zn₂alpha₅-form
Authors : Eicken, C.; Pennella, M.A.; Chen, X.; Koshlap, K.M.; VanZile, M.L.; Sacchetti, J.C.; Giedroc, D.P.
Deposited on : 2003-09-25
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	122	 44% 23% 8% • 24%
1	B	122	 51% 23% 5% • 20%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 1567 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transcriptional repressor smtB.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
1	A	93	734	465	135	134	0	0	0
1	B	97	771	484	147	140	0	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	14	SER	CYS	engineered mutation	UNP P30340
A	61	SER	CYS	engineered mutation	UNP P30340
A	121	SER	CYS	engineered mutation	UNP P30340
B	14	SER	CYS	engineered mutation	UNP P30340
B	61	SER	CYS	engineered mutation	UNP P30340
B	121	SER	CYS	engineered mutation	UNP P30340

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Zn		
2	A	1	1	1	0	0
2	B	1	1	1	0	0

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	23	23	23	0	0
3	B	37	37	37	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: Transcriptional repressor smtB



- Molecule 1: Transcriptional repressor smtB



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	64.45Å 124.68Å 25.91Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.30	Depositor
% Data completeness (in resolution range)	94.4 (50.00-2.30)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.229 , 0.290	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	1567	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:
ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.30	7/745 (0.9%)	1.14	7/1011 (0.7%)
1	B	1.37	2/783 (0.3%)	1.28	9/1059 (0.8%)
All	All	1.33	9/1528 (0.6%)	1.21	16/2070 (0.8%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	97	HIS	CE1-NE2	13.19	1.62	1.32
1	B	61	SER	CA-CB	7.93	1.64	1.52
1	B	100	TYR	CD2-CE2	6.06	1.48	1.39
1	A	34	GLN	CB-CG	5.39	1.67	1.52
1	A	121	SER	CA-CB	5.33	1.60	1.52
1	A	31	GLU	CD-OE1	5.32	1.31	1.25
1	A	31	GLU	CD-OE2	5.28	1.31	1.25
1	A	38	GLU	CD-OE2	5.19	1.31	1.25
1	A	34	GLN	CG-CD	5.01	1.62	1.51

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	92	ARG	NE-CZ-NH2	-7.41	116.59	120.30
1	B	92	ARG	NE-CZ-NH1	7.28	123.94	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	97	HIS	ND1-CG-CD2	7.20	118.88	108.80
1	A	64	ASP	CB-CG-OD2	7.14	124.73	118.30
1	B	87	ARG	NE-CZ-NH1	7.06	123.83	120.30
1	A	104	ASP	CB-CG-OD1	7.03	124.63	118.30
1	B	84	ARG	NE-CZ-NH2	-6.84	116.88	120.30
1	A	97	HIS	CG-ND1-CE1	-6.51	97.23	105.70
1	A	45	ASP	CB-CG-OD2	5.63	123.37	118.30
1	B	80	LEU	CA-CB-CG	5.42	127.77	115.30
1	A	97	HIS	CB-CG-ND1	-5.39	109.73	123.20
1	B	84	ARG	NE-CZ-NH1	5.30	122.95	120.30
1	B	88	LEU	CA-CB-CG	-5.18	103.38	115.30
1	A	50	ARG	NE-CZ-NH2	-5.17	117.71	120.30
1	B	80	LEU	CB-CG-CD1	5.15	119.76	111.00
1	B	116	ASP	CB-CG-OD2	5.14	122.93	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	120	GLU	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	734	0	733	29	0
1	B	771	0	773	26	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	23	0	0	2	0
3	B	37	0	0	3	0
All	All	1567	0	1506	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:72:SER:O	1:B:73:GLU:CB	2.19	0.90
1:A:117:HIS:CD2	1:B:107:ILE:HD11	2.08	0.87
1:B:72:SER:O	1:B:73:GLU:HB3	1.86	0.74
1:A:42:VAL:HG21	1:A:86:LEU:HD13	1.72	0.70
1:B:76:VAL:HG12	1:B:80:LEU:HD22	1.74	0.70
1:A:25:LEU:HD22	1:B:57:ARG:HH12	1.58	0.68
1:B:72:SER:O	1:B:73:GLU:HB2	1.99	0.63
1:A:97:HIS:CD2	1:A:97:HIS:N	2.67	0.63
1:A:39:PHE:O	1:A:42:VAL:HG12	1.98	0.62
1:B:57:ARG:O	1:B:58:SER:HB3	2.01	0.61
1:B:43:LEU:HD21	1:B:88:LEU:HD22	1.85	0.58
1:A:51:LEU:HD13	1:A:80:LEU:CD1	2.35	0.56
1:B:45:ASP:OD2	1:B:48:ARG:HG3	2.04	0.56
1:A:58:SER:OG	1:A:59:GLU:O	2.24	0.56
1:B:101:GLN:HG3	3:B:1007:HOH:O	2.06	0.56
1:A:42:VAL:HG23	1:A:48:ARG:NH1	2.21	0.55
1:B:57:ARG:O	1:B:58:SER:CB	2.55	0.55
1:B:59:GLU:OE1	1:B:92:ARG:HD3	2.07	0.54
1:B:39:PHE:HE2	1:B:110:LEU:HD11	1.71	0.54
1:A:87:ARG:NH2	1:B:120:GLU:OE2	2.41	0.53
1:A:84:ARG:HG3	1:A:85:ASN:N	2.24	0.53
1:A:117:HIS:CE1	1:B:103:GLN:HG2	2.43	0.53
1:A:117:HIS:HD2	1:B:107:ILE:HD11	1.70	0.51
1:A:48:ARG:HD2	3:A:1005:HOH:O	2.12	0.49
1:A:62:VAL:O	1:A:76:VAL:HG21	2.13	0.49
1:A:119:GLN:HE21	1:A:119:GLN:N	2.12	0.48
1:A:121:SER:HA	3:A:1016:HOH:O	2.14	0.47
1:A:54:LEU:O	1:A:57:ARG:HG2	2.13	0.47
1:A:48:ARG:NH2	1:A:82:SER:OG	2.47	0.47
1:B:87:ARG:HD3	3:B:1002:HOH:O	2.15	0.47
1:B:62:VAL:HG12	1:B:73:GLU:HG3	1.97	0.47
1:A:83:LEU:HA	1:A:86:LEU:HD12	1.97	0.46
1:B:39:PHE:CE2	1:B:110:LEU:HD11	2.51	0.45
1:A:45:ASP:HA	1:A:46:PRO:HD3	1.87	0.45
1:A:86:LEU:O	1:A:87:ARG:HB2	2.18	0.44
1:A:50:ARG:O	1:A:53:SER:OG	2.31	0.43
1:A:76:VAL:O	1:A:80:LEU:HD22	2.19	0.43
1:A:84:ARG:CG	1:A:85:ASN:N	2.82	0.43
1:B:26:GLN:O	1:B:27:ALA:HB3	2.19	0.42
1:A:91:TYR:HA	1:A:99:TYR:O	2.19	0.42
1:B:69:ILE:HG22	1:B:71:VAL:HG13	2.01	0.42
1:A:72:SER:O	1:A:76:VAL:HG22	2.20	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:60:LEU:HB3	1:B:64:ASP:HB2	2.01	0.41
1:B:57:ARG:NH1	3:B:1057:HOH:O	2.54	0.41
1:A:107:ILE:N	1:A:107:ILE:HD13	2.35	0.41
1:B:62:VAL:HG13	1:B:76:VAL:HB	2.03	0.41
1:B:34:GLN:O	1:B:38:GLU:HG3	2.21	0.40
1:A:117:HIS:HE1	1:B:103:GLN:HG2	1.86	0.40
1:A:119:GLN:HE21	1:A:119:GLN:CA	2.33	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	89/122 (73%)	83 (93%)	5 (6%)	1 (1%)	14	15
1	B	95/122 (78%)	88 (93%)	4 (4%)	3 (3%)	4	2
All	All	184/244 (75%)	171 (93%)	9 (5%)	4 (2%)	6	5

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	73	GLU
1	A	87	ARG
1	B	58	SER
1	B	121	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	77/102 (76%)	66 (86%)	11 (14%)	3	3
1	B	80/102 (78%)	77 (96%)	3 (4%)	33	47
All	All	157/204 (77%)	143 (91%)	14 (9%)	9	11

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	25	LEU
1	A	31	GLU
1	A	58	SER
1	A	73	GLU
1	A	74	SER
1	A	76	VAL
1	A	80	LEU
1	A	92	ARG
1	A	97	HIS
1	A	98	VAL
1	A	119	GLN
1	B	31	GLU
1	B	67	GLN
1	B	80	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	79	GLN
1	A	97	HIS
1	A	112	GLN
1	A	119	GLN
1	B	34	GLN
1	B	85	ASN
1	B	103	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.