



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 1, 2024 – 01:29 pm BST

PDB ID : 8R2J  
Title : X-ray crystallographic structure of SwaQ2 in complex with NADP<sup>+</sup> and doxorubicin  
Authors : Schnell, R.; Schneider, G.  
Deposited on : 2023-11-06  
Resolution : 2.40 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.37.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	292	
1	B	292	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	DM2	B	301	-	-	X	-

## 2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 4765 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NmrA family transcriptional regulator.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	274	Total 2117	C 1328	N 388	O 397	S 4	0	4	0
1	B	275	Total 2115	C 1326	N 389	O 396	S 4	0	3	0

- Molecule 2 is DOXORUBICIN (three-letter code: DM2) (formula:  $C_{27}H_{29}NO_{11}$ ) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	Total 39	C 27	N 1	O 11	0	0
2	A	1	Total 39	C 27	N 1	O 11	0	0
2	B	1	Total 39	C 27	N 1	O 11	0	0
2	B	1	Total 39	C 27	N 1	O 11	0	0




### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: NmrA family transcriptional regulator



- Molecule 1: NmrA family transcriptional regulator



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 65	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	87.31Å 87.31Å 174.38Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	29.08 – 2.40 29.06 – 2.40	Depositor EDS
% Data completeness (in resolution range)	93.6 (29.08-2.40) 93.6 (29.06-2.40)	Depositor EDS
$R_{merge}$	0.17	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.13 (at 2.39Å)	Xtrriage
Refinement program	REFMAC 5.8.0257	Depositor
R, $R_{free}$	0.161 , 0.203 0.174 , 0.214	Depositor DCC
$R_{free}$ test set	1444 reflections (5.25%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	27.1	Xtrriage
Anisotropy	0.832	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 41.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.35$	Xtrriage
Estimated twinning fraction	0.061 for h,-h-k,-l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4765	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	44.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.50% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAP, DM2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.75	0/2173	0.90	0/2954
1	B	0.76	1/2168 (0.0%)	0.90	0/2947
All	All	0.76	1/4341 (0.0%)	0.90	0/5901

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	22	GLU	CD-OE2	5.14	1.31	1.25

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2117	0	2090	13	1
1	B	2115	0	2087	13	2
2	A	78	0	56	13	0
2	B	78	0	57	21	0
3	A	48	0	25	1	0
3	B	48	0	25	0	0
4	A	138	0	0	2	0
4	B	143	0	0	0	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	4765	0	4340	58	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:301:DM2:O14	2:B:301:DM2:H5'	1.23	1.31
2:B:301:DM2:H2'1	2:B:302:DM2:C6'	1.75	1.16
2:B:301:DM2:O14	2:B:301:DM2:C5'	1.98	1.12
2:B:301:DM2:H2'1	2:B:302:DM2:H6'1	1.16	1.12
2:B:301:DM2:H5'	2:B:301:DM2:C14	1.96	0.96
2:B:301:DM2:C2'	2:B:302:DM2:H6'1	1.96	0.95
1:B:19:GLN:O	1:B:23[B]:ARG:HG3	1.73	0.88
2:A:301:DM2:N3'	2:A:302:DM2:H142	1.89	0.88
2:A:301:DM2:H111	2:A:301:DM2:H3'	1.54	0.87
2:B:301:DM2:C2'	2:B:302:DM2:C6'	2.57	0.81
1:B:10:GLY:O	1:B:14[B]:ARG:HG3	1.82	0.79
2:B:301:DM2:O14	2:B:301:DM2:C6'	2.31	0.78
2:A:301:DM2:H3'	2:A:301:DM2:C11	2.16	0.76
2:B:301:DM2:H2'1	2:B:302:DM2:H6'2	1.73	0.71
2:A:301:DM2:H112	2:A:301:DM2:O5'	1.91	0.70
1:B:208:ARG:NH1	1:B:210:GLU:OE1	2.26	0.69
1:A:148:ARG:HD2	1:A:220:ILE:HD13	1.77	0.65
2:B:301:DM2:C4'	2:B:302:DM2:H142	2.29	0.63
2:B:301:DM2:H3'	2:B:302:DM2:H142	1.80	0.62
2:B:301:DM2:H4'	2:B:302:DM2:H142	1.80	0.62
2:A:301:DM2:H6'3	2:A:301:DM2:H2'2	1.81	0.60
2:B:301:DM2:H6'3	2:B:301:DM2:N3'	2.17	0.60
2:A:301:DM2:H6'3	2:A:301:DM2:C2'	2.30	0.59
2:B:301:DM2:H3'	2:B:302:DM2:C14	2.34	0.57
1:A:10:GLY:O	1:A:14[A]:ARG:HG3	2.05	0.57
1:B:127:GLY:HA2	2:B:301:DM2:H213	1.88	0.56
2:B:301:DM2:C3'	2:B:302:DM2:H142	2.35	0.55
2:A:301:DM2:H111	2:A:301:DM2:C3'	2.32	0.55
2:B:301:DM2:H111	2:B:301:DM2:O5'	2.07	0.54
2:A:301:DM2:C11	2:A:301:DM2:C3'	2.85	0.54
1:B:18:ARG:NH2	1:B:162[A]:GLN:OE1	2.39	0.54
1:B:162[A]:GLN:NE2	1:B:165:ALA:HB3	2.23	0.53
2:B:301:DM2:H5'	2:B:301:DM2:C13	2.39	0.53

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:227:PHE:HE2	2:A:301:DM2:O13	1.93	0.52
2:A:301:DM2:H2'2	2:A:302:DM2:H6'1	1.93	0.50
1:B:139:PRO:HB2	1:B:140:PRO:HD3	1.94	0.48
2:B:301:DM2:C2'	2:B:302:DM2:H6'2	2.38	0.48
1:A:262:LEU:HD23	1:A:262:LEU:HA	1.71	0.47
2:A:301:DM2:C11	2:A:301:DM2:C2'	2.93	0.47
2:A:301:DM2:C2'	2:A:302:DM2:H6'1	2.45	0.47
1:A:90:ILE:O	1:A:121:TRP:HA	2.16	0.46
1:B:154:ALA:O	1:B:191:THR:HA	2.16	0.46
1:A:10:GLY:HA2	3:A:303:NAP:O1A	2.17	0.44
1:A:19:GLN:OE1	1:A:162[A]:GLN:NE2	2.49	0.44
2:B:301:DM2:C6'	2:B:301:DM2:N3'	2.81	0.44
1:B:90:ILE:O	1:B:121:TRP:HA	2.18	0.44
1:B:23[A]:ARG:HH11	1:B:23[A]:ARG:HG3	1.82	0.44
1:A:288:ASP:OD1	1:A:291:ARG:NH1	2.51	0.43
1:B:7:GLY:O	1:B:13:GLY:HA3	2.18	0.43
1:A:18:ARG:NH2	4:A:406:HOH:O	2.51	0.43
1:A:139:PRO:HB2	1:A:140:PRO:HD3	2.01	0.43
1:B:149:GLU:OE2	1:B:150:PRO:HD2	2.19	0.42
1:B:281:GLN:NE2	1:B:285:ASP:OD1	2.50	0.41
2:A:301:DM2:H111	2:A:301:DM2:C2'	2.51	0.41
2:B:301:DM2:H5'	2:B:301:DM2:C12	2.48	0.41
1:A:165:ALA:O	1:A:169:VAL:HG23	2.21	0.41
1:A:188:GLU:HG3	4:A:451:HOH:O	2.21	0.40
1:A:157:PHE:CZ	1:A:189:LEU:HB2	2.56	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:23[A]:ARG:NH1	1:B:105:ASP:OD1[6_554]	2.00	0.20
1:A:218:ARG:NH1	1:B:223:ALA:O[6_454]	2.05	0.15

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	274/292 (94%)	268 (98%)	6 (2%)	0	100	100
1	B	274/292 (94%)	264 (96%)	10 (4%)	0	100	100
All	All	548/584 (94%)	532 (97%)	16 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	212/221 (96%)	207 (98%)	5 (2%)	49	68
1	B	211/221 (96%)	209 (99%)	2 (1%)	78	90
All	All	423/442 (96%)	416 (98%)	7 (2%)	60	78

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	SER
1	A	35	ARG
1	A	57	LYS
1	A	188	GLU
1	A	218	ARG
1	B	1	SER
1	B	57	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	DM2	A	301	-	41,43,43	0.95	3 (7%)	55,67,67	1.49	7 (12%)
2	DM2	B	301	-	41,43,43	0.90	0	55,67,67	1.05	3 (5%)
2	DM2	A	302	-	41,43,43	0.83	1 (2%)	55,67,67	0.87	3 (5%)
2	DM2	B	302	-	41,43,43	0.56	0	55,67,67	0.78	1 (1%)
3	NAP	B	303	-	45,52,52	0.98	3 (6%)	56,80,80	1.54	9 (16%)
3	NAP	A	303	-	45,52,52	0.95	1 (2%)	56,80,80	1.55	11 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	DM2	A	301	-	-	6/13/60/60	0/5/5/5
2	DM2	B	301	-	-	8/13/60/60	0/5/5/5
2	DM2	A	302	-	-	2/13/60/60	0/5/5/5
2	DM2	B	302	-	-	2/13/60/60	0/5/5/5
3	NAP	B	303	-	-	5/31/67/67	0/5/5/5
3	NAP	A	303	-	-	4/31/67/67	0/5/5/5

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	303	NAP	O4B-C1B	3.16	1.45	1.41
2	A	301	DM2	C7-C8	2.59	1.46	1.41
3	A	303	NAP	O4B-C1B	2.57	1.44	1.41
2	A	302	DM2	O6-C6	2.37	1.27	1.22
3	B	303	NAP	P2B-O2B	2.19	1.63	1.59
3	B	303	NAP	C5A-N7A	-2.17	1.31	1.39
2	A	301	DM2	C7-C6	2.11	1.52	1.47
2	A	301	DM2	C8-C9	2.04	1.43	1.40

All (34) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	301	DM2	C8-C9-C10	6.80	129.25	118.59
2	A	301	DM2	C16-C9-C10	-4.16	114.19	120.96
3	A	303	NAP	C5B-C4B-C3B	-4.13	99.69	115.18
3	B	303	NAP	O7N-C7N-N7N	-3.81	117.16	122.58
3	B	303	NAP	C5B-C4B-C3B	-3.66	101.46	115.18
3	B	303	NAP	N3A-C2A-N1A	-3.54	123.14	128.68
3	B	303	NAP	O2N-PN-O1N	3.47	129.37	112.24
3	A	303	NAP	O2N-PN-O1N	3.37	128.92	112.24
3	B	303	NAP	O4B-C4B-C5B	3.27	120.14	109.37
2	B	301	DM2	C8-C9-C10	3.25	123.69	118.59
3	A	303	NAP	O4B-C4B-C5B	3.05	119.41	109.37
3	A	303	NAP	N3A-C2A-N1A	-2.92	124.12	128.68
3	B	303	NAP	O7N-C7N-C3N	2.90	123.11	119.63
3	B	303	NAP	O4B-C1B-C2B	-2.85	101.64	106.59
3	A	303	NAP	O7N-C7N-N7N	-2.79	118.61	122.58
3	B	303	NAP	O3X-P2B-O2X	2.77	118.21	107.64
3	A	303	NAP	C3D-C2D-C1D	2.74	105.10	100.98
3	A	303	NAP	O4B-C1B-C2B	-2.60	102.08	106.59
2	A	302	DM2	O4-C4-C5	2.59	119.46	115.85
3	B	303	NAP	C3D-C2D-C1D	2.57	104.85	100.98
2	B	301	DM2	O17-C17-C18	-2.55	115.29	120.98
2	A	302	DM2	C14-C13-C12	2.50	120.58	117.73
2	A	301	DM2	C8-C7-C6	2.43	124.53	120.45
2	B	302	DM2	O12-C12-C11	2.38	114.79	109.41
2	A	301	DM2	C15-C16-C17	-2.35	114.86	119.23
3	A	303	NAP	O3X-P2B-O2X	2.35	116.62	107.64
2	A	301	DM2	C18-C7-C6	-2.35	116.84	120.01
3	A	303	NAP	C6N-N1N-C2N	-2.30	119.88	121.97
2	A	302	DM2	C21-O4-C4	2.18	120.82	117.53
2	A	301	DM2	C2'-C3'-C4'	2.15	113.28	110.04

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	301	DM2	O19-C19-C20	-2.11	117.82	120.91
3	A	303	NAP	PN-O3-PA	-2.08	125.69	132.83
3	A	303	NAP	C3N-C7N-N7N	2.01	120.17	117.75
2	B	301	DM2	C21-O4-C4	2.01	120.56	117.53

There are no chirality outliers.

All (27) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	301	DM2	C11-C10-O10-C1'
2	A	301	DM2	C2'-C1'-O10-C10
2	A	301	DM2	C12-C13-C14-O14
2	A	301	DM2	O13-C13-C14-O14
2	A	302	DM2	C12-C13-C14-O14
2	A	302	DM2	O13-C13-C14-O14
2	B	301	DM2	C9-C10-O10-C1'
2	B	301	DM2	C12-C13-C14-O14
2	B	301	DM2	O13-C13-C14-O14
2	B	302	DM2	C12-C13-C14-O14
2	B	302	DM2	O13-C13-C14-O14
3	A	303	NAP	C5B-O5B-PA-O1A
3	A	303	NAP	C5B-O5B-PA-O2A
3	B	303	NAP	C5B-O5B-PA-O1A
3	B	303	NAP	C3D-C4D-C5D-O5D
2	B	301	DM2	C15-C12-C13-C14
3	B	303	NAP	O4D-C4D-C5D-O5D
2	A	301	DM2	O12-C12-C13-O13
3	B	303	NAP	C5B-O5B-PA-O2A
2	B	301	DM2	C3-C4-O4-C21
3	A	303	NAP	C3D-C4D-C5D-O5D
2	B	301	DM2	C11-C12-C13-C14
2	A	301	DM2	O5'-C1'-O10-C10
2	B	301	DM2	C5-C4-O4-C21
3	A	303	NAP	C5B-O5B-PA-O3
3	B	303	NAP	C5B-O5B-PA-O3
2	B	301	DM2	C11-C10-O10-C1'



There are no ring outliers.

5 monomers are involved in 35 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	301	DM2	13	0
2	B	301	DM2	21	0
2	A	302	DM2	3	0
2	B	302	DM2	11	0
3	A	303	NAP	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	274/292 (93%)	-0.17	14 (5%) 28 26	21, 35, 88, 107	0
1	B	275/292 (94%)	-0.11	18 (6%) 18 17	20, 35, 80, 102	0
All	All	549/584 (94%)	-0.14	32 (5%) 23 22	20, 35, 87, 107	0

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	292	ALA	7.6
1	B	36	PRO	5.5
1	A	238	PHE	4.1
1	B	239	GLU	3.9
1	B	292	ALA	3.8
1	B	238	PHE	3.6
1	A	223	ALA	3.5
1	B	264	PRO	3.3
1	B	261	SER	3.2
1	B	262	LEU	3.2
1	B	263	GLY	3.0
1	B	37	GLY	2.8
1	B	234	PHE	2.8
1	A	239	GLU	2.8
1	A	215	GLN	2.7
1	B	243	GLY	2.7
1	A	224	GLN	2.6
1	A	219	GLU	2.6
1	B	240	ASP	2.5
1	B	215	GLN	2.5
1	B	35	ARG	2.5
1	A	204	GLY	2.3
1	A	206	GLU	2.3
1	A	229	ALA	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	242	GLY	2.2
1	A	41	GLU	2.2
1	A	234	PHE	2.1
1	B	143	ALA	2.1
1	B	33	ARG	2.1
1	B	242	GLY	2.1
1	B	38	ALA	2.1
1	A	261	SER	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	DM2	B	301	39/39	0.86	0.23	50,62,82,89	12
2	DM2	A	301	39/39	0.88	0.24	49,60,88,90	12
2	DM2	A	302	39/39	0.92	0.14	51,61,66,76	0
2	DM2	B	302	39/39	0.94	0.15	45,58,66,74	0
3	NAP	A	303	48/48	0.98	0.10	20,28,33,36	0
3	NAP	B	303	48/48	0.98	0.11	22,32,35,37	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around DM2 A 302:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around DM2 B 302:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around NAP A 303:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around NAP B 303:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.