



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 6, 2024 – 03:34 pm GMT

PDB ID : 8R2T  
Title : Crystal structure of 4-hydroxybenzoate-1-hydroxylase from *Gelatoporia subvermispora* (GsMNX1)  
Authors : Zahn, M.; Kuatsjah, E.; Salvachua, D.  
Deposited on : 2023-11-07  
Resolution : 1.82 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 3.0  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.003 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.82 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	9242 (1.84-1.80)
Clashscore	180529	1080 (1.82-1.82)
Ramachandran outliers	177936	1073 (1.82-1.82)
Sidechain outliers	177891	1073 (1.82-1.82)
RSRZ outliers	164620	9241 (1.84-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	483	 2% 77% 8% • 14%
1	B	483	 19% 78% 5% • 15%

## 2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 6921 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called FAD-binding domain-containing protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	417	3253	2054	589	594	16	0	4	0
1	B	410	3181	2008	575	582	16	0	0	0

There are 28 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	470	ALA	-	expression tag	UNP M2QGN5
A	471	GLU	-	expression tag	UNP M2QGN5
A	472	ASN	-	expression tag	UNP M2QGN5
A	473	LEU	-	expression tag	UNP M2QGN5
A	474	TYR	-	expression tag	UNP M2QGN5
A	475	PHE	-	expression tag	UNP M2QGN5
A	476	GLN	-	expression tag	UNP M2QGN5
A	477	GLY	-	expression tag	UNP M2QGN5
A	478	HIS	-	expression tag	UNP M2QGN5
A	479	HIS	-	expression tag	UNP M2QGN5
A	480	HIS	-	expression tag	UNP M2QGN5
A	481	HIS	-	expression tag	UNP M2QGN5
A	482	HIS	-	expression tag	UNP M2QGN5
A	483	HIS	-	expression tag	UNP M2QGN5
B	470	ALA	-	expression tag	UNP M2QGN5
B	471	GLU	-	expression tag	UNP M2QGN5
B	472	ASN	-	expression tag	UNP M2QGN5
B	473	LEU	-	expression tag	UNP M2QGN5
B	474	TYR	-	expression tag	UNP M2QGN5
B	475	PHE	-	expression tag	UNP M2QGN5
B	476	GLN	-	expression tag	UNP M2QGN5
B	477	GLY	-	expression tag	UNP M2QGN5
B	478	HIS	-	expression tag	UNP M2QGN5
B	479	HIS	-	expression tag	UNP M2QGN5
B	480	HIS	-	expression tag	UNP M2QGN5

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	481	HIS	-	expression tag	UNP M2QGN5
B	482	HIS	-	expression tag	UNP M2QGN5
B	483	HIS	-	expression tag	UNP M2QGN5

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula:  $C_{27}H_{33}N_9O_{15}P_2$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	53	27	9	15	2	0	0
2	B	1	53	27	9	15	2	0	0

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Cl		
3	A	1	1	1	0	0
3	B	1	1	1	0	0


- Molecule 4 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
4	A	274	Total 274	O 274	0	0
4	B	105	Total 105	O 105	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: FAD-binding domain-containing protein



- Molecule 1: FAD-binding domain-containing protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	112.22Å 64.79Å 160.10Å 90.00° 93.90° 90.00°	Depositor
Resolution (Å)	159.73 – 1.82 159.73 – 1.82	Depositor EDS
% Data completeness (in resolution range)	56.3 (159.73-1.82) 56.3 (159.73-1.82)	Depositor EDS
$R_{merge}$	0.18	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.34 (at 1.82Å)	Xtrriage
Refinement program	REFMAC 5.8.0405	Depositor
R, $R_{free}$	0.179 , 0.225 0.187 , 0.232	Depositor DCC
$R_{free}$ test set	2912 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	25.6	Xtrriage
Anisotropy	0.016	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 34.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	6921	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	37.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.27% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: FAD, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.58	3/3336 (0.1%)	0.86	4/4522 (0.1%)
1	B	0.43	0/3249	0.76	0/4402
All	All	0.51	3/6585 (0.0%)	0.81	4/8924 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	9
1	B	0	2
All	All	0	11

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	242	GLU	CD-OE1	6.63	1.32	1.25
1	A	439	GLU	CD-OE1	5.94	1.32	1.25
1	A	450	GLU	CD-OE2	5.02	1.31	1.25

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	43	ARG	NE-CZ-NH1	5.73	123.17	120.30
1	A	125	ARG	NE-CZ-NH1	5.57	123.08	120.30
1	A	108	ARG	NE-CZ-NH1	-5.30	117.65	120.30
1	A	72	ARG	CB-CG-CD	5.05	124.72	111.60

There are no chirality outliers.



All (11) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	183	ARG	Sidechain
1	A	191	SER	Peptide
1	A	297	ARG	Sidechain
1	A	309[A]	ARG	Sidechain
1	A	309[B]	ARG	Sidechain
1	A	374[A]	ARG	Sidechain
1	A	374[B]	ARG	Sidechain
1	A	426	GLY	Peptide
1	A	84	ARG	Sidechain
1	B	210	ARG	Sidechain
1	B	374	ARG	Sidechain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3253	0	3241	18	0
1	B	3181	0	3151	19	0
2	A	53	0	31	1	0
2	B	53	0	31	1	0
3	A	1	0	0	1	0
3	B	1	0	0	1	0
4	A	274	0	0	2	0
4	B	105	0	0	2	0
All	All	6921	0	6454	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:382:LEU:HD11	1:A:426:GLY:HA2	1.44	0.99
1:A:382:LEU:HD11	1:A:426:GLY:CA	2.09	0.82
1:A:154:GLU:OE1	4:A:601:HOH:O	1.99	0.80

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:84:ARG:HH11	1:B:84:ARG:HB3	1.59	0.67
1:B:276:GLU:OE1	4:B:601:HOH:O	2.13	0.66
1:B:177:VAL:CG2	1:B:295:MET:HG2	2.28	0.64
1:B:322:PRO:HA	3:B:502:CL:CL	2.40	0.58
1:A:463:SER:O	1:A:464:TYR:C	2.45	0.55
1:B:186:LEU:HD11	1:B:300:LEU:HD22	1.88	0.55
1:A:214:GLU:OE2	4:A:602:HOH:O	2.17	0.55
1:B:150:SER:HB3	1:B:159:ILE:CG1	2.37	0.54
1:A:322:PRO:HA	3:A:502:CL:CL	2.45	0.54
1:B:360:GLU:OE2	1:B:460:ARG:NH2	2.42	0.52
1:B:150:SER:HB3	1:B:159:ILE:HG12	1.92	0.51
1:B:352:LYS:HB3	1:B:353:PRO:HD3	1.93	0.50
1:A:352:LYS:HB3	1:A:353:PRO:HD3	1.96	0.47
1:A:363:LEU:HB3	1:A:364:PRO:HD3	1.95	0.47
1:A:195:THR:HG21	1:A:321:LEU:HD21	1.96	0.47
1:B:327:GLY:HA3	2:B:501:FAD:H1'2	1.96	0.47
1:B:354:LEU:HD23	1:B:461:ILE:HD13	1.96	0.46
1:B:177:VAL:HG21	1:B:295:MET:HG2	1.98	0.46
1:A:267:LYS:NZ	1:A:271:ASP:OD1	2.48	0.44
1:A:327:GLY:HA3	2:A:501:FAD:H1'2	2.00	0.43
1:B:210:ARG:HD2	1:B:216:ARG:HG3	1.99	0.43
1:A:199:VAL:HA	1:A:248:LEU:O	2.19	0.43
1:B:320:MET:HB3	1:B:370:GLN:HG2	2.01	0.43
1:B:435:GLN:H	1:B:435:GLN:CD	2.22	0.43
1:B:296:ASP:HB3	4:B:604:HOH:O	2.18	0.42
1:B:313:LEU:HD23	1:B:313:LEU:C	2.39	0.42
1:B:199:VAL:HA	1:B:248:LEU:O	2.19	0.42
1:A:231:GLN:HA	1:A:381:HIS:HB3	2.02	0.42
1:A:313:LEU:C	1:A:313:LEU:HD23	2.40	0.42
1:A:436:LYS:HB3	1:A:436:LYS:HE3	1.90	0.41
1:A:320:MET:HB3	1:A:370[A]:GLN:HG2	2.03	0.41
1:B:231:GLN:HA	1:B:381:HIS:HB3	2.01	0.41
1:A:39:TYR:CZ	1:A:43:ARG:HD3	2.56	0.41
1:A:90:PRO:HG3	1:A:122:HIS:CE1	2.57	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	415/483 (86%)	405 (98%)	10 (2%)	0	100	100
1	B	402/483 (83%)	389 (97%)	13 (3%)	0	100	100
All	All	817/966 (85%)	794 (97%)	23 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	334/380 (88%)	329 (98%)	5 (2%)	60	47
1	B	323/380 (85%)	315 (98%)	8 (2%)	42	25
All	All	657/760 (86%)	644 (98%)	13 (2%)	50	35

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	72	ARG
1	A	276	GLU
1	A	401	VAL
1	A	460	ARG
1	A	463	SER
1	B	72	ARG
1	B	84	ARG
1	B	276	GLU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	287	LYS
1	B	295	MET
1	B	375	SER
1	B	460	ARG
1	B	461	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	434	HIS
1	B	435	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	FAD	B	501	-	53,58,58	0.66	0	68,89,89	0.81	3 (4%)
2	FAD	A	501	-	53,58,58	0.81	0	68,89,89	0.94	3 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	B	501	-	-	1/30/50/50	0/6/6/6
2	FAD	A	501	-	-	1/30/50/50	0/6/6/6

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	FAD	O4B-C1B-C2B	-2.87	102.72	106.93
2	A	501	FAD	O2A-PA-O1A	2.71	125.62	112.24
2	B	501	FAD	O2P-P-O1P	2.32	123.70	112.24
2	B	501	FAD	C4-N3-C2	-2.19	121.59	125.64
2	B	501	FAD	C5A-C6A-N6A	2.14	123.60	120.35
2	A	501	FAD	C5A-C6A-N6A	2.09	123.53	120.35

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	FAD	O4B-C4B-C5B-O5B
2	B	501	FAD	O4B-C4B-C5B-O5B


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	501	FAD	1	0
2	A	501	FAD	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and

any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	417/483 (86%)	-0.24	12 (2%) 54 53	13, 23, 45, 87	4 (0%)
1	B	410/483 (84%)	1.18	94 (22%) 2 2	24, 47, 77, 97	0
All	All	827/966 (85%)	0.46	106 (12%) 9 8	13, 33, 73, 97	4 (0%)

All (106) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	261	ALA	6.8
1	B	189	PRO	6.2
1	A	401	VAL	5.6
1	A	254	GLY	5.5
1	A	17	PRO	5.3
1	B	159	ILE	5.1
1	B	160	LEU	5.1
1	B	165	VAL	5.0
1	A	464	TYR	4.9
1	B	295	MET	4.8
1	A	402	VAL	4.7
1	B	44	VAL	4.6
1	B	426	GLY	4.5
1	A	263	GLY	4.5
1	B	144	LEU	4.4
1	B	307	ALA	4.4
1	B	18	LYS	4.0
1	B	161	ALA	3.9
1	B	348	ILE	3.8
1	B	456	VAL	3.8
1	B	57	ILE	3.7
1	B	151	VAL	3.7
1	B	464	TYR	3.5
1	B	304	VAL	3.4

*Continued on next page...*



*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	B	254	GLY	3.4
1	B	262	GLU	3.3
1	B	21	LEU	3.2
1	B	47	LYS	3.2
1	B	163	GLY	3.2
1	B	146	ALA	3.2
1	B	180	PHE	3.1
1	B	461	ILE	3.1
1	B	345	ILE	3.0
1	A	462	GLY	2.9
1	B	187	GLY	2.9
1	B	167	HIS	2.9
1	B	310	VAL	2.9
1	B	459	SER	2.9
1	B	186	LEU	2.9
1	B	300	LEU	2.9
1	B	149	VAL	2.9
1	A	426	GLY	2.9
1	B	170	LEU	2.9
1	B	290	LEU	2.9
1	B	50	VAL	2.9
1	B	31	GLY	2.9
1	B	188	LYS	2.9
1	B	457	GLY	2.8
1	B	85	LYS	2.8
1	B	297	ARG	2.8
1	A	463	SER	2.8
1	B	267	LYS	2.8
1	B	148	VAL	2.8
1	B	19	ALA	2.8
1	B	51	LEU	2.8
1	B	22	LYS	2.8
1	B	306	PRO	2.7
1	B	174	ALA	2.7
1	B	185	VAL	2.7
1	A	84	ARG	2.6
1	B	241	ALA	2.6
1	B	342	PHE	2.6
1	B	29	GLY	2.6
1	B	162	THR	2.6
1	B	23	LEU	2.6
1	B	24	HIS	2.5

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	B	54	ALA	2.5
1	B	356	TYR	2.5
1	B	312	LEU	2.5
1	B	158	ALA	2.5
1	B	340	VAL	2.5
1	B	56	THR	2.5
1	B	171	ILE	2.4
1	B	20	THR	2.4
1	B	26	LEU	2.4
1	B	210	ARG	2.3
1	B	346	SER	2.3
1	B	263	GLY	2.3
1	B	28	VAL	2.3
1	B	55	THR	2.3
1	B	355	LEU	2.3
1	B	157	SER	2.3
1	B	363	LEU	2.3
1	A	253	ASP	2.2
1	B	449	ALA	2.2
1	B	253	ASP	2.2
1	B	166	VAL	2.2
1	B	452	TRP	2.2
1	B	453	TRP	2.2
1	B	45	GLY	2.2
1	B	139	TYR	2.2
1	B	196	GLY	2.2
1	B	137	ALA	2.2
1	A	461	ILE	2.2
1	B	147	THR	2.1
1	B	351	LEU	2.1
1	B	448	ALA	2.1
1	B	27	ILE	2.1
1	B	142	LEU	2.1
1	B	293	ARG	2.1
1	B	135	LEU	2.1
1	B	289	THR	2.1
1	B	462	GLY	2.1
1	B	296	ASP	2.0
1	B	184	VAL	2.0
1	B	291	LYS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	FAD	B	501	53/53	0.94	0.09	35,45,57,58	0
3	CL	B	502	1/1	0.97	0.04	36,36,36,36	0
2	FAD	A	501	53/53	0.98	0.04	14,17,18,19	0
3	CL	A	502	1/1	1.00	0.04	18,18,18,18	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around FAD B 501:**


$2mF_o-DF_c$  (at 0.7 rnsd) in gray  
 $mF_o-DF_c$  (at 3 rnsd) in purple (negative)  
and green (positive)




**Electron density around CL B 502:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







## 6.5 Other polymers ⓘ

There are no such residues in this entry.