



# Full wwPDB X-ray Structure Validation Report ⓘ

May 14, 2020 – 02:37 am BST

PDB ID : 3R5J  
Title : Crystal structure of active caspase-2 bound with Ac-ADVAD-CHO  
Authors : Tang, Y.; Wells, J.; Arkin, M.  
Deposited on : 2011-03-18  
Resolution : 1.77 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.77 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	9185 (1.80-1.76)
Clashscore	141614	10184 (1.80-1.76)
Ramachandran outliers	138981	10051 (1.80-1.76)
Sidechain outliers	138945	10050 (1.80-1.76)
RSRZ outliers	127900	9032 (1.80-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	160	 3% 91% 5% 3%
1	C	160	 3% 89% 10% 2%
2	B	112	 14% 77% 9% 13%
2	D	112	 10% 79% 7% 13%
3	E	6	 17% 83% 17%
3	F	6	 83% 17%

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4368 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Caspase-2 subunit p18.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	154	1195	763	208	215	9	0	2	0
1	C	159	1228	781	211	227	9	0	2	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	174	MET	-	EXPRESSION TAG	UNP P42575
C	174	MET	-	EXPRESSION TAG	UNP P42575

- Molecule 2 is a protein called Caspase-2 subunit p12.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	97	771	486	138	132	15	3	3	0
2	D	97	777	491	140	132	14	1	3	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	453	LEU	-	EXPRESSION TAG	UNP P42575
B	454	GLU	-	EXPRESSION TAG	UNP P42575
B	455	HIS	-	EXPRESSION TAG	UNP P42575
B	456	HIS	-	EXPRESSION TAG	UNP P42575
B	457	HIS	-	EXPRESSION TAG	UNP P42575
B	458	HIS	-	EXPRESSION TAG	UNP P42575
B	459	HIS	-	EXPRESSION TAG	UNP P42575
B	460	HIS	-	EXPRESSION TAG	UNP P42575
D	453	LEU	-	EXPRESSION TAG	UNP P42575
D	454	GLU	-	EXPRESSION TAG	UNP P42575

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
D	455	HIS	-	EXPRESSION TAG	UNP P42575
D	456	HIS	-	EXPRESSION TAG	UNP P42575
D	457	HIS	-	EXPRESSION TAG	UNP P42575
D	458	HIS	-	EXPRESSION TAG	UNP P42575
D	459	HIS	-	EXPRESSION TAG	UNP P42575
D	460	HIS	-	EXPRESSION TAG	UNP P42575

- Molecule 3 is a protein called Peptide Inhibitor (ACE)ADVAD-CHO.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	F	6	Total	C	N	O	0	0	0
			36	21	5	10			
3	E	6	Total	C	N	O	0	0	0
			36	21	5	10			


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	97	Total	O	0	0
			97	97		
4	B	55	Total	O	0	0
			55	55		
4	C	107	Total	O	0	0
			107	107		
4	D	63	Total	O	0	0
			63	63		
4	F	2	Total	O	0	0
			2	2		
4	E	1	Total	O	0	0
			1	1		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Caspase-2 subunit p18




- Molecule 1: Caspase-2 subunit p18




- Molecule 2: Caspase-2 subunit p12




- Molecule 2: Caspase-2 subunit p12



- Molecule 3: Peptide Inhibitor (ACE)ADVAD-CHO



- Molecule 3: Peptide Inhibitor (ACE)ADVAD-CHO



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	63.30Å 97.01Å 98.15Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 1.77 46.64 – 1.77	Depositor EDS
% Data completeness (in resolution range)	99.3 (50.00-1.77) 99.3 (46.64-1.77)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	6.10 (at 1.77Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.184 , 0.218 0.183 , 0.215	Depositor DCC
$R_{free}$ test set	2982 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	23.8	Xtriage
Anisotropy	0.056	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 57.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.011 for -h,l,k	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4368	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.05% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ASA, ACE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.57	0/1225	0.63	0/1658
1	C	0.58	0/1258	0.67	0/1703
2	B	0.58	0/801	0.73	1/1079 (0.1%)
2	D	0.66	0/804	0.70	0/1080
3	E	0.43	0/25	0.66	0/34
3	F	0.43	0/25	0.69	0/34
All	All	0.59	0/4138	0.68	1/5588 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
2	B	417	ARG	NE-CZ-NH2	-5.03	117.79	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1195	0	1159	9	0
1	C	1228	0	1183	19	0
2	B	771	0	756	9	0
2	D	777	0	779	8	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	E	36	0	30	2	0
3	F	36	0	30	3	0
4	A	97	0	0	1	0
4	B	55	0	0	3	0
4	C	107	0	0	0	0
4	D	63	0	0	2	0
4	E	1	0	0	1	0
4	F	2	0	0	0	0
All	All	4368	0	3937	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:320:CYS:SG	3:F:406:ASA:C	2.27	1.23
2:D:366[A]:CYS:SG	4:D:562:HOH:O	2.14	1.05
1:A:194:LEU:HD22	1:A:199:ARG:HD2	1.48	0.95
1:C:320:CYS:SG	3:F:406:ASA:HXT	2.07	0.94
1:A:320:CYS:SG	3:E:406:ASA:C	2.56	0.93
2:B:366[A]:CYS:SG	4:B:554:HOH:O	2.28	0.91
1:A:190:LEU:O	4:A:590:HOH:O	2.08	0.70
1:C:190:LEU:HD23	2:D:442:HIS:HD2	1.59	0.67
2:B:424:THR:HG23	2:B:426:PHE:H	1.65	0.61
1:A:194:LEU:CD2	1:A:199:ARG:HD2	2.26	0.60
1:C:320:CYS:SG	3:F:406:ASA:CA	2.96	0.53
1:A:331[B]:GLN:NE2	1:C:310:ASN:ND2	2.57	0.52
1:C:194:LEU:HD22	2:D:445:LEU:HA	1.91	0.52
1:C:205:LEU:HD23	1:C:243:LEU:HB2	1.91	0.51
1:C:187:HIS:O	1:C:190:LEU:HD22	2.10	0.51
1:A:331[B]:GLN:NE2	1:C:310:ASN:HD21	2.09	0.50
1:C:194:LEU:HD23	1:C:199:ARG:HD2	1.94	0.50
2:B:441[B]:ARG:NH1	4:B:540:HOH:O	2.45	0.48
2:D:378:ARG:HA	2:D:384:SER:HA	1.94	0.48
1:C:205:LEU:CD2	1:C:243:LEU:HB2	2.43	0.48
1:C:274:LEU:HD12	1:C:317:ILE:HD12	1.96	0.47
2:B:417:ARG:HD3	4:E:501:HOH:O	2.14	0.47
1:A:320:CYS:SG	3:E:406:ASA:O	2.72	0.46
1:C:190:LEU:HD23	2:D:442:HIS:CD2	2.47	0.46
1:C:194:LEU:CD2	1:C:199:ARG:HD2	2.45	0.46

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:517:HOH:O	2:D:356:MET:HG3	2.15	0.46
1:C:200:GLY:HA2	1:C:267:THR:HB	1.98	0.45
2:B:378:ARG:HA	2:B:384:SER:HA	1.98	0.45
2:B:421:ALA:O	2:B:424:THR:HG22	2.18	0.44
1:C:205:LEU:HD23	1:C:243:LEU:CB	2.48	0.44
1:C:306:PRO:HA	1:C:309:GLN:HG3	2.00	0.43
1:C:205:LEU:CD2	1:C:243:LEU:CB	2.96	0.43
1:A:179[A]:CYS:SG	2:B:449:HIS:HB2	2.59	0.42
2:D:392[B]:GLN:HG3	4:D:522:HOH:O	2.20	0.42
2:B:417:ARG:HG2	2:B:418:GLU:N	2.34	0.41
2:B:417:ARG:HG2	2:B:418:GLU:H	1.86	0.41
1:A:248:ALA:HA	1:A:251:MET:HE2	2.03	0.40
1:C:229:LEU:HD11	2:D:387:ILE:HD12	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	152/160 (95%)	149 (98%)	3 (2%)	0	100	100
1	C	158/160 (99%)	154 (98%)	4 (2%)	0	100	100
2	B	99/112 (88%)	96 (97%)	3 (3%)	0	100	100
2	D	98/112 (88%)	96 (98%)	2 (2%)	0	100	100
3	E	4/6 (67%)	4 (100%)	0	0	100	100
3	F	4/6 (67%)	4 (100%)	0	0	100	100
All	All	515/556 (93%)	503 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	126/140 (90%)	125 (99%)	1 (1%)	81	76
1	C	130/140 (93%)	126 (97%)	4 (3%)	40	22
2	B	81/96 (84%)	80 (99%)	1 (1%)	71	62
2	D	83/96 (86%)	83 (100%)	0	100	100
3	E	2/2 (100%)	2 (100%)	0	100	100
3	F	2/2 (100%)	2 (100%)	0	100	100
All	All	424/476 (89%)	418 (99%)	6 (1%)	71	56

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	194	LEU
2	B	416	ASP
1	C	194	LEU
1	C	325	THR
1	C	332[A]	GLN
1	C	332[B]	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	257	ASN
1	A	332	GLN
1	C	257	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	ASA	F	406	3	3,7,7	0.82	0	1,8,8	0.50	0
3	ASA	E	406	3	3,7,7	0.74	0	1,8,8	0.22	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ASA	F	406	3	-	3/3/6/6	-
3	ASA	E	406	3	-	2/3/6/6	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	F	406	ASA	O-C-CA-CB
3	F	406	ASA	C-CA-CB-CG
3	E	406	ASA	C-CA-CB-CG
3	E	406	ASA	N-CA-CB-CG
3	F	406	ASA	N-CA-CB-CG

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	F	406	ASA	3	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	E	406	ASA	2	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	154/160 (96%)	0.18	5 (3%) 47 46	15, 26, 37, 43	3 (1%)
1	C	159/160 (99%)	0.13	5 (3%) 49 47	15, 26, 37, 48	2 (1%)
2	B	97/112 (86%)	0.89	16 (16%) 1 1	13, 23, 47, 51	0
2	D	97/112 (86%)	0.43	11 (11%) 5 4	12, 21, 36, 44	1 (1%)
3	E	4/6 (66%)	1.53	1 (25%) 0 0	40, 42, 43, 44	0
3	F	4/6 (66%)	0.12	0 100 100	27, 29, 32, 33	0
All	All	515/556 (92%)	0.35	38 (7%) 14 14	12, 25, 40, 51	6 (1%)

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	421	ALA	5.8
2	B	451	PRO	5.6
2	B	420	TYR	4.9
1	A	179[A]	CYS	4.7
2	D	451	PRO	4.2
2	B	424	THR	4.1
1	C	174	MET	3.8
2	B	380	THR	3.6
2	B	423	GLY	3.4
2	B	422	PRO	3.4
2	D	440	CYS	3.3
2	D	355	LYS	3.2
2	D	436	CYS	3.1
2	D	365	ILE	3.1
2	D	356	MET	2.9
2	B	365	ILE	2.9
1	A	216	LEU	2.9
2	B	435	TYR	2.9
1	C	332[A]	GLN	2.9

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	B	450	PRO	2.7
3	E	402	ALA	2.6
2	B	407	LEU	2.6
2	D	435	TYR	2.6
2	B	426	PHE	2.5
1	C	213	GLU	2.4
2	D	439	LEU	2.4
1	A	174	MET	2.4
2	D	403	VAL	2.3
2	D	437	SER	2.3
2	B	355	LYS	2.3
1	C	214	LYS	2.2
1	C	197	ARG	2.1
2	B	408	VAL	2.1
2	D	407	LEU	2.1
2	B	366[A]	CYS	2.1
1	A	180	THR	2.0
1	A	181	PRO	2.0
2	B	436	CYS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	ASA	E	406	8/8	0.79	0.25	45,46,47,47	0
3	ASA	F	406	8/8	0.92	0.11	32,34,34,35	0

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers

There are no such residues in this entry.