



# wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 21, 2024 – 04:10 pm BST

PDB ID : 8R6L  
Title : Crystal structure of *Candida glabrata* Bdf1 bromodomain 2 in the unbound state  
Authors : Petosa, C.; Wei, K.; McKenna, C.E.; Govin, J.  
Deposited on : 2023-11-22  
Resolution : 1.96 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.003 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*






The reported resolution of this entry is 1.96 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	3187 (1.96-1.96)
Clashscore	180529	3412 (1.96-1.96)
Ramachandran outliers	177936	3390 (1.96-1.96)
Sidechain outliers	177891	3390 (1.96-1.96)
RSRZ outliers	164620	3186 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	118	 3% 92% 5% ..
1	B	118	 2% 96% ..
1	C	118	 3% 94% ..
1	D	118	 3% 89% 5% 6%
1	E	118	 3% 93% ..

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	118	 3% 92% 6% •
1	G	118	 3% 92% 6% •

## 2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 6973 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Candida glabrata strain CBS138 chromosome C complete sequence.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	116	950	609	156	176	9	0	0	0
1	B	117	959	614	157	179	9	0	0	0
1	C	116	950	609	156	176	9	0	0	0
1	D	111	909	582	151	167	9	0	0	0
1	E	116	950	609	156	176	9	0	0	0
1	F	116	950	609	156	176	9	0	0	0
1	G	116	950	609	156	176	9	0	0	0

There are 21 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	301	ALA	-	expression tag	UNP Q6FWV7
A	302	MET	-	expression tag	UNP Q6FWV7
A	303	GLY	-	expression tag	UNP Q6FWV7
B	301	ALA	-	expression tag	UNP Q6FWV7
B	302	MET	-	expression tag	UNP Q6FWV7
B	303	GLY	-	expression tag	UNP Q6FWV7
C	301	ALA	-	expression tag	UNP Q6FWV7
C	302	MET	-	expression tag	UNP Q6FWV7
C	303	GLY	-	expression tag	UNP Q6FWV7
D	301	ALA	-	expression tag	UNP Q6FWV7
D	302	MET	-	expression tag	UNP Q6FWV7
D	303	GLY	-	expression tag	UNP Q6FWV7
E	301	ALA	-	expression tag	UNP Q6FWV7
E	302	MET	-	expression tag	UNP Q6FWV7

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	303	GLY	-	expression tag	UNP Q6FWV7
F	301	ALA	-	expression tag	UNP Q6FWV7
F	302	MET	-	expression tag	UNP Q6FWV7
F	303	GLY	-	expression tag	UNP Q6FWV7
G	301	ALA	-	expression tag	UNP Q6FWV7
G	302	MET	-	expression tag	UNP Q6FWV7
G	303	GLY	-	expression tag	UNP Q6FWV7

- Molecule 2 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: C<sub>6</sub>H<sub>14</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 8 6 2	0	0
2	A	1	Total C O 8 6 2	0	0
2	B	1	Total C O 8 6 2	0	0
2	B	1	Total C O 8 6 2	0	0
2	C	1	Total C O 8 6 2	0	0
2	D	1	Total C O 8 6 2	0	0
2	D	1	Total C O 8 6 2	0	0

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	E	1	8	6	2	0	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
3	A	1	6	3	3	0	0
3	A	1	6	3	3	0	0
3	A	1	6	3	3	0	0
3	A	1	6	3	3	0	0
3	A	1	6	3	3	0	0
3	B	1	6	3	3	0	0
3	B	1	6	3	3	0	0
3	B	1	6	3	3	0	0
3	C	1	6	3	3	0	0
3	C	1	6	3	3	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	D	1	Total	C	O	0	0
			6	3	3		
3	E	1	Total	C	O	0	0
			6	3	3		
3	E	1	Total	C	O	0	0
			6	3	3		
3	E	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is ACETATE ION (three-letter code: ACT) (formula: C<sub>2</sub>H<sub>3</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		
4	E	1	Total	C	O	0	0
			4	2	2		
4	F	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	63	Total O 63 63	0	0
5	B	38	Total O 38 38	0	0
5	C	30	Total O 30 30	0	0
5	D	25	Total O 25 25	0	0
5	E	22	Total O 22 22	0	0
5	G	1	Total O 1 1	0	0



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: *Candida glabrata* strain CBS138 chromosome C complete sequence




- Molecule 1: *Candida glabrata* strain CBS138 chromosome C complete sequence




- Molecule 1: *Candida glabrata* strain CBS138 chromosome C complete sequence




- Molecule 1: *Candida glabrata* strain CBS138 chromosome C complete sequence




- Molecule 1: *Candida glabrata* strain CBS138 chromosome C complete sequence



- Molecule 1: *Candida glabrata* strain CBS138 chromosome C complete sequence



- Molecule 1: *Candida glabrata* strain CBS138 chromosome C complete sequence



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	80.29Å 67.69Å 96.62Å 90.00° 95.52° 90.00°	Depositor
Resolution (Å)	44.37 – 1.96 44.37 – 1.96	Depositor EDS
% Data completeness (in resolution range)	98.9 (44.37-1.96) 99.0 (44.37-1.96)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.68 (at 1.73Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, $R_{free}$	0.200 , 0.227 0.195 , 0.223	Depositor DCC
$R_{free}$ test set	3616 reflections (4.92%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	40.5	Xtrriage
Anisotropy	0.392	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 48.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	6973	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	76.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.08% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MPD, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.80	1/976 (0.1%)	0.69	0/1322
1	B	0.64	0/985	0.65	0/1334
1	C	0.46	0/976	0.56	1/1322 (0.1%)
1	D	0.52	0/934	0.56	0/1264
1	E	0.39	0/976	0.51	0/1322
1	F	0.30	0/976	0.44	0/1322
1	G	0.31	0/976	0.47	0/1322
All	All	0.52	1/6799 (0.0%)	0.56	1/9208 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	310	CYS	CB-SG	-5.25	1.73	1.81

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	360	LEU	CA-CB-CG	5.21	127.29	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	950	0	909	4	0
1	B	959	0	915	2	0
1	C	950	0	909	2	0
1	D	909	0	871	4	0
1	E	950	0	909	5	0
1	F	950	0	909	4	0
1	G	950	0	909	5	0
2	A	16	0	28	0	0
2	B	16	0	28	0	0
2	C	8	0	14	0	0
2	D	16	0	28	2	0
2	E	8	0	14	1	0
3	A	30	0	39	0	0
3	B	18	0	23	0	0
3	C	12	0	16	0	0
3	D	6	0	8	0	0
3	E	18	0	24	1	0
4	A	8	0	6	1	0
4	B	8	0	6	0	0
4	C	4	0	3	0	0
4	E	4	0	3	0	0
4	F	4	0	3	0	0
5	A	63	0	0	0	0
5	B	38	0	0	0	0
5	C	30	0	0	0	0
5	D	25	0	0	0	0
5	E	22	0	0	0	0
5	G	1	0	0	1	0
All	All	6973	0	6574	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 24 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:398:ARG:HH11	2:D:902:MPD:H31	1.43	0.81
1:D:398:ARG:NH1	2:D:902:MPD:H31	2.14	0.61
1:G:412:LEU:O	5:G:501:HOH:O	2.18	0.56
1:E:310:CYS:HB3	1:E:360:LEU:HD21	1.91	0.51
1:A:348:LYS:NZ	4:A:904:ACT:OXT	2.39	0.49

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	114/118 (97%)	113 (99%)	1 (1%)	0	100	100
1	B	115/118 (98%)	113 (98%)	2 (2%)	0	100	100
1	C	114/118 (97%)	112 (98%)	2 (2%)	0	100	100
1	D	109/118 (92%)	108 (99%)	1 (1%)	0	100	100
1	E	114/118 (97%)	112 (98%)	2 (2%)	0	100	100
1	F	114/118 (97%)	113 (99%)	1 (1%)	0	100	100
1	G	114/118 (97%)	113 (99%)	1 (1%)	0	100	100
All	All	794/826 (96%)	784 (99%)	10 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	104/106 (98%)	104 (100%)	0	100	100
1	B	105/106 (99%)	105 (100%)	0	100	100
1	C	104/106 (98%)	104 (100%)	0	100	100
1	D	99/106 (93%)	98 (99%)	1 (1%)	73	72
1	E	104/106 (98%)	103 (99%)	1 (1%)	73	72

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	F	104/106 (98%)	103 (99%)	1 (1%)	73	72
1	G	104/106 (98%)	104 (100%)	0	100	100
All	All	724/742 (98%)	721 (100%)	3 (0%)	89	89

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	388	ASP
1	E	310	CYS
1	F	374	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	380	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

29 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	MPD	D	902	-	7,7,7	0.35	0	9,10,10	0.62	0
2	MPD	E	901	-	7,7,7	0.38	0	9,10,10	0.39	0
4	ACT	A	904	-	3,3,3	2.10	1 (33%)	3,3,3	1.40	0
2	MPD	C	901	-	7,7,7	0.29	0	9,10,10	0.24	0
2	MPD	A	901	-	7,7,7	0.40	0	9,10,10	0.34	0
3	GOL	A	903	-	5,5,5	1.01	0	5,5,5	0.94	0
3	GOL	C	903	-	5,5,5	0.94	0	5,5,5	1.13	0
3	GOL	A	909	-	5,5,5	1.15	1 (20%)	5,5,5	0.76	0
4	ACT	F	501	-	3,3,3	1.28	0	3,3,3	1.52	1 (33%)
3	GOL	A	906	-	5,5,5	1.27	1 (20%)	5,5,5	1.22	1 (20%)
3	GOL	C	902	-	5,5,5	1.10	0	5,5,5	0.84	0
2	MPD	B	501	-	7,7,7	0.48	0	9,10,10	1.00	0
2	MPD	B	503	-	7,7,7	0.29	0	9,10,10	0.32	0
4	ACT	B	504	-	3,3,3	1.34	0	3,3,3	1.39	0
2	MPD	A	902	-	7,7,7	0.26	0	9,10,10	0.68	0
4	ACT	C	904	-	3,3,3	1.31	0	3,3,3	1.31	0
3	GOL	E	903	-	5,5,5	1.16	1 (20%)	5,5,5	1.16	0
3	GOL	B	505	-	5,5,5	1.00	0	5,5,5	0.94	0
3	GOL	D	903	-	5,5,5	0.88	0	5,5,5	1.07	0
3	GOL	E	902	-	5,5,5	1.00	0	5,5,5	0.89	0
4	ACT	B	507	-	3,3,3	1.52	1 (33%)	3,3,3	1.44	1 (33%)
3	GOL	A	908	-	5,5,5	1.00	0	5,5,5	0.98	0
3	GOL	B	502	-	5,5,5	1.42	1 (20%)	5,5,5	0.79	0
3	GOL	A	905	-	5,5,5	1.14	1 (20%)	5,5,5	1.11	0
3	GOL	B	506	-	5,5,5	0.90	0	5,5,5	1.06	0
4	ACT	A	907	-	3,3,3	1.18	0	3,3,3	1.22	0
4	ACT	E	905	-	3,3,3	1.16	0	3,3,3	1.41	0
2	MPD	D	901	-	7,7,7	0.30	0	9,10,10	0.32	0
3	GOL	E	904	-	5,5,5	0.97	0	5,5,5	0.92	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MPD	D	902	-	-	3/5/5/5	-
2	MPD	E	901	-	-	1/5/5/5	-

*Continued on next page...*



Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MPD	C	901	-	-	0/5/5/5	-
2	MPD	A	901	-	-	1/5/5/5	-
3	GOL	A	903	-	-	2/4/4/4	-
3	GOL	C	903	-	-	2/4/4/4	-
3	GOL	A	909	-	-	0/4/4/4	-
3	GOL	A	906	-	-	0/4/4/4	-
3	GOL	C	902	-	-	4/4/4/4	-
2	MPD	B	501	-	-	0/5/5/5	-
2	MPD	B	503	-	-	3/5/5/5	-
2	MPD	A	902	-	-	1/5/5/5	-
3	GOL	E	903	-	-	3/4/4/4	-
3	GOL	B	505	-	-	0/4/4/4	-
3	GOL	D	903	-	-	4/4/4/4	-
3	GOL	E	902	-	-	0/4/4/4	-
3	GOL	A	908	-	-	2/4/4/4	-
3	GOL	B	502	-	-	2/4/4/4	-
3	GOL	A	905	-	-	3/4/4/4	-
3	GOL	B	506	-	-	0/4/4/4	-
2	MPD	D	901	-	-	0/5/5/5	-
3	GOL	E	904	-	-	2/4/4/4	-

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	904	ACT	CH3-C	3.34	1.63	1.49
3	B	502	GOL	O2-C2	-2.67	1.35	1.43
3	A	906	GOL	C1-C2	2.44	1.61	1.51
3	A	909	GOL	O2-C2	-2.20	1.36	1.43
3	A	905	GOL	C3-C2	2.15	1.60	1.51

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	906	GOL	C3-C2-C1	-2.35	102.56	111.70
4	F	501	ACT	OXT-C-O	2.04	129.58	122.05
4	B	507	ACT	OXT-C-O	2.01	129.46	122.05

There are no chirality outliers.

5 of 33 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	902	GOL	C1-C2-C3-O3
3	D	903	GOL	C1-C2-C3-O3
3	E	904	GOL	O1-C1-C2-C3
3	E	904	GOL	O1-C1-C2-O2
3	A	903	GOL	O1-C1-C2-C3

There are no ring outliers.

4 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	902	MPD	2	0
2	E	901	MPD	1	0
4	A	904	ACT	1	0
3	E	903	GOL	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	116/118 (98%)	-0.12	4 (3%) 48 55	32, 44, 68, 73	0
1	B	117/118 (99%)	0.09	2 (1%) 69 75	33, 53, 82, 93	0
1	C	116/118 (98%)	0.23	3 (2%) 57 63	35, 70, 105, 120	0
1	D	111/118 (94%)	0.47	4 (3%) 46 54	44, 75, 120, 133	0
1	E	116/118 (98%)	0.45	4 (3%) 48 55	45, 85, 127, 137	0
1	F	116/118 (98%)	0.66	4 (3%) 48 55	76, 97, 115, 126	0
1	G	116/118 (98%)	0.57	4 (3%) 48 55	78, 102, 120, 127	0
All	All	808/826 (97%)	0.33	25 (3%) 51 58	32, 78, 118, 137	0

The worst 5 of 25 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	411	PRO	4.9
1	F	301	ALA	4.4
1	E	301	ALA	4.3
1	G	301	ALA	4.0
1	D	301	ALA	3.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	ACT	A	904	4/4	0.73	0.25	62,66,68,69	0
4	ACT	B	504	4/4	0.74	0.17	87,89,89,89	0
4	ACT	C	904	4/4	0.82	0.18	93,93,94,94	0
4	ACT	F	501	4/4	0.82	0.13	83,85,86,88	0
3	GOL	A	905	6/6	0.83	0.14	51,55,55,55	0
3	GOL	B	506	6/6	0.83	0.16	72,73,74,75	0
3	GOL	E	904	6/6	0.84	0.10	98,99,99,99	0
4	ACT	B	507	4/4	0.85	0.18	71,77,77,79	0
3	GOL	D	903	6/6	0.85	0.11	52,55,57,59	0
2	MPD	A	902	8/8	0.85	0.22	84,87,90,90	0
3	GOL	A	903	6/6	0.86	0.14	92,94,96,98	0
3	GOL	A	908	6/6	0.86	0.15	85,88,88,89	0
3	GOL	E	903	6/6	0.87	0.10	49,53,57,57	0
3	GOL	A	909	6/6	0.87	0.16	72,78,79,80	0
3	GOL	E	902	6/6	0.87	0.14	97,98,98,99	0
4	ACT	A	907	4/4	0.87	0.14	46,55,55,62	0
2	MPD	B	503	8/8	0.88	0.17	85,87,90,91	0
4	ACT	E	905	4/4	0.88	0.17	100,101,102,102	0
3	GOL	C	903	6/6	0.88	0.11	43,53,55,55	0
3	GOL	B	502	6/6	0.89	0.18	67,75,77,81	0
3	GOL	C	902	6/6	0.89	0.14	77,78,79,80	0
3	GOL	A	906	6/6	0.90	0.12	51,52,54,54	0
2	MPD	D	902	8/8	0.91	0.21	92,94,95,96	0
3	GOL	B	505	6/6	0.93	0.09	74,80,83,86	0
2	MPD	D	901	8/8	0.93	0.13	64,71,74,76	0
2	MPD	E	901	8/8	0.93	0.12	69,77,81,83	0
2	MPD	B	501	8/8	0.94	0.10	38,45,48,50	0
2	MPD	A	901	8/8	0.94	0.12	48,56,63,66	0
2	MPD	C	901	8/8	0.95	0.12	61,66,69,72	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.