



wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 24, 2024 – 12:13 am GMT

PDB ID : 8R70
Title : Polysaccharide lyase BtPL33HA (BT4410) Y291A with HA dp4
Authors : Cartmell, A.
Deposited on : 2023-11-23
Resolution : 1.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.40


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	7108 (1.80-1.80)
Clashscore	180529	8162 (1.80-1.80)
Ramachandran outliers	177936	8077 (1.80-1.80)
Sidechain outliers	177891	8076 (1.80-1.80)
RSRZ outliers	164620	7108 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	644	 86% 9% . .
1	B	644	 83% 10% . 6%
2	C	3	 100%
3	D	4	 100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	MES	A	702	-	-	X	-

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 20540 atoms, of which 9811 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Heparinase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	618	9900	3178	4901	855	932	34	123	3	0
1	B	608	9703	3117	4802	835	915	34	120	2	0

There are 54 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	initiating methionine	UNP Q89ZG7
A	2	GLY	-	expression tag	UNP Q89ZG7
A	3	SER	-	expression tag	UNP Q89ZG7
A	4	SER	-	expression tag	UNP Q89ZG7
A	5	HIS	-	expression tag	UNP Q89ZG7
A	6	HIS	-	expression tag	UNP Q89ZG7
A	7	HIS	-	expression tag	UNP Q89ZG7
A	8	HIS	-	expression tag	UNP Q89ZG7
A	9	HIS	-	expression tag	UNP Q89ZG7
A	10	HIS	-	expression tag	UNP Q89ZG7
A	11	SER	-	expression tag	UNP Q89ZG7
A	12	SER	-	expression tag	UNP Q89ZG7
A	13	GLY	-	expression tag	UNP Q89ZG7
A	14	LEU	-	expression tag	UNP Q89ZG7
A	15	VAL	-	expression tag	UNP Q89ZG7
A	16	PRO	-	expression tag	UNP Q89ZG7
A	17	ARG	-	expression tag	UNP Q89ZG7
A	18	GLY	-	expression tag	UNP Q89ZG7
A	19	SER	-	expression tag	UNP Q89ZG7
A	20	HIS	-	expression tag	UNP Q89ZG7
A	21	MET	-	expression tag	UNP Q89ZG7
A	22	ALA	-	expression tag	UNP Q89ZG7
A	23	SER	-	expression tag	UNP Q89ZG7
A	24	ASN	-	expression tag	UNP Q89ZG7
A	25	ALA	-	expression tag	UNP Q89ZG7

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	26	TYR	-	expression tag	UNP Q89ZG7
A	291	ALA	TYR	engineered mutation	UNP Q89ZG7
B	1	MET	-	initiating methionine	UNP Q89ZG7
B	2	GLY	-	expression tag	UNP Q89ZG7
B	3	SER	-	expression tag	UNP Q89ZG7
B	4	SER	-	expression tag	UNP Q89ZG7
B	5	HIS	-	expression tag	UNP Q89ZG7
B	6	HIS	-	expression tag	UNP Q89ZG7
B	7	HIS	-	expression tag	UNP Q89ZG7
B	8	HIS	-	expression tag	UNP Q89ZG7
B	9	HIS	-	expression tag	UNP Q89ZG7
B	10	HIS	-	expression tag	UNP Q89ZG7
B	11	SER	-	expression tag	UNP Q89ZG7
B	12	SER	-	expression tag	UNP Q89ZG7
B	13	GLY	-	expression tag	UNP Q89ZG7
B	14	LEU	-	expression tag	UNP Q89ZG7
B	15	VAL	-	expression tag	UNP Q89ZG7
B	16	PRO	-	expression tag	UNP Q89ZG7
B	17	ARG	-	expression tag	UNP Q89ZG7
B	18	GLY	-	expression tag	UNP Q89ZG7
B	19	SER	-	expression tag	UNP Q89ZG7
B	20	HIS	-	expression tag	UNP Q89ZG7
B	21	MET	-	expression tag	UNP Q89ZG7
B	22	ALA	-	expression tag	UNP Q89ZG7
B	23	SER	-	expression tag	UNP Q89ZG7
B	24	ASN	-	expression tag	UNP Q89ZG7
B	25	ALA	-	expression tag	UNP Q89ZG7
B	26	TYR	-	expression tag	UNP Q89ZG7
B	291	ALA	TYR	engineered mutation	UNP Q89ZG7

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-beta-D-glucopyranuronic acid-(1-3)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	H	N	O			
2	C	3	71	22	30	2	17	6	0	0


- Molecule 3 is an oligosaccharide called beta-D-glucopyranuronic acid-(1-3)-2-acetamido-2-d

eoxy-beta-D-glucopyranose-(1-4)-beta-D-glucopyranuronic acid-(1-3)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
			Total	C	H	N				O
3	D	4	90	28	37	2	23	8	0	0

- Molecule 4 is polyethylene glycol (three-letter code: P4K) (formula: C₃₀H₆₂O₁₅).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
4	A	1	28	8	16	4	1	0
4	B	1	21	6	12	3	1	0

- Molecule 5 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: C₆H₁₃NO₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	H	N	O			S
5	A	1	25	6	13	1	4	1	0	0

- Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Zn		
6	A	1	1	1	0	0
6	B	1	1	1	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
7	A	434	434	434	0	0
7	B	266	266	266	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Heparinase



- Molecule 1: Heparinase



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-beta-D-glucopyranuronic acid-(1-3)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: beta-D-glucopyranuronic acid-(1-3)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-beta-D-glucopyranuronic acid-(1-3)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D:

100%

NAG1
BDF2
NAG3
BDF4

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	53.64Å 137.30Å 202.84Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	113.70 – 1.80 113.70 – 1.80	Depositor EDS
% Data completeness (in resolution range)	87.0 (113.70-1.80) 92.0 (113.70-1.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.50 (at 1.78Å)	Xtrriage
Refinement program	REFMAC 5.8.0419	Depositor
R, R_{free}	0.181 , 0.223 0.183 , 0.224	Depositor DCC
R_{free} test set	7064 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	31.9	Xtrriage
Anisotropy	0.335	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 30.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	20540	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.03% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, NAG, P4K, MES, BDP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.72	2/5120 (0.0%)	1.17	26/6927 (0.4%)
1	B	0.62	0/5019	1.13	25/6792 (0.4%)
All	All	0.67	2/10139 (0.0%)	1.15	51/13719 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	7
1	B	0	6
All	All	0	13

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	260	ARG	NE-CZ	5.27	1.40	1.33
1	A	327	GLU	CD-OE1	-5.19	1.20	1.25

The worst 5 of 51 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	60	ARG	NE-CZ-NH1	16.69	128.65	120.30
1	B	254	MET	CG-SD-CE	13.56	121.89	100.20
1	A	60	ARG	NE-CZ-NH2	-12.80	113.90	120.30
1	B	324	ARG	NE-CZ-NH2	-12.12	114.24	120.30
1	B	324	ARG	NE-CZ-NH1	11.44	126.02	120.30

There are no chirality outliers.

5 of 13 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	114	ARG	Sidechain
1	A	260	ARG	Sidechain
1	A	267[A]	ARG	Sidechain
1	A	29	ARG	Sidechain
1	A	60	ARG	Sidechain

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4999	4901	4889	39	1
1	B	4901	4802	4790	35	1
2	C	41	30	33	0	0
3	D	53	37	40	0	0
4	A	12	16	0	1	0
4	B	9	12	0	0	0
5	A	12	13	13	11	0
6	A	1	0	0	0	0
6	B	1	0	0	0	0
7	A	434	0	0	11	0
7	B	266	0	0	11	0
All	All	10729	9811	9765	73	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 73 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:593:ASP:HB3	7:B:1042:HOH:O	1.41	1.20
1:B:419:GLU:HG3	7:B:829:HOH:O	1.48	1.11
1:A:572:ASN:HB3	7:A:1088:HOH:O	1.48	1.10
1:A:327:GLU:OE1	4:A:701:P4K:O9	1.74	1.03
1:A:277:ASN:HD22	5:A:702:MES:H72	1.45	0.82

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the sym-

metry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:577:SER:HG	1:B:599:LYS:HZ3[2_454]	1.27	0.33

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	619/644 (96%)	595 (96%)	24 (4%)	0	100	100
1	B	606/644 (94%)	579 (96%)	27 (4%)	0	100	100
All	All	1225/1288 (95%)	1174 (96%)	51 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	533/551 (97%)	522 (98%)	11 (2%)	48	38
1	B	523/551 (95%)	509 (97%)	14 (3%)	40	28
All	All	1056/1102 (96%)	1031 (98%)	25 (2%)	44	32

5 of 25 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	88	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	344	ASP
1	B	599	LYS
1	B	179	TYR
1	B	359	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	510	ASN
1	B	231	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

7 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	C	1	2	15,15,15	1.84	3 (20%)	21,21,21	1.96	8 (38%)
2	BDP	C	2	2	12,12,13	1.77	4 (33%)	14,17,19	2.38	3 (21%)
2	NAG	C	3	2	14,14,15	2.98	9 (64%)	17,19,21	2.83	6 (35%)
3	NAG	D	1	3	15,15,15	1.35	3 (20%)	21,21,21	0.99	0
3	BDP	D	2	3	12,12,13	2.48	6 (50%)	14,17,19	2.30	6 (42%)
3	NAG	D	3	3	14,14,15	3.04	9 (64%)	17,19,21	2.83	9 (52%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	BDP	D	4	3	12,12,13	3.62	7 (58%)	14,17,19	3.21	10 (71%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	2	-	2/6/26/26	0/1/1/1
2	BDP	C	2	2	-	1/4/21/24	0/1/1/1
2	NAG	C	3	2	-	2/6/23/26	0/1/1/1
3	NAG	D	1	3	-	0/6/26/26	0/1/1/1
3	BDP	D	2	3	-	0/4/21/24	0/1/1/1
3	NAG	D	3	3	-	2/6/23/26	0/1/1/1
3	BDP	D	4	3	-	3/4/21/24	0/1/1/1

The worst 5 of 41 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	3	NAG	C8-C7	-7.52	1.34	1.50
2	C	3	NAG	C8-C7	-7.36	1.35	1.50
3	D	4	BDP	O5-C1	6.46	1.54	1.43
3	D	4	BDP	C5-C6	5.28	1.64	1.53
3	D	4	BDP	O6A-C6	5.24	1.38	1.22

The worst 5 of 42 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	3	NAG	C8-C7-N2	7.47	128.75	116.10
2	C	2	BDP	C1-C2-C3	6.30	117.41	109.67
2	C	3	NAG	O7-C7-N2	-5.97	110.97	121.95
3	D	4	BDP	O5-C5-C6	5.85	125.56	106.31
3	D	3	NAG	O3-C3-C2	5.05	119.91	109.47

There are no chirality outliers.

5 of 10 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	D	4	BDP	C4-C5-C6-O6A
3	D	4	BDP	C4-C5-C6-O6B

Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	C	3	NAG	C8-C7-N2-C2
2	C	3	NAG	O7-C7-N2-C2
3	D	3	NAG	C8-C7-N2-C2

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 2 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	P4K	B	701	-	8,8,44	0.54	0	7,7,43	0.67	0
5	MES	A	702	-	12,12,12	0.99	0	14,16,16	0.91	0
4	P4K	A	701	-	11,11,44	0.50	0	10,10,43	0.76	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	P4K	B	701	-	-	0/6/6/42	-
5	MES	A	702	-	-	6/6/14/14	0/1/1/1
4	P4K	A	701	-	-	5/9/9/42	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 11 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	A	702	MES	N4-C7-C8-S
5	A	702	MES	C7-C8-S-O3S
4	A	701	P4K	O8-C15-C16-O9
4	A	701	P4K	C14-C13-O7-C12
5	A	702	MES	C8-C7-N4-C3

There are no ring outliers.

2 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	702	MES	11	0
4	A	701	P4K	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	618/644 (95%)	-0.39	4 (0%) 85 85	13, 30, 54, 74	3 (0%)
1	B	608/644 (94%)	-0.03	11 (1%) 67 66	18, 38, 63, 96	2 (0%)
All	All	1226/1288 (95%)	-0.21	15 (1%) 76 76	13, 34, 60, 96	5 (0%)

The worst 5 of 15 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	481	THR	4.0
1	B	469	THR	3.7
1	B	482	ILE	2.9
1	B	26	TYR	2.9
1	A	291	ALA	2.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	BDP	D	4	12/13	0.64	0.18	30,78,93,97	2
3	NAG	D	3	14/15	0.71	0.16	30,64,70,72	2
2	NAG	C	3	14/15	0.78	0.14	30,49,58,75	2
3	BDP	D	2	12/13	0.85	0.11	30,51,60,67	2
3	NAG	D	1	15/15	0.89	0.10	30,58,66,67	2


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	NAG	C	1	15/15	0.89	0.10	30,41,49,59	2
2	BDP	C	2	12/13	0.93	0.08	27,36,51,59	2

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	MES	A	702	12/12	0.66	0.17	48,60,132,136	0
4	P4K	A	701	12/45	0.88	0.11	27,34,45,47	1
4	P4K	B	701	9/45	0.89	0.11	30,45,48,50	1
6	ZN	B	702	1/1	0.99	0.03	35,35,35,35	0
6	ZN	A	703	1/1	1.00	0.02	30,30,30,30	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.