



# wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 3, 2024 – 12:06 am GMT

PDB ID : 8R96  
Title : Crystal structure of the cryorhodopsin CryoR2 at pH 4.6, type A crystals  
Authors : Kovalev, K.; Lamm, G.H.U.; Marin, E.  
Deposited on : 2023-11-30  
Resolution : 2.46 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 3.0  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.003 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.40


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.46 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	1096 (2.46-2.46)
Clashscore	180529	1178 (2.46-2.46)
Ramachandran outliers	177936	1170 (2.46-2.46)
Sidechain outliers	177891	1170 (2.46-2.46)
RSRZ outliers	164620	1096 (2.46-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	327	 3% 73% 14% • 12%
1	B	327	 3% 72% 15% • 11%
1	C	327	 % 69% 17% • 12%
1	D	327	 2% 72% 16% • 12%
1	E	327	 3% 70% 16% • 13%

## 2 Entry composition [i](#)


There are 6 unique types of molecules in this entry. The entry contains 11044 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called cryorhodopsin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	288	Total 2155	C 1415	N 350	O 384	S 6	0	0	0
1	B	290	Total 2162	C 1419	N 349	O 388	S 6	0	0	0
1	C	287	Total 2138	C 1408	N 344	O 380	S 6	0	0	0
1	D	289	Total 2140	C 1409	N 341	O 384	S 6	0	0	0
1	E	286	Total 2143	C 1408	N 347	O 382	S 6	0	0	0

- Molecule 2 is EICOSANE (three-letter code: LFA) (formula: C<sub>20</sub>H<sub>42</sub>).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total 11	C 11	0	0

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C 7 7	0	0
2	A	1	Total C 7 7	0	0
2	A	1	Total C 10 10	0	0
2	A	1	Total C 7 7	0	0
2	B	1	Total C 11 11	0	0
2	B	1	Total C 12 12	0	0
2	C	1	Total C 6 6	0	0
2	C	1	Total C 6 6	0	0
2	D	1	Total C 10 10	0	0
2	D	1	Total C 11 11	0	0
2	D	1	Total C 8 8	0	0
2	E	1	Total C 11 11	0	0
2	E	1	Total C 13 13	0	0

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	P	0	0
			5	4	1		
3	B	1	Total	O	P	0	0
			5	4	1		
3	C	1	Total	O	P	0	0
			5	4	1		
3	D	1	Total	O	P	0	0
			5	4	1		
3	E	1	Total	O	P	0	0
			5	4	1		

- Molecule 4 is RETINAL (three-letter code: RET) (formula: C<sub>20</sub>H<sub>28</sub>O) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C 20 20	0	0
4	B	1	Total C 20 20	0	0
4	C	1	Total C 20 20	0	0
4	D	1	Total C 20 20	0	0
4	E	1	Total C 20 20	0	0

- Molecule 5 is OLEIC ACID (three-letter code: OLA) (formula:  $C_{18}H_{34}O_2$ ).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	D	1	Total	C O	0	0
			20	18 2		

- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	4	Total	O	0	0
			4	4		
6	B	9	Total	O	0	0
			9	9		
6	C	6	Total	O	0	0
			6	6		
6	D	7	Total	O	0	0
			7	7		
6	E	5	Total	O	0	0
			5	5		

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: cryorhodopsin

Chain A: 



- Molecule 1: cryorhodopsin

Chain B: 



- Molecule 1: cryorhodopsin

Chain C: 





ARG  
THR  
LYS  
PRO  
PRO  
THR  
SER  
ALA  
ALA  
VAL  
ALA  
SER  
PRO  
GLU  
ALA  
LEU  
LEU  
PRO  
PRO  
ASP  
SER  
SER  
THR  
PRO  
PRO  
LEU  
ASP  
GLY  
GLY  
TYR

• Molecule 1: cryorhodopsin



MET ARG L3 T4 N5 V6 S7 V8 W10 Y24 V28 T40 Y41 I42 R58 L59 G60 M61 L62 Y69 I73 F76 A84 N95 I96 F97 S98 V108 P109 L110 L111 T157 L162 I166 V169 I173 V177 L178 L179 R180 A181 V182 R183

Q184 S185 P187 T188 L189 M196 W208 V210 V211 P212 I213 V214 L217 P218 A222 V241 K242 L243 G244 F245 S246 V252 R256 Y289 L290 A291 GLU GLY ALA VAL HIS ASP ARG THR LYS PRO THR SER ALA VAL ALA SER PRO THR

LEU  
PRO  
PRO  
LEU  
THR  
SER  
PRO  
LEU  
ASP  
GLY  
GLY  
TYR

• Molecule 1: cryorhodopsin



MET ARG LEU THR N5 V6 S7 V8 W9 Y24 V27 V28 T40 Y41 I42 R58 L59 G60 M61 L62 F76 A84 G85 G86 N95 I96 F97 S98 V108 P109 L110 L111 L122 G123 V124 Q125 T157 L162 I166 V169 I173 V177 L178 V182

L186 P187 T188 L189 E192 M196 V209 V210 V211 P212 I213 V214 L217 P218 L219 F220 G221 A222 V239 I240 V241 K242 L243 G244 F245 S246 R251 V252 L255 R256 K278 D281 E287 V288 Y289 L290 ALA GLU GLY ALA VAL HIS ASP ARG THR LYS PRO

PRO  
THR  
SER  
ALA  
ALA  
VAL  
ALA  
SER  
PRO  
GLU  
ALA  
PRO  
LEU  
PRO  
PRO  
ASP  
SER  
THR  
PRO  
LEU  
LEU  
ASP  
GLY  
GLY  
TYR

## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	85.56Å 295.67Å 185.50Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.46 20.00 – 2.46	Depositor EDS
% Data completeness (in resolution range)	65.4 (20.00-2.46) 65.6 (20.00-2.46)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.25 (at 2.47Å)	Xtrriage
Refinement program	REFMAC 5.8.0267	Depositor
R, $R_{free}$	0.239 , 0.254 0.246 , 0.263	Depositor DCC
$R_{free}$ test set	4229 reflections (4.89%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	66.5	Xtrriage
Anisotropy	0.050	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 47.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	11044	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	63.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.24% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, LFA, RET, OLA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.66	0/2203	0.66	0/3025
1	B	0.65	0/2210	0.66	0/3035
1	C	0.65	0/2186	0.66	0/3003
1	D	0.66	0/2188	0.66	0/3009
1	E	0.66	0/2191	0.66	0/3009
All	All	0.66	0/10978	0.66	0/15081

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2155	0	2198	36	0
1	B	2162	0	2199	38	0
1	C	2138	0	2176	46	0
1	D	2140	0	2168	45	0
1	E	2143	0	2187	46	0
2	A	42	0	79	0	0
2	B	23	0	44	0	0
2	C	12	0	22	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	29	0	55	1	0
2	E	24	0	46	0	0
3	A	5	0	0	0	0
3	B	5	0	0	0	0
3	C	5	0	0	0	0
3	D	5	0	0	0	0
3	E	5	0	0	0	0
4	A	20	0	27	5	0
4	B	20	0	27	4	0
4	C	20	0	27	4	0
4	D	20	0	27	5	0
4	E	20	0	27	4	0
5	D	20	0	33	1	0
6	A	4	0	0	0	0
6	B	9	0	0	0	0
6	C	6	0	0	0	0
6	D	7	0	0	0	0
6	E	5	0	0	0	0
All	All	11044	0	11342	213	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 213 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:31:LEU:O	1:C:34:VAL:HG12	1.49	1.09
1:E:5:ASN:HB2	1:E:157:THR:OG1	1.53	1.07
1:E:289:TYR:O	1:E:290:LEU:HD23	1.73	0.88
1:D:58:ARG:HD2	1:D:246:SER:HB2	1.62	0.81
1:E:5:ASN:HB2	1:E:157:THR:HG1	1.50	0.76

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	286/327 (88%)	276 (96%)	10 (4%)	0	100	100
1	B	288/327 (88%)	272 (94%)	16 (6%)	0	100	100
1	C	285/327 (87%)	272 (95%)	13 (5%)	0	100	100
1	D	287/327 (88%)	273 (95%)	14 (5%)	0	100	100
1	E	284/327 (87%)	272 (96%)	12 (4%)	0	100	100
All	All	1430/1635 (88%)	1365 (96%)	65 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	224/261 (86%)	219 (98%)	5 (2%)	47	62
1	B	224/261 (86%)	220 (98%)	4 (2%)	54	68
1	C	221/261 (85%)	217 (98%)	4 (2%)	54	68
1	D	221/261 (85%)	217 (98%)	4 (2%)	54	68
1	E	224/261 (86%)	219 (98%)	5 (2%)	47	62
All	All	1114/1305 (85%)	1092 (98%)	22 (2%)	50	65

5 of 22 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	61	MET
1	E	5	ASN
1	D	243	LEU
1	E	42	ILE
1	B	81	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	19	HIS
1	D	175	ASN
1	E	175	ASN
1	B	175	ASN
1	A	175	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

25 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	PO4	D	402	-	4,4,4	0.71	0	6,6,6	0.42	0
2	LFA	A	404	-	6,6,19	0.14	0	5,5,18	0.11	0
2	LFA	B	401	-	10,10,19	0.12	0	9,9,18	0.09	0
2	LFA	A	406	-	6,6,19	0.13	0	5,5,18	0.09	0
2	LFA	B	402	-	11,11,19	0.12	0	10,10,18	0.07	0
2	LFA	A	402	-	6,6,19	0.15	0	5,5,18	0.09	0
2	LFA	E	403	-	12,12,19	0.08	0	11,11,18	0.08	0
2	LFA	D	404	-	10,10,19	0.11	0	9,9,18	0.07	0
2	LFA	C	401	-	5,5,19	0.16	0	4,4,18	0.11	0
3	PO4	A	403	-	4,4,4	0.65	0	6,6,6	0.44	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	OLA	D	401	-	19,19,19	0.52	0	19,19,19	0.47	0
3	PO4	B	403	-	4,4,4	0.69	0	6,6,6	0.42	0
2	LFA	A	401	-	10,10,19	0.12	0	9,9,18	0.08	0
4	RET	C	404	1	20,20,21	1.73	3 (15%)	27,27,28	1.06	1 (3%)
4	RET	E	404	1	20,20,21	1.67	3 (15%)	27,27,28	1.07	1 (3%)
4	RET	D	406	1	20,20,21	1.69	3 (15%)	27,27,28	1.06	1 (3%)
4	RET	B	404	1	20,20,21	1.68	3 (15%)	27,27,28	1.06	1 (3%)
2	LFA	C	403	-	5,5,19	0.16	0	4,4,18	0.11	0
2	LFA	A	405	-	9,9,19	0.12	0	8,8,18	0.07	0
2	LFA	E	401	-	10,10,19	0.12	0	9,9,18	0.08	0
2	LFA	D	403	-	9,9,19	0.12	0	8,8,18	0.08	0
3	PO4	C	402	-	4,4,4	0.66	0	6,6,6	0.44	0
3	PO4	E	402	-	4,4,4	0.69	0	6,6,6	0.41	0
4	RET	A	407	1	20,20,21	1.65	3 (15%)	27,27,28	1.08	1 (3%)
2	LFA	D	405	-	7,7,19	0.13	0	6,6,18	0.09	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	LFA	A	404	-	-	0/4/4/17	-
2	LFA	B	401	-	-	4/8/8/17	-
2	LFA	A	406	-	-	2/4/4/17	-
2	LFA	B	402	-	-	8/9/9/17	-
2	LFA	A	402	-	-	0/4/4/17	-
2	LFA	E	403	-	-	2/10/10/17	-
2	LFA	D	404	-	-	1/8/8/17	-
2	LFA	C	401	-	-	1/3/3/17	-
5	OLA	D	401	-	-	7/17/17/17	-
2	LFA	A	401	-	-	3/8/8/17	-
4	RET	C	404	1	-	0/13/30/31	0/1/1/1
4	RET	E	404	1	-	0/13/30/31	0/1/1/1
4	RET	D	406	1	-	0/13/30/31	0/1/1/1
4	RET	B	404	1	-	0/13/30/31	0/1/1/1
2	LFA	C	403	-	-	0/3/3/17	-
2	LFA	A	405	-	-	1/7/7/17	-
2	LFA	E	401	-	-	1/8/8/17	-
2	LFA	D	403	-	-	0/7/7/17	-

Continued on next page...

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	RET	A	407	1	-	0/13/30/31	0/1/1/1
2	LFA	D	405	-	-	1/5/5/17	-

The worst 5 of 15 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	404	RET	C10-C9	4.21	1.41	1.35
4	D	406	RET	C10-C9	4.08	1.41	1.35
4	B	404	RET	C10-C9	4.03	1.41	1.35
4	A	407	RET	C10-C9	3.99	1.41	1.35
4	D	406	RET	C14-C13	3.93	1.36	1.33

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	E	404	RET	C19-C9-C10	-3.97	117.36	122.92
4	A	407	RET	C19-C9-C10	-3.90	117.46	122.92
4	B	404	RET	C19-C9-C10	-3.88	117.48	122.92
4	D	406	RET	C19-C9-C10	-3.86	117.52	122.92
4	C	404	RET	C19-C9-C10	-3.69	117.75	122.92

There are no chirality outliers.

5 of 31 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	402	LFA	C5-C6-C7-C8
2	B	402	LFA	C7-C8-C9-C10
5	D	401	OLA	C13-C14-C15-C16
2	A	401	LFA	C3-C4-C5-C6
2	A	406	LFA	C15-C16-C17-C18

There are no ring outliers.

7 monomers are involved in 24 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	D	401	OLA	1	0
4	C	404	RET	4	0
4	E	404	RET	4	0
4	D	406	RET	5	0
4	B	404	RET	4	0
4	A	407	RET	5	0


*Continued on next page...*









*Continued from previous page...*


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	405	LFA	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	288/327 (88%)	0.19	10 (3%) 47 49	33, 62, 101, 134	0
1	B	290/327 (88%)	0.06	9 (3%) 51 54	30, 56, 96, 132	0
1	C	287/327 (87%)	0.10	4 (1%) 73 75	37, 60, 98, 119	0
1	D	289/327 (88%)	0.21	6 (2%) 63 65	38, 60, 98, 117	0
1	E	286/327 (87%)	0.18	9 (3%) 51 54	35, 61, 97, 117	0
All	All	1440/1635 (88%)	0.15	38 (2%) 57 60	30, 60, 99, 134	0

The worst 5 of 38 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	3	LEU	5.0
1	E	85	GLY	4.3
1	E	84	ALA	3.5
1	C	4	THR	3.4
1	C	5	ASN	3.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	LFA	D	403	10/20	0.56	0.13	44,85,89,89	0
2	LFA	D	405	8/20	0.56	0.22	44,56,65,66	0
2	LFA	C	403	6/20	0.72	0.10	37,54,60,62	0
2	LFA	D	404	11/20	0.75	0.13	45,75,86,86	0
2	LFA	A	402	7/20	0.78	0.14	40,58,65,68	0
2	LFA	A	406	7/20	0.78	0.12	61,66,73,78	0
3	PO4	C	402	5/5	0.80	0.10	102,106,117,124	0
3	PO4	A	403	5/5	0.82	0.11	88,100,108,114	0
2	LFA	E	401	11/20	0.82	0.12	33,47,52,53	0
2	LFA	E	403	13/20	0.83	0.11	70,79,96,98	0
2	LFA	A	404	7/20	0.83	0.11	34,41,50,52	0
2	LFA	A	405	10/20	0.83	0.13	40,67,69,71	0
3	PO4	E	402	5/5	0.83	0.11	88,91,104,105	0
2	LFA	B	401	11/20	0.84	0.11	34,46,55,56	0
3	PO4	D	402	5/5	0.84	0.10	92,92,99,99	0
2	LFA	A	401	11/20	0.84	0.14	44,70,72,73	0
3	PO4	B	403	5/5	0.85	0.08	90,92,102,103	0
2	LFA	C	401	6/20	0.87	0.12	30,40,55,57	0
5	OLA	D	401	20/20	0.87	0.15	71,78,95,98	0
2	LFA	B	402	12/20	0.88	0.15	44,68,80,81	0
4	RET	E	404	20/21	0.92	0.15	61,72,78,78	0
4	RET	A	407	20/21	0.93	0.15	58,72,86,87	0
4	RET	B	404	20/21	0.94	0.11	50,58,62,65	0
4	RET	D	406	20/21	0.94	0.12	58,63,73,75	0
4	RET	C	404	20/21	0.95	0.11	54,63,67,68	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around LFA D 405:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around LFA D 404:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around LFA A 406:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around LFA E 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around LFA E 403:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around LFA A 405:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around LFA B 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around LFA A 401:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around LFA C 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around OLA D 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around LFA B 402:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around RET E 404:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around RET A 407:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around RET B 404:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around RET D 406:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around RET C 404:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.