



Full wwPDB X-ray Structure Validation Report ⓘ

May 25, 2020 – 01:48 am BST

PDB ID : 5RAS
Title : PanDDA analysis group deposition – Crystal Structure of JMJD1B in complex with XS036302b
Authors : Snee, M.; Nowak, R.; Johansson, C.; Burgess-Brown, N.A.; Arrowsmith, C.H.; Bountra, C.; Edwards, A.M.; Oppermann, U.
Deposited on : 2020-03-16
Resolution : 1.58 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.58 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5534 (1.60-1.56)
Clashscore	141614	5861 (1.60-1.56)
Ramachandran outliers	138981	5708 (1.60-1.56)
Sidechain outliers	138945	5703 (1.60-1.56)
RSRZ outliers	127900	5431 (1.60-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.


Mol	Chain	Length	Quality of chain
1	A	372	
1	B	372	

ENTRY-COMPOSITION INFOmissingINFO


2 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Lysine-specific demethylase 3B



- Molecule 1: Lysine-specific demethylase 3B



3 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	58.03Å 93.63Å 93.80Å 90.00° 108.19° 90.00°	Depositor
Resolution (Å)	89.11 – 1.58 89.11 – 1.58	Depositor EDS
% Data completeness (in resolution range)	98.3 (89.11-1.58) 98.4 (89.11-1.58)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.66 (at 1.58Å)	Xtrriage
Refinement program	REFMAC 5.8.0238	Depositor
R, R_{free}	0.192 , 0.216 0.204 , 0.223	Depositor DCC
R_{free} test set	6410 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	22.5	Xtrriage
Anisotropy	0.065	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 46.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.024 for h,-k,-h-l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6293	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.71% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

4 Model quality [i](#)

4.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, MN, S3J

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.90	0/2841	0.95	5/3854 (0.1%)
1	B	0.85	1/2872 (0.0%)	0.87	2/3895 (0.1%)
All	All	0.87	1/5713 (0.0%)	0.91	7/7749 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	1708	GLU	CD-OE2	6.47	1.32	1.25

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1540	MET	CG-SD-CE	-7.69	87.90	100.20
1	A	1626	ARG	NE-CZ-NH2	-5.77	117.41	120.30
1	A	1470	CYS	N-CA-CB	5.75	120.94	110.60
1	A	1661	ARG	NE-CZ-NH2	-5.59	117.50	120.30
1	B	1544	TYR	CB-CG-CD1	5.26	124.16	121.00
1	A	1661	ARG	CG-CD-NE	-5.09	101.11	111.80
1	B	1708	GLU	CB-CG-CD	5.05	127.84	114.20

There are no chirality outliers.

There are no planarity outliers.

4.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2768	0	2650	18	0
1	B	2798	0	2685	24	0
2	A	4	0	0	0	0
2	B	1	0	0	1	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	B	16	0	0	0	0
5	A	377	0	0	8	1
5	B	327	0	0	11	1
All	All	6293	0	5335	42	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1407:GLU:HG2	5:A:2193:HOH:O	1.59	1.00
1:A:1476:GLU:CB	5:A:2169:HOH:O	2.17	0.93
1:B:1715:ARG:HG3	5:B:2111:HOH:O	1.77	0.84
1:B:1602:ARG:O	5:B:1901:HOH:O	1.96	0.83
1:B:1516:LYS:HE3	5:B:1943:HOH:O	1.79	0.83
1:B:1517[B]:ARG:HG3	1:B:1517[B]:ARG:HH11	1.43	0.83
1:B:1644:HIS:NE2	1:B:1649:ASP:OD2	2.16	0.79
1:B:1518:ASP:HB3	5:B:2153:HOH:O	1.82	0.79
1:B:1517[B]:ARG:HH11	1:B:1517[B]:ARG:CG	1.97	0.77
1:A:1406[A]:ARG:NH1	5:A:1903:HOH:O	2.20	0.74
1:A:1495:ARG:NH2	5:A:1904:HOH:O	2.21	0.74
1:B:1650:GLN:OE1	5:B:1902:HOH:O	2.05	0.74
1:B:1517[B]:ARG:HG3	1:B:1517[B]:ARG:NH1	2.03	0.71
1:A:1582:HIS:CE1	5:A:1956:HOH:O	2.48	0.66
1:A:1639:GLU:O	5:A:1902:HOH:O	2.13	0.65
1:B:1552:ARG:NH1	1:B:1592:GLU:OE1	2.31	0.62
1:A:1407:GLU:O	1:A:1410:LYS:HG2	2.01	0.61
1:A:1406[A]:ARG:NH2	5:A:1907:HOH:O	2.34	0.61
1:A:1385:LEU:HD12	1:A:1392[B]:CYS:SG	2.41	0.60
1:A:1502:PHE:CE2	1:A:1506:MET:HE2	2.37	0.58
1:B:1708:GLU:HG3	5:B:2209:HOH:O	2.05	0.56
1:B:1505:LEU:HD22	1:B:1700:VAL:HG11	1.87	0.56
1:B:1715:ARG:HD3	5:B:2190:HOH:O	2.06	0.55
1:A:1385:LEU:HG	1:A:1392[B]:CYS:SG	2.49	0.53
1:B:1494:PHE:CZ	1:B:1498:MET:HE3	2.46	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1388:GLY:N	5:B:1908:HOH:O	2.38	0.51
1:A:1385:LEU:CD1	1:A:1392[B]:CYS:SG	2.99	0.50
1:A:1582:HIS:CG	1:A:1582:HIS:O	2.65	0.50
1:A:1624:LYS:NZ	5:A:1901:HOH:O	1.87	0.49
1:B:1587:LEU:HD12	1:B:1587:LEU:O	2.15	0.46
1:B:1708:GLU:CG	5:B:2209:HOH:O	2.61	0.46
1:B:1517[B]:ARG:HH11	1:B:1517[B]:ARG:CB	2.28	0.46
1:B:1518:ASP:OD1	5:B:1904:HOH:O	2.21	0.45
1:A:1631:LYS:HE2	1:A:1666:TYR:OH	2.16	0.45
1:B:1516:LYS:CE	5:B:1943:HOH:O	2.53	0.45
1:B:1500:THR:OG1	2:B:1802:CL:CL	2.63	0.44
1:A:1661:ARG:HG3	1:A:1665:GLU:OE2	2.17	0.44
1:B:1602:ARG:HD3	1:B:1676:PHE:CE1	2.53	0.43
1:A:1644:HIS:NE2	1:A:1649:ASP:OD2	2.49	0.43
1:A:1488:TRP:HA	1:A:1489:PRO:C	2.39	0.42
1:B:1468:ILE:HD13	1:B:1468:ILE:N	2.34	0.42
1:B:1506:MET:CE	1:B:1538:PRO:HG3	2.49	0.42

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:2192:HOH:O	5:B:1937:HOH:O[2_557]	2.19	0.01

4.3 Torsion angles [i](#)

4.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	343/372 (92%)	335 (98%)	7 (2%)	1 (0%)	41 21
1	B	342/372 (92%)	330 (96%)	12 (4%)	0	100 100
All	All	685/744 (92%)	665 (97%)	19 (3%)	1 (0%)	51 28

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1581	ALA

4.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	299/331 (90%)	292 (98%)	7 (2%)	50	23
1	B	302/331 (91%)	294 (97%)	8 (3%)	46	19
All	All	601/662 (91%)	586 (98%)	15 (2%)	49	20

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1406[A]	ARG
1	A	1406[B]	ARG
1	A	1407	GLU
1	A	1579	GLU
1	A	1603	ILE
1	A	1644	HIS
1	A	1717	THR
1	B	1577	ILE
1	B	1582	HIS
1	B	1587	LEU
1	B	1588	LYS
1	B	1605	ASP
1	B	1607	LYS
1	B	1643	ASP
1	B	1724	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1650	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	1650	GLN
1	B	1656	GLN

4.3.3 RNA [i](#)

There are no RNA molecules in this entry.

4.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

4.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 1 could not be matched to an existing wwPDB Chemical Component Dictionary definition at this stage and 7 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.

The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



4.7 Other polymers [i](#)

There are no such residues in this entry.

4.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

5 Fit of model and data

5.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	340/372 (91%)	0.12	20 (5%) 22 22	14, 24, 58, 127	5 (1%)
1	B	344/372 (92%)	0.08	24 (6%) 16 16	16, 28, 57, 123	15 (4%)
All	All	684/744 (91%)	0.10	44 (6%) 19 19	14, 26, 58, 127	20 (2%)

All (44) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1581	ALA	15.5
1	A	1577	ILE	12.8
1	A	1578	GLY	11.2
1	B	1581	ALA	10.4
1	A	1392[A]	CYS	10.2
1	B	1603	ILE	8.3
1	A	1580	GLY	8.2
1	A	1379[A]	MET	7.6
1	A	1582	HIS	7.1
1	B	1577	ILE	6.9
1	B	1587	LEU	6.1
1	B	1606	HIS	6.0
1	B	1598	VAL	5.9
1	A	1564[A]	SER	5.9
1	A	1378	SER	5.4
1	B	1517[A]	ARG	5.3
1	A	1588[A]	LYS	5.3
1	A	1406[A]	ARG	5.2
1	A	1717	THR	5.1
1	B	1605	ASP	4.9
1	B	1599	THR	4.7
1	A	1606	HIS	4.6
1	A	1579	GLU	4.6
1	B	1604	HIS	4.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	1583	ASP	3.9
1	B	1582	HIS	3.8
1	B	1596	ASP	3.6
1	B	1600	LYS	3.6
1	B	1664[A]	GLU	3.5
1	B	1597	GLU	3.5
1	B	1601	GLU	3.4
1	A	1642	PRO	3.3
1	B	1602	ARG	3.1
1	B	1590	ILE	3.1
1	B	1607	LYS	3.0
1	A	1603	ILE	2.9
1	B	1642	PRO	2.6
1	A	1470	CYS	2.6
1	A	1576	PRO	2.4
1	A	1639	GLU	2.2
1	B	1584	GLU	2.1
1	B	1387	ASP	2.1
1	B	1592	GLU	2.0
1	B	1583	ASP	2.0

5.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.4 Ligands [i](#)

LIGAND-RSR INFOmissingINFO

5.5 Other polymers [i](#)

There are no such residues in this entry.