



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 29, 2024 – 09:24 AM EDT

PDB ID : 3RC3
Title : Human Mitochondrial Helicase Suv3
Authors : Dauter, Z.; Jedrzejczak, R.; Dauter, M.; Szczesny, R.; Stepień, P.
Deposited on : 2011-03-30
Resolution : 2.08 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.08 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	7574 (2.10-2.06)
Clashscore	180529	8325 (2.10-2.06)
Ramachandran outliers	177936	8271 (2.10-2.06)
Sidechain outliers	177891	8272 (2.10-2.06)
RSRZ outliers	164620	7574 (2.10-2.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	677	 76% 12% • 10%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 5405 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called ATP-dependent RNA helicase SUPV3L1, mitochondrial.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	607	4857	3120	827	882	14	14	0	1	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	46	GLY	-	expression tag	UNP Q8IYB8

- Molecule 2 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (three-letter code: ANP) (formula: $C_{10}H_{17}N_6O_{12}P_3$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	31	10	6	12	3	0	0

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Na 1 1	0	0

- Molecule 4 is AZIDE ION (three-letter code: AZI) (formula: N₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total N 3 3	0	0

- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total Cl 1 1	0	0


- Molecule 6 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	512	Total O 512 512	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: ATP-dependent RNA helicase SUPV3L1, mitochondrial

Chain A:  76% 12% 10%



4 Data and refinement statistics i

Property	Value	Source
Space group	P 32	Depositor
Cell constants a, b, c, α , β , γ	89.39Å 89.39Å 88.07Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	39.84 – 2.08 39.84 – 2.08	Depositor EDS
% Data completeness (in resolution range)	100.0 (39.84-2.08) 99.8 (39.84-2.08)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.27 (at 2.08Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.160 , 0.203 0.161 , 0.203	Depositor DCC
R_{free} test set	965 reflections (2.05%)	wwPDB-VP
Wilson B-factor (Å ²)	32.5	Xtrriage
Anisotropy	0.020	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 56.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.011 for -h,-k,l 0.028 for h,-h-k,-l 0.015 for -k,-h,-l	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	5405	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.12% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: CL, NA, AZI, ANP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.14	7/4962 (0.1%)	0.99	11/6699 (0.2%)

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	663	ASP	CB-CG	7.11	1.66	1.51
1	A	287	VAL	CB-CG2	6.10	1.65	1.52
1	A	333	MSE	SE-CE	-6.02	1.59	1.95
1	A	281	VAL	CB-CG2	5.71	1.64	1.52
1	A	272	VAL	CB-CG2	5.64	1.64	1.52
1	A	663	ASP	CG-OD2	5.20	1.37	1.25
1	A	396	TYR	CD2-CE2	5.02	1.46	1.39

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	347	ARG	NE-CZ-NH2	-11.02	114.79	120.30
1	A	648	MSE	CG-SE-CE	10.34	121.65	98.90
1	A	301	ARG	NE-CZ-NH1	8.19	124.39	120.30
1	A	347	ARG	NE-CZ-NH1	7.76	124.18	120.30
1	A	147	ASP	CB-CG-OD1	7.32	124.89	118.30
1	A	301	ARG	NE-CZ-NH2	-6.46	117.07	120.30
1	A	568	LEU	CA-CB-CG	-6.10	101.27	115.30
1	A	625	LEU	CB-CG-CD2	6.10	121.36	111.00
1	A	595	ARG	NE-CZ-NH1	-5.17	117.71	120.30
1	A	603	LEU	CA-CB-CG	5.17	127.18	115.30
1	A	200	ARG	NE-CZ-NH2	-5.09	117.75	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4857	0	4889	57	0
2	A	31	0	13	1	0
3	A	1	0	0	0	0
4	A	3	0	0	0	0
5	A	1	0	0	0	0
6	A	512	0	0	10	0
All	All	5405	0	4902	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:646:MSE:HG3	6:A:1046:HOH:O	1.65	0.96
1:A:58:ILE:HG23	1:A:59:PRO:HD3	1.51	0.92
1:A:118:ARG:HD2	6:A:1134:HOH:O	1.71	0.90
1:A:608:LEU:HD23	1:A:655:ILE:HD12	1.53	0.90
1:A:70:LYS:HE2	6:A:1093:HOH:O	1.84	0.77
1:A:164:ARG:HD2	6:A:1156:HOH:O	1.86	0.76
1:A:518:GLU:HA	1:A:568:LEU:HD22	1.69	0.74
1:A:572:TYR:O	1:A:576:THR:HG23	1.86	0.74
1:A:366:PRO:HB3	1:A:419:LYS:HG2	1.71	0.72
1:A:238:HIS:HD2	6:A:1027:HOH:O	1.74	0.71
1:A:608:LEU:HD23	1:A:655:ILE:CD1	2.22	0.69
1:A:358:LEU:HB3	1:A:486:MSE:HE3	1.79	0.64
1:A:686:LEU:HB2	6:A:969:HOH:O	2.00	0.61
1:A:489:GLU:H	1:A:489:GLU:CD	2.09	0.56
1:A:602:PRO:HG3	1:A:648:MSE:HE2	1.85	0.56
1:A:347:ARG:NH2	1:A:469:GLY:O	2.39	0.56
1:A:521:ALA:HB2	1:A:568:LEU:HD13	1.89	0.55
1:A:333:MSE:HE2	1:A:333:MSE:N	2.22	0.54
1:A:238:HIS:HE1	1:A:258:GLU:OE1	1.92	0.53
1:A:596:GLN:NE2	1:A:601:GLU:HG2	2.23	0.53
1:A:146:ASN:O	1:A:150:PHE:HD1	1.92	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:285:TYR:O	1:A:314:ALA:HA	2.09	0.52
1:A:87:PRO:HD2	6:A:1056:HOH:O	2.08	0.52
1:A:185:LEU:CD2	1:A:516:GLN:HE22	2.24	0.51
1:A:187:ILE:N	1:A:188:PRO:HD3	2.27	0.50
1:A:518:GLU:CA	1:A:568:LEU:HD22	2.39	0.49
1:A:332:LEU:HD23	1:A:545:GLN:HG3	1.95	0.49
2:A:801:ANP:PB	2:A:801:ANP:H5'2	2.52	0.49
1:A:185:LEU:HD21	1:A:516:GLN:HE22	1.78	0.49
1:A:236:LEU:CD1	1:A:429:MSE:HE2	2.43	0.49
1:A:129:ASN:CG	1:A:133:GLN:NE2	2.67	0.48
1:A:58:ILE:CG2	1:A:59:PRO:HD3	2.34	0.48
1:A:642:SER:HA	1:A:649:PHE:HB2	1.94	0.48
1:A:202:ILE:HG12	1:A:333:MSE:SE	2.65	0.47
1:A:445:LYS:C	6:A:1178:HOH:O	2.53	0.47
1:A:129:ASN:OD1	1:A:133:GLN:NE2	2.48	0.47
1:A:521:ALA:CB	1:A:568:LEU:HD13	2.45	0.47
1:A:366:PRO:HB3	1:A:419:LYS:CG	2.43	0.47
1:A:227:GLY:HA2	1:A:287:VAL:O	2.16	0.46
1:A:265:ASN:OD1	1:A:267:LYS:HB2	2.17	0.45
1:A:104:LYS:HE2	1:A:105:GLU:OE2	2.17	0.44
1:A:215:TYR:O	1:A:219:GLN:HG2	2.18	0.44
1:A:383:ARG:O	1:A:387:ILE:HG23	2.18	0.43
1:A:623:LYS:O	1:A:623:LYS:HD3	2.19	0.43
1:A:257:GLU:C	1:A:258:GLU:HG3	2.38	0.43
1:A:376:ASN:HB3	1:A:688:LEU:O	2.18	0.43
1:A:239:GLU:HG3	6:A:1113:HOH:O	2.18	0.43
1:A:129:ASN:ND2	1:A:133:GLN:NE2	2.68	0.42
1:A:228:VAL:HG23	1:A:272:VAL:HG13	2.02	0.42
1:A:357:ALA:HB2	1:A:488:HIS:HA	2.03	0.41
1:A:608:LEU:HD22	1:A:638:TYR:CD2	2.56	0.41
1:A:614:TRP:CZ2	1:A:661:GLU:HG2	2.56	0.41
1:A:569:ARG:HD2	1:A:569:ARG:HA	1.89	0.41
1:A:654:LEU:HD23	6:A:1001:HOH:O	2.20	0.40
1:A:288:ALA:HB1	1:A:312:LEU:HD13	2.03	0.40
1:A:373:PHE:CZ	1:A:459:ILE:HG21	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	602/677 (89%)	595 (99%)	7 (1%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	539/575 (94%)	519 (96%)	20 (4%)	29 30

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	108	LYS
1	A	136	SER
1	A	168	GLN
1	A	220	LYS
1	A	234	LYS
1	A	259	ARG
1	A	297	ARG
1	A	333	MSE
1	A	387	ILE
1	A	401	PRO
1	A	419	LYS
1	A	429	MSE
1	A	486	MSE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	489	GLU
1	A	511	HIS
1	A	520	PHE
1	A	545	GLN
1	A	568	LEU
1	A	625	LEU
1	A	648	MSE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	133	GLN
1	A	238	HIS
1	A	268	GLN
1	A	415	ASN
1	A	511	HIS
1	A	596	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	ANP	A	801	-	29,33,33	2.30	9 (31%)	31,52,52	2.25	10 (32%)
4	AZI	A	902	-	2,2,2	0.37	0	0,1,1	-	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ANP	A	801	-	-	4/14/38/38	0/3/3/3

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	801	ANP	PG-O1G	5.09	1.53	1.46
2	A	801	ANP	PG-N3B	5.01	1.76	1.63
2	A	801	ANP	PB-N3B	4.83	1.76	1.63
2	A	801	ANP	PB-O1B	4.38	1.52	1.46
2	A	801	ANP	O4'-C1'	3.49	1.45	1.40
2	A	801	ANP	PB-O3A	3.48	1.63	1.59
2	A	801	ANP	C2-N1	2.80	1.38	1.33
2	A	801	ANP	C2-N3	2.71	1.36	1.32
2	A	801	ANP	PA-O5'	2.42	1.68	1.59

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	801	ANP	O1G-PG-N3B	-5.55	103.60	111.77
2	A	801	ANP	N3-C2-N1	-5.52	121.19	128.67
2	A	801	ANP	O2B-PB-O1B	4.90	120.37	109.87
2	A	801	ANP	O2B-PB-O3A	4.55	119.81	104.64
2	A	801	ANP	O1B-PB-N3B	-2.71	107.77	111.77
2	A	801	ANP	N6-C6-N1	2.71	124.12	118.33
2	A	801	ANP	O2A-PA-O1A	2.37	123.47	112.44
2	A	801	ANP	O2G-PG-O3G	2.31	113.80	107.59
2	A	801	ANP	O2A-PA-O3A	2.19	113.19	107.27
2	A	801	ANP	O3A-PB-N3B	-2.07	100.84	106.59

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	801	ANP	PG-N3B-PB-O1B
2	A	801	ANP	PG-N3B-PB-O3A
2	A	801	ANP	PA-O3A-PB-O2B
2	A	801	ANP	PB-O3A-PA-O1A

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	801	ANP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	593/677 (87%)	-0.48	1 (0%) 92 92	19, 35, 57, 79	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	58	ILE	2.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	AZI	A	902	3/3	0.83	0.18	34,34,42,50	0
2	ANP	A	801	31/31	0.96	0.07	26,37,47,51	0
3	NA	A	901	1/1	0.98	0.04	37,37,37,37	0
5	CL	A	903	1/1	0.98	0.06	32,32,32,32	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers

as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.