



# Full wwPDB X-ray Structure Validation Report ⓘ

May 15, 2020 – 08:42 pm BST

PDB ID : 1RDA  
Title : CRYSTAL STRUCTURES OF RIBONUCLEASE HI ACTIVE SITE MUTANTS FROM ESCHERICHIA COLI  
Authors : Katayanagi, K.; Morikawa, K.  
Deposited on : 1993-06-23  
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtrriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	1585 (2.16-2.16)
Ramachandran outliers	138981	1560 (2.16-2.16)
Sidechain outliers	138945	1559 (2.16-2.16)
RSRZ outliers	127900	1456 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	155	

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1338 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RIBONUCLEASE H.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	155	1238	776	228	227	7	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	10	ASN	ASP	CONFLICT	UNP P0A7Y4


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
2	A	100	100	100	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: RIBONUCLEASE H



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	44.17Å 87.23Å 35.27Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 2.15 9.93 – 2.15	Depositor EDS
% Data completeness (in resolution range)	(Not available) (10.00-2.15) 80.5 (9.93-2.15)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$	-	Xtrriage
Refinement program	PROLSQ	Depositor
R, $R_{free}$	0.189 , (Not available) 0.180 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	17.9	Xtrriage
Anisotropy	0.515	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.25 , 81.5	EDS
L-test for twinning <sup>1</sup>	$\langle  L  \rangle = 0.41$ , $\langle L^2 \rangle = 0.24$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	1338	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	17.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.39% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.95	0/1266	1.77	29/1710 (1.7%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	106	ARG	NE-CZ-NH2	-14.97	112.82	120.30
1	A	106	ARG	NE-CZ-NH1	12.34	126.47	120.30
1	A	28	TYR	CA-CB-CG	9.27	131.02	113.40
1	A	27	ARG	NE-CZ-NH2	-8.06	116.27	120.30
1	A	41	ARG	NE-CZ-NH2	-7.78	116.41	120.30
1	A	138	ARG	NE-CZ-NH1	7.24	123.92	120.30
1	A	149	THR	CA-CB-CG2	7.02	122.23	112.40
1	A	75	ARG	NE-CZ-NH2	-6.94	116.83	120.30
1	A	132	ARG	NE-CZ-NH2	-6.79	116.91	120.30
1	A	64	GLU	CA-CB-CG	6.62	127.97	113.40
1	A	153	VAL	C-N-CA	6.56	138.11	121.70
1	A	70	ASP	CB-CG-OD1	6.19	123.87	118.30
1	A	129	GLU	OE1-CD-OE2	-6.10	115.98	123.30
1	A	66	ILE	O-C-N	6.01	132.32	122.70
1	A	67	LEU	CB-CA-C	6.00	121.61	110.20
1	A	88	ARG	NE-CZ-NH2	-5.79	117.41	120.30
1	A	46	ARG	NE-CZ-NH1	-5.71	117.44	120.30
1	A	154	GLU	CG-CD-OE1	5.62	129.55	118.30

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	14	LEU	CB-CA-C	5.39	120.45	110.20
1	A	24	ALA	C-N-CA	5.33	135.03	121.70
1	A	155	VAL	CA-CB-CG1	5.32	118.88	110.90
1	A	131	GLU	CG-CD-OE1	5.26	128.81	118.30
1	A	127	HIS	N-CA-CB	5.22	120.00	110.60
1	A	28	TYR	CB-CG-CD2	5.17	124.10	121.00
1	A	154	GLU	CA-CB-CG	5.15	124.72	113.40
1	A	127	HIS	O-C-N	5.12	130.83	121.10
1	A	66	ILE	N-CA-CB	5.08	122.49	110.80
1	A	116	ILE	N-CA-C	5.05	124.63	111.00
1	A	132	ARG	CD-NE-CZ	-5.01	116.58	123.60

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	132	ARG	Sidechain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1238	0	1219	37	1
2	A	100	0	0	8	0
All	All	1338	0	1219	37	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:63:CYS:H	1:A:114:HIS:HD2	1.17	0.89
1:A:111:LEU:HD23	1:A:116:ILE:HD12	1.69	0.74
1:A:67:LEU:HD22	1:A:118:TRP:CZ3	2.25	0.71

Continued on next page...

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:84:ASN:HB2	2:A:190:HOH:O	1.90	0.69
1:A:77:GLY:HA3	1:A:104:TRP:CH2	2.29	0.68
1:A:114:HIS:HE1	2:A:236:HOH:O	1.76	0.67
1:A:29:ARG:HB3	2:A:223:HOH:O	1.99	0.62
1:A:63:CYS:H	1:A:114:HIS:CD2	2.09	0.61
1:A:127:HIS:HB3	1:A:128:PRO:HD2	1.84	0.59
1:A:111:LEU:CD2	1:A:116:ILE:HD12	2.32	0.59
1:A:44:ASN:O	1:A:48:GLU:HG3	2.05	0.57
1:A:49:LEU:HD23	1:A:74:VAL:HG22	1.87	0.56
1:A:28:TYR:HD1	1:A:33:LYS:HZ1	1.53	0.56
1:A:30:GLY:N	2:A:223:HOH:O	2.40	0.54
1:A:154:GLU:O	1:A:155:VAL:HG23	2.09	0.53
1:A:72:GLN:HB2	2:A:254:HOH:O	2.09	0.53
1:A:149:THR:O	1:A:151:TYR:N	2.42	0.52
1:A:75:ARG:HA	1:A:120:TRP:CH2	2.47	0.50
1:A:8:PHE:CD1	1:A:129:GLU:HB2	2.48	0.48
1:A:153:VAL:HB	1:A:154:GLU:OE2	2.15	0.47
1:A:49:LEU:HD23	1:A:74:VAL:CG2	2.44	0.47
1:A:28:TYR:O	1:A:29:ARG:HB2	2.14	0.47
1:A:9:THR:HB	1:A:51:ALA:HB1	1.97	0.46
1:A:9:THR:HB	1:A:51:ALA:CB	2.46	0.46
1:A:6:GLU:O	1:A:26:LEU:HA	2.16	0.44
1:A:155:VAL:HG11	2:A:242:HOH:O	2.17	0.44
1:A:67:LEU:HD22	1:A:118:TRP:HZ3	1.76	0.44
1:A:81:TRP:HB3	1:A:85:TRP:CZ2	2.53	0.43
1:A:139:ALA:O	1:A:142:MET:HB2	2.19	0.43
1:A:63:CYS:HB3	2:A:215:HOH:O	2.18	0.43
1:A:48:GLU:OE1	1:A:71:SER:OG	2.36	0.43
1:A:44:ASN:HB3	2:A:251:HOH:O	2.20	0.41
1:A:6:GLU:HG2	1:A:66:ILE:CG2	2.51	0.41
1:A:39:TYR:HA	1:A:146:LEU:O	2.21	0.41
1:A:142:MET:HE2	1:A:142:MET:HB3	1.86	0.41
1:A:102:ASP:OD2	1:A:150:GLY:HA3	2.22	0.40
1:A:75:ARG:HG3	1:A:120:TRP:CE2	2.55	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:94:ASP:O	1:A:132:ARG:CD[3_555]	2.17	0.03



## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	153/155 (99%)	140 (92%)	9 (6%)	4 (3%)	<b>5</b> <b>1</b>

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	154	GLU
1	A	150	GLY
1	A	123	GLY
1	A	29	ARG

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	127/127 (100%)	113 (89%)	14 (11%)	<b>6</b> <b>3</b>

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	4	GLN
1	A	12	SER
1	A	26	LEU
1	A	31	ARG
1	A	44	ASN
1	A	60	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	63	CYS
1	A	64	GLU
1	A	72	GLN
1	A	116	ILE
1	A	135	GLU
1	A	149	THR
1	A	154	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	44	ASN
1	A	45	ASN
1	A	72	GLN
1	A	84	ASN
1	A	105	GLN
1	A	114	HIS
1	A	152	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	155/155 (100%)	-0.39	0 <b>100</b> <b>100</b>	6, 15, 27, 36	0

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.