



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 14, 2018 – 02:19 pm GMT

PDB ID : 1RFD  
Title : ANTI-COCAINE ANTIBODY M82G2  
Authors : Pozharski, E.; Hewagama, A.; Shanafelt, A.B.; Petsko, G.A.; Ringe, D.  
Deposited on : 2003-11-07  
Resolution : 2.09 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : trunk31020  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk31020


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.09 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	4608 (2.10-2.10)
Clashscore	122126	5109 (2.10-2.10)
Ramachandran outliers	120053	5059 (2.10-2.10)
Sidechain outliers	120020	5060 (2.10-2.10)
RSRZ outliers	108989	4497 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	L	218	<p>4% 80% 19% •</p>
2	H	223	<p>3% 81% 17% ••</p>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3576 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fab M82G2, Light Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	L	218	1680	1050	282	341	7	0	0	0

- Molecule 2 is a protein called Fab M82G2, Heavy Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	H	220	1659	1042	279	327	11	0	4	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	L	102	Total	O	0	0
			102	102		
3	H	135	Total	O	0	0
			135	135		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Fab M82G2, Light Chain



- Molecule 2: Fab M82G2, Heavy Chain



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	74.92Å 74.92Å 151.27Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.00 – 2.09 27.42 – 2.09	Depositor EDS
% Data completeness (in resolution range)	90.2 (30.00-2.09) 90.6 (27.42-2.09)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.66 (at 2.10Å)	Xtrriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.183 , 0.226 0.193 , 0.237	Depositor DCC
$R_{free}$ test set	1311 reflections (4.87%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	29.0	Xtrriage
Anisotropy	0.002	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 51.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.037 for -h,-k,l	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3576	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	32.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.30% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	L	0.55	0/1722	0.76	0/2341
2	H	0.56	0/1700	0.78	0/2319
All	All	0.55	0/3422	0.77	0/4660

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	1680	0	1600	31	0
2	H	1659	0	1606	44	0
3	H	135	0	0	5	0
3	L	102	0	0	0	0
All	All	3576	0	3206	72	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (72) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:5:GLN:HA	2:H:105:GLN:HE22	1.25	0.99

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:131:ALA:HA	2:H:135:GLY:HA2	1.60	0.82
2:H:34:VAL:HG11	2:H:78:VAL:HG21	1.64	0.78
1:L:182:THR:OG1	1:L:185:GLU:HG3	1.85	0.77
2:H:5:GLN:HA	2:H:105:GLN:NE2	2.00	0.76
2:H:105:GLN:CD	2:H:105:GLN:H	1.91	0.74
1:L:33:LEU:HD22	1:L:71:PHE:CG	2.29	0.68
2:H:58:LYS:HG2	3:H:289:HOH:O	1.94	0.67
2:H:131:ALA:HA	2:H:135:GLY:CA	2.24	0.67
2:H:179:GLN:HG3	2:H:180:SER:H	1.59	0.66
2:H:179:GLN:HG3	2:H:180:SER:N	2.12	0.65
2:H:64:LYS:HE2	3:H:267:HOH:O	1.97	0.65
2:H:131:ALA:CA	2:H:135:GLY:HA2	2.26	0.63
1:L:89:MET:HE3	2:H:100(B):PHE:CE2	2.33	0.63
1:L:13:VAL:HG11	1:L:19:VAL:HG21	1.83	0.60
2:H:2:VAL:HG11	2:H:102:TYR:CD1	2.38	0.59
2:H:130:GLY:O	2:H:131:ALA:HB2	2.03	0.58
2:H:99:GLY:O	2:H:100:ARG:HG2	2.02	0.58
2:H:97:GLN:C	2:H:99:GLY:H	2.07	0.58
1:L:187:GLU:HA	1:L:211:ARG:NE	2.19	0.58
2:H:226:VAL:HG22	3:H:316:HOH:O	2.04	0.57
1:L:125:LEU:O	1:L:183:LYS:HD2	2.04	0.56
1:L:89:MET:CE	2:H:100(B):PHE:CE2	2.89	0.55
1:L:61:ARG:HD2	1:L:77:ARG:O	2.07	0.54
2:H:22[A]:CYS:HB3	2:H:78:VAL:HG23	1.90	0.54
1:L:136:LEU:HD21	1:L:196:ALA:HB2	1.90	0.53
1:L:212:GLY:O	1:L:213:GLU:HG3	2.10	0.52
1:L:33:LEU:HD22	1:L:71:PHE:CD2	2.45	0.52
2:H:59:TYR:HB2	2:H:64:LYS:HG2	1.92	0.51
2:H:2:VAL:HG11	2:H:102:TYR:CG	2.46	0.51
2:H:52:ARG:HD2	2:H:56:ALA:HB3	1.92	0.51
1:L:149:LYS:HB2	1:L:193:THR:HB	1.92	0.51
2:H:105:GLN:N	2:H:105:GLN:CD	2.62	0.51
1:L:182:THR:HG23	1:L:185:GLU:OE2	2.12	0.50
1:L:5:THR:HB	1:L:24:ARG:HG2	1.93	0.49
1:L:151:ASP:OD2	1:L:189:HIS:HB3	2.13	0.49
1:L:182:THR:HG1	1:L:185:GLU:HG3	1.78	0.49
1:L:89:MET:HE3	2:H:100(B):PHE:HE2	1.76	0.47
2:H:119:PRO:HB3	2:H:147:TYR:HB3	1.96	0.47
1:L:104:LEU:HD21	1:L:106:ILE:HD11	1.96	0.47
2:H:95:VAL:HG22	2:H:100(B):PHE:HD1	1.79	0.46
2:H:179:GLN:CG	2:H:180:SER:N	2.75	0.45

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:22[B]:CYS:HB3	2:H:78:VAL:HG23	1.98	0.45
1:L:37:LEU:HD13	1:L:86:TYR:CZ	2.51	0.45
2:H:97:GLN:C	2:H:99:GLY:N	2.68	0.45
2:H:130:GLY:O	2:H:131:ALA:CB	2.65	0.44
2:H:180:SER:O	2:H:181:ASP:HB2	2.18	0.43
2:H:114:ALA:HB3	2:H:148:PHE:CE2	2.53	0.43
2:H:12:VAL:HG11	2:H:82(C):LEU:HD13	2.00	0.43
2:H:34:VAL:HB	2:H:78:VAL:HG11	2.00	0.43
1:L:19:VAL:HG12	1:L:75:PHE:HB2	2.01	0.43
1:L:80:ALA:O	1:L:83:VAL:HG13	2.19	0.42
2:H:98:LEU:HD23	2:H:98:LEU:HA	1.76	0.42
2:H:100:ARG:HG3	2:H:100:ARG:HH11	1.83	0.42
2:H:100:ARG:NH2	2:H:102:TYR:OH	2.52	0.42
2:H:96:PRO:HB3	3:H:279:HOH:O	2.19	0.42
1:L:10:SER:HA	1:L:103:LYS:O	2.19	0.42
1:L:13:VAL:CG1	1:L:19:VAL:HG21	2.49	0.42
2:H:14:PRO:HG3	2:H:111:VAL:HG12	2.01	0.42
1:L:136:LEU:N	1:L:136:LEU:HD12	2.34	0.41
2:H:179:GLN:HB3	3:H:266:HOH:O	2.20	0.41
1:L:122:SER:O	1:L:126:THR:HG23	2.21	0.41
2:H:57:THR:O	2:H:58:LYS:HD2	2.20	0.41
2:H:35:ASP:OD1	2:H:50:GLU:HB2	2.21	0.41
2:H:34:VAL:CG1	2:H:78:VAL:HG21	2.44	0.41
1:L:150:ILE:O	1:L:151:ASP:HB2	2.20	0.41
1:L:118:PHE:HA	1:L:119:PRO:HD3	1.96	0.40
2:H:71:ARG:HB3	2:H:78:VAL:HG12	2.04	0.40
1:L:136:LEU:CD2	1:L:196:ALA:HB2	2.51	0.40
1:L:199:LYS:HB2	1:L:199:LYS:HE2	1.89	0.40
1:L:187:GLU:O	1:L:211:ARG:NH2	2.54	0.40
1:L:142:GLN:HB3	1:L:173:TYR:CE2	2.57	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	216/218 (99%)	212 (98%)	3 (1%)	1 (0%)	31	27
2	H	220/223 (99%)	215 (98%)	4 (2%)	1 (0%)	31	27
All	All	436/441 (99%)	427 (98%)	7 (2%)	2 (0%)	31	27

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	H	99	GLY
1	L	68	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	192/192 (100%)	188 (98%)	4 (2%)	56	62
2	H	186/185 (100%)	182 (98%)	4 (2%)	55	60
All	All	378/377 (100%)	370 (98%)	8 (2%)	56	62

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	122	SER
1	L	157	SER
1	L	199	LYS
1	L	202	THR
2	H	105	GLN
2	H	149	PRO
2	H	180	SER
2	H	209	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	L	137	ASN
2	H	54	ASN
2	H	82(A)	ASN
2	H	105	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	L	218/218 (100%)	-0.23	8 (3%) 41 48	19, 30, 59, 88	0
2	H	220/223 (98%)	-0.31	6 (2%) 54 61	19, 27, 54, 82	0
All	All	438/441 (99%)	-0.27	14 (3%) 47 55	19, 29, 56, 88	0

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	H	131	ALA	4.8
2	H	1	GLU	4.0
1	L	212	GLY	3.9
1	L	7	ALA	3.7
2	H	128	SER	3.3
1	L	202	THR	3.2
2	H	99	GLY	2.9
1	L	213	GLU	2.9
1	L	188	ARG	2.8
1	L	150	ILE	2.2
2	H	129	GLY	2.2
2	H	100(B)	PHE	2.2
1	L	183	LYS	2.1
1	L	203	SER	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands

There are no ligands in this entry.

## 6.5 Other polymers

There are no such residues in this entry.