



# Full wwPDB X-ray Structure Validation Report ⓘ

May 27, 2020 – 10:11 pm BST

PDB ID : 2RFN  
Title : x-ray structure of c-Met with inhibitor.  
Authors : Bellon, S.F.; Kaplan-Lefko, P.; Yang, Y.; Zhang, Y.; Moriguchi, J.; Dussault, I.  
Deposited on : 2007-10-01  
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	310	<div style="display: flex; align-items: center;"> <div style="width: 3%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 68%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 17%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 14%; height: 10px; background-color: grey;"></div> </div> <p style="margin-left: 30px;">3%      68%      17%      •      14%</p>
1	B	310	<div style="display: flex; align-items: center;"> <div style="width: 4%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 71%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 13%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 13%; height: 10px; background-color: grey;"></div> </div> <p style="margin-left: 30px;">4%      71%      13%      •      13%</p>

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 4490 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Hepatocyte growth factor receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	268	2114	1370	355	375	14	0	0	0
1	B	269	2119	1373	357	375	14	0	0	0

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1047	MET	-	INITIATING METHIONINE	UNP P08581
A	1272	LEU	VAL	CONFLICT	UNP P08581
A	1352	HIS	-	EXPRESSION TAG	UNP P08581
A	1353	HIS	-	EXPRESSION TAG	UNP P08581
A	1354	HIS	-	EXPRESSION TAG	UNP P08581
A	1355	HIS	-	EXPRESSION TAG	UNP P08581
A	1356	HIS	-	EXPRESSION TAG	UNP P08581
B	1047	MET	-	INITIATING METHIONINE	UNP P08581
B	1272	LEU	VAL	CONFLICT	UNP P08581
B	1352	HIS	-	EXPRESSION TAG	UNP P08581
B	1353	HIS	-	EXPRESSION TAG	UNP P08581
B	1354	HIS	-	EXPRESSION TAG	UNP P08581
B	1355	HIS	-	EXPRESSION TAG	UNP P08581
B	1356	HIS	-	EXPRESSION TAG	UNP P08581

- Molecule 2 is 2-benzyl-5-(3-fluoro-4-{{6-methoxy-7-(3-morpholin-4-ylpropoxy)quinolin-4-yl}oxy}phenyl)-3-methylpyrimidin-4(3H)-one (three-letter code: AM7) (formula: C<sub>35</sub>H<sub>35</sub>FN<sub>4</sub>O<sub>5</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	F	N	O		
2	A	1	45	35	1	4	5	0	0
2	B	1	45	35	1	4	5	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	81	81	81	0	0
3	B	86	86	86	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Hepatocyte growth factor receptor



- Molecule 1: Hepatocyte growth factor receptor



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	74.51Å 78.92Å 129.13Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.50 19.68 – 2.38	Depositor EDS
% Data completeness (in resolution range)	97.6 (20.00-2.50) 96.5 (19.68-2.38)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.82 (at 2.38Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.238 , 0.294 0.231 , 0.282	Depositor DCC
$R_{free}$ test set	1512 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	46.9	Xtrriage
Anisotropy	0.570	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.28 , 45.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	4490	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	51.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 24.43 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.8387e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: AM7

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.44	0/2166	0.61	0/2938
1	B	0.44	0/2172	0.59	0/2947
All	All	0.44	0/4338	0.60	0/5885

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	1089	PHE	Peptide

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2114	0	2122	33	0
1	B	2119	0	2121	32	0
2	A	45	0	35	1	0
2	B	45	0	35	2	0
3	A	81	0	0	1	0
3	B	86	0	0	3	0
All	All	4490	0	4313	65	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (65) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1300:ARG:HG2	1:B:1300:ARG:HH21	1.15	1.03
1:A:1160:MET:HG2	1:A:1211:MET:HB3	1.67	0.77
1:B:1062:LEU:O	1:B:1066:VAL:HG23	1.86	0.76
1:B:1300:ARG:NH2	1:B:1300:ARG:HG2	1.95	0.72
1:B:1122:SER:O	1:B:1126:THR:HG22	1.89	0.71
1:B:1066:VAL:HG22	1:B:1129:ILE:HD11	1.71	0.70
1:B:1248:LYS:HD3	1:B:1284:TYR:HB2	1.74	0.70
1:B:1300:ARG:CG	1:B:1300:ARG:HH21	2.01	0.69
1:A:1284:TYR:HB3	1:A:1287:VAL:CG1	2.25	0.66
1:B:1212:LEU:HD22	1:B:1216:PHE:HA	1.82	0.61
1:A:1284:TYR:HB3	1:A:1287:VAL:HG13	1.81	0.61
1:A:1171:ASN:OD1	1:A:1174:HIS:ND1	2.35	0.59
1:B:1059:ASN:HD22	1:B:1062:LEU:H	1.50	0.59
1:B:1284:TYR:HB3	1:B:1287:VAL:HG13	1.83	0.59
1:B:1284:TYR:HB3	1:B:1287:VAL:CG1	2.33	0.58
1:A:1140:LEU:CD1	1:A:1160:MET:HE1	2.33	0.57
1:B:1222:ASP:N	1:B:1222:ASP:OD1	2.38	0.57
1:B:1112:LEU:HD23	1:B:1155:VAL:HG23	1.87	0.56
1:A:1161:LYS:HD2	1:A:1214:GLU:HB2	1.87	0.56
1:B:1066:VAL:HG21	1:B:1125:LEU:HD13	1.87	0.55
1:B:1132:LYS:HE3	3:B:1400:HOH:O	2.05	0.55
1:A:1315:VAL:HG22	1:A:1336:ARG:HD3	1.89	0.53
1:A:1066:VAL:HG11	1:A:1125:LEU:HD22	1.91	0.52
1:A:1105:ILE:HD11	1:A:1143:LEU:HD13	1.92	0.52
1:A:1121:VAL:O	1:A:1125:LEU:HG	2.10	0.52
1:B:1131:MET:SD	2:B:1:AM7:H6	2.50	0.52
1:A:1318:LYS:HA	1:A:1321:HIS:CD2	2.45	0.51
1:B:1089:PHE:O	1:B:1110:LYS:NZ	2.34	0.51

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1059:ASN:HD22	1:A:1062:LEU:H	1.60	0.50
1:B:1058:LEU:HD12	1:B:1118:ILE:HG13	1.93	0.50
1:B:1112:LEU:CD2	1:B:1155:VAL:HG23	2.41	0.50
1:A:1066:VAL:HG22	1:A:1129:ILE:HD11	1.94	0.49
1:B:1066:VAL:HG11	1:B:1125:LEU:HD22	1.94	0.49
1:B:1059:ASN:HD21	1:B:1061:GLU:HB2	1.78	0.48
1:A:1277:MET:O	1:A:1307:TYR:HB2	2.13	0.48
1:B:1169:ILE:HD12	1:B:1279:ARG:HA	1.95	0.48
1:B:1145:ILE:HD11	2:B:1:AM7:H8	1.96	0.48
1:A:1311:PRO:O	1:A:1315:VAL:HG23	2.15	0.46
1:B:1146:CYS:HB2	1:B:1154:LEU:HB2	1.98	0.46
1:A:1282:PRO:HA	1:A:1283:PRO:HD3	1.89	0.46
1:B:1141:SER:HB3	3:B:1398:HOH:O	2.16	0.46
1:A:1298:GLN:NE2	1:A:1300:ARG:HD2	2.32	0.45
1:A:1288:ASN:HA	3:A:1430:HOH:O	2.16	0.45
1:B:1278:THR:HG23	1:B:1305:PRO:HB3	1.99	0.44
1:A:1082:GLU:HB3	1:A:1094:HIS:HD2	1.82	0.44
1:A:1310:ASP:N	1:A:1311:PRO:HD2	2.32	0.44
1:A:1248:LYS:NZ	1:A:1287:VAL:O	2.45	0.44
1:B:1126:THR:HA	1:B:1129:ILE:HD12	1.99	0.44
1:A:1222:ASP:N	1:A:1222:ASP:OD1	2.50	0.43
1:A:1059:ASN:HD21	1:A:1061:GLU:HB2	1.82	0.43
1:A:1123:GLN:HG2	3:B:1382:HOH:O	2.19	0.43
1:A:1062:LEU:O	1:A:1066:VAL:HG23	2.19	0.42
1:A:1323:LYS:HB3	1:A:1326:MET:HG3	2.01	0.42
1:B:1300:ARG:CG	1:B:1300:ARG:NH2	2.68	0.42
1:A:1325:GLU:H	1:A:1325:GLU:CD	2.23	0.42
1:A:1115:ILE:CG2	1:A:1116:THR:N	2.83	0.42
1:B:1212:LEU:CD2	1:B:1216:PHE:HA	2.49	0.42
1:A:1160:MET:HE2	1:A:1160:MET:HB2	1.69	0.41
1:B:1308:CYS:HA	1:B:1309:PRO:HD3	1.84	0.41
1:A:1069:VAL:HG12	1:A:1145:ILE:HG12	2.02	0.41
1:A:1158:PRO:O	2:A:1:AM7:H27	2.21	0.41
1:B:1140:LEU:HD21	1:B:1157:LEU:HD22	2.03	0.41
1:A:1069:VAL:O	1:A:1069:VAL:HG12	2.20	0.40
1:B:1138:ASN:O	1:B:1220:VAL:HG23	2.22	0.40
1:A:1267:TRP:CD1	1:A:1267:TRP:C	2.95	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	262/310 (84%)	258 (98%)	4 (2%)	0	100	100
1	B	263/310 (85%)	259 (98%)	3 (1%)	1 (0%)	34	54
All	All	525/620 (85%)	517 (98%)	7 (1%)	1 (0%)	47	68

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	1258	GLN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	235/276 (85%)	221 (94%)	14 (6%)	19	37
1	B	235/276 (85%)	216 (92%)	19 (8%)	11	23
All	All	470/552 (85%)	437 (93%)	33 (7%)	15	29

All (33) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1074	SER
1	A	1091	CYS
1	A	1105	ILE
1	A	1107	CYS
1	A	1112	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	1148	ARG
1	A	1157	LEU
1	A	1171	ASN
1	A	1173	THR
1	A	1222	ASP
1	A	1257	THR
1	A	1258	GLN
1	A	1287	VAL
1	A	1293	THR
1	B	1058	LEU
1	B	1070	VAL
1	B	1105	ILE
1	B	1107	CYS
1	B	1115	ILE
1	B	1116	THR
1	B	1126	THR
1	B	1146	CYS
1	B	1212	LEU
1	B	1222	ASP
1	B	1257	THR
1	B	1258	GLN
1	B	1262	THR
1	B	1278	THR
1	B	1287	VAL
1	B	1290	PHE
1	B	1292	ILE
1	B	1300	ARG
1	B	1336	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1059	ASN
1	A	1123	GLN
1	A	1167	ASN
1	A	1171	ASN
1	A	1256	GLN
1	A	1298	GLN
1	A	1321	HIS
1	B	1059	ASN
1	B	1256	GLN
1	B	1258	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	AM7	A	1	-	49,50,50	1.23	5 (10%)	58,69,69	1.36	9 (15%)
2	AM7	B	1	-	49,50,50	1.21	3 (6%)	58,69,69	1.14	5 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	AM7	A	1	-	-	4/20/29/29	0/6/6/6
2	AM7	B	1	-	-	2/20/29/29	0/6/6/6

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1	AM7	C11-C18	-4.58	1.41	1.49

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1	AM7	C11-C18	-4.58	1.41	1.49
2	A	1	AM7	C13-N2	2.62	1.42	1.36
2	A	1	AM7	C12-N2	2.53	1.42	1.38
2	B	1	AM7	C12-N2	2.51	1.42	1.38
2	A	1	AM7	C26-N4	-2.05	1.33	1.37
2	A	1	AM7	C22-C23	2.04	1.40	1.36
2	B	1	AM7	C25-C24	2.02	1.40	1.36

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1	AM7	C1-N1-C4	4.18	118.24	108.83
2	A	1	AM7	C2-C1-N1	3.42	115.28	110.10
2	B	1	AM7	C27-N4-C26	2.98	121.54	116.93
2	B	1	AM7	C11-C12-N2	2.89	119.93	116.33
2	A	1	AM7	O1-C3-C4	-2.77	105.69	111.80
2	B	1	AM7	C21-C26-N4	-2.54	120.13	122.83
2	A	1	AM7	C27-N4-C26	2.51	120.81	116.93
2	A	1	AM7	O2-C23-C22	-2.35	122.20	125.24
2	A	1	AM7	C28-C29-C21	-2.33	117.06	121.15
2	A	1	AM7	C11-C12-N2	2.29	119.19	116.33
2	B	1	AM7	O2-C23-C22	-2.25	122.32	125.24
2	A	1	AM7	C28-C27-N4	-2.24	121.11	124.58
2	B	1	AM7	C28-C27-N4	-2.10	121.33	124.58
2	A	1	AM7	O2-C23-C24	2.07	118.29	115.41

There are no chirality outliers.

All (6) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	1	AM7	C24-C23-O2-C35
2	B	1	AM7	C22-C23-O2-C35
2	A	1	AM7	C24-C23-O2-C35
2	A	1	AM7	C32-C33-N1-C4
2	A	1	AM7	C32-C33-N1-C1
2	A	1	AM7	C22-C23-O2-C35

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1	AM7	1	0
2	B	1	AM7	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	268/310 (86%)	0.22	10 (3%) 41 45	39, 52, 60, 66	0
1	B	269/310 (86%)	0.19	12 (4%) 33 36	39, 52, 60, 66	0
All	All	537/620 (86%)	0.21	22 (4%) 37 40	39, 52, 60, 66	0

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1290	PHE	4.8
1	A	1326	MET	4.5
1	B	1257	THR	3.7
1	A	1181	LEU	3.1
1	A	1100	ASN	2.7
1	A	1319	CYS	2.6
1	A	1257	THR	2.6
1	A	1057	ALA	2.5
1	B	1326	MET	2.5
1	B	1181	LEU	2.3
1	A	1325	GLU	2.3
1	B	1115	ILE	2.3
1	B	1266	VAL	2.3
1	A	1256	GLN	2.3
1	B	1090	GLY	2.3
1	B	1298	GLN	2.2
1	B	1307	TYR	2.2
1	B	1116	THR	2.2
1	A	1348	HIS	2.2
1	B	1351	HIS	2.2
1	B	1114	ARG	2.1
1	A	1089	PHE	2.1



## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	AM7	A	1	45/45	0.94	0.13	31,34,48,49	0
2	AM7	B	1	45/45	0.94	0.13	35,38,49,50	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers

There are no such residues in this entry.