



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2024 – 10:55 PM EST

PDB ID : 3RFC
Title : Crystal structure of D-alanine-D-alanine ligase A from *Xanthomonas oryzae* pathovar *oryzae* with ADP
Authors : Doan, T.T.N.; Kim, J.K.; Ahn, Y.J.; Kang, L.W.
Deposited on : 2011-04-06
Resolution : 2.10 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	384	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 2763 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called D-alanine–D-alanine ligase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	344	2597	1639	474	475	9	0	0	0

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-16	HIS	-	expression tag	UNP Q2P8P8
A	-15	HIS	-	expression tag	UNP Q2P8P8
A	-14	HIS	-	expression tag	UNP Q2P8P8
A	-13	HIS	-	expression tag	UNP Q2P8P8
A	-12	HIS	-	expression tag	UNP Q2P8P8
A	-11	HIS	-	expression tag	UNP Q2P8P8
A	-10	HIS	-	expression tag	UNP Q2P8P8
A	-9	SER	-	expression tag	UNP Q2P8P8
A	-8	SER	-	expression tag	UNP Q2P8P8
A	-7	GLU	-	expression tag	UNP Q2P8P8
A	-6	ASN	-	expression tag	UNP Q2P8P8
A	-5	LEU	-	expression tag	UNP Q2P8P8
A	-4	TYR	-	expression tag	UNP Q2P8P8
A	-3	PHE	-	expression tag	UNP Q2P8P8
A	-2	GLN	-	expression tag	UNP Q2P8P8
A	-1	GLY	-	expression tag	UNP Q2P8P8
A	0	HIS	-	expression tag	UNP Q2P8P8
A	327	VAL	MET	engineered mutation	UNP Q2P8P8

- Molecule 2 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mg	0	0
			1	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	138	Total	O	0	0
			138	138		

4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	83.24Å 83.24Å 98.28Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.32 – 2.10 38.32 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.7 (38.32-2.10) 99.7 (38.32-2.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.77 (at 2.10Å)	Xtrriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.204 , 0.282 0.213 , 0.286	Depositor DCC
R_{free} test set	1063 reflections (5.14%)	wwPDB-VP
Wilson B-factor (Å ²)	32.3	Xtrriage
Anisotropy	0.080	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 46.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2763	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.98% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.01	0/2638	0.93	2/3583 (0.1%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	210	LEU	CA-CB-CG	5.63	128.26	115.30
1	A	31	LEU	CA-CB-CG	5.21	127.29	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2597	0	2642	62	0
2	A	27	0	12	0	0
3	A	1	0	0	0	0
4	A	138	0	0	4	0
All	All	2763	0	2654	62	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:340:ARG:HH11	1:A:340:ARG:HG3	0.99	1.08
1:A:330:LYS:HG3	4:A:425:HOH:O	1.73	0.88
1:A:13:LYS:HG2	1:A:45:LYS:O	1.77	0.84
1:A:66:ARG:HD3	4:A:467:HOH:O	1.75	0.83
1:A:340:ARG:HG3	1:A:340:ARG:NH1	1.76	0.83
1:A:190:GLY:HA2	1:A:191:SER:OG	1.84	0.77
1:A:340:ARG:HH11	1:A:340:ARG:CG	1.91	0.74
1:A:188:ASN:OD1	1:A:188:ASN:C	2.31	0.68
1:A:119:LEU:HD23	1:A:119:LEU:C	2.14	0.67
1:A:139:MET:HE3	1:A:188:ASN:HD21	1.60	0.67
1:A:308:ASP:N	1:A:308:ASP:OD1	2.29	0.64
1:A:276:ALA:O	1:A:280:GLN:HG2	1.98	0.63
1:A:161:PHE:HD1	1:A:165:THR:HG23	1.64	0.63
1:A:189:GLN:NE2	1:A:315:GLU:OE1	2.32	0.62
1:A:190:GLY:HA2	1:A:191:SER:CB	2.27	0.61
1:A:94:GLN:N	4:A:393:HOH:O	2.34	0.60
1:A:90:ILE:O	1:A:91:GLN:HB2	2.02	0.59
1:A:182:LEU:HB2	1:A:198:VAL:HG22	1.85	0.58
1:A:30:ALA:O	1:A:340:ARG:HG3	2.03	0.58
1:A:139:MET:CE	1:A:188:ASN:HD21	2.16	0.58
1:A:87:LEU:HD13	1:A:125:LEU:HD11	1.84	0.58
1:A:87:LEU:CD1	1:A:125:LEU:HD11	2.35	0.56
1:A:227:ARG:NH1	1:A:246:GLU:OE1	2.37	0.56
1:A:162:ASP:H	1:A:165:THR:HG22	1.71	0.56
1:A:189:GLN:O	1:A:189:GLN:HG3	2.06	0.54
1:A:227:ARG:NH2	1:A:274:ILE:HG22	2.23	0.54
1:A:26:ASN:OD1	1:A:324:ARG:N	2.41	0.53
1:A:212:LEU:HG	1:A:218:VAL:HG21	1.91	0.53
1:A:30:ALA:O	1:A:340:ARG:NH1	2.41	0.52
1:A:317:ASN:ND2	1:A:319:LEU:H	2.06	0.52
1:A:244:CYS:H	1:A:279:GLN:NE2	2.09	0.51
1:A:351:LEU:O	1:A:355:THR:HG23	2.11	0.51
1:A:171:VAL:HG13	1:A:204:PHE:CD2	2.46	0.51
1:A:308:ASP:N	1:A:309:GLY:HA2	2.26	0.51
1:A:188:ASN:OD1	1:A:190:GLY:N	2.44	0.50
1:A:188:ASN:OD1	1:A:188:ASN:O	2.29	0.50
1:A:227:ARG:HD2	1:A:246:GLU:OE2	2.11	0.49
1:A:4:ILE:O	1:A:36:PHE:HA	2.13	0.49
1:A:118:GLY:O	1:A:122:MET:HG2	2.13	0.48
1:A:139:MET:O	1:A:139:MET:HG3	2.13	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:145:LYS:HE3	1:A:221:GLU:OE1	2.13	0.48
1:A:168:HIS:HD2	4:A:444:HOH:O	1.96	0.48
1:A:238:VAL:HG13	1:A:238:VAL:O	2.15	0.47
1:A:361:ARG:O	1:A:361:ARG:HG3	2.14	0.47
1:A:57:PHE:CE2	1:A:58:LEU:HG	2.50	0.47
1:A:162:ASP:OD2	1:A:165:THR:HB	2.14	0.46
1:A:348:GLU:O	1:A:352:GLU:HG3	2.16	0.46
1:A:306:CYS:O	1:A:309:GLY:HA3	2.17	0.45
1:A:87:LEU:HD12	1:A:87:LEU:N	2.31	0.45
1:A:119:LEU:HD23	1:A:119:LEU:O	2.16	0.45
1:A:139:MET:CE	1:A:188:ASN:ND2	2.79	0.45
1:A:142:ASP:HB2	1:A:219:LEU:HD11	1.98	0.45
1:A:328:TYR:HB3	1:A:329:PRO:HD3	2.00	0.44
1:A:90:ILE:O	1:A:91:GLN:CB	2.66	0.43
1:A:244:CYS:H	1:A:279:GLN:HE21	1.67	0.43
1:A:317:ASN:HD22	1:A:318:THR:N	2.15	0.43
1:A:162:ASP:HA	1:A:216:HIS:O	2.18	0.43
1:A:119:LEU:C	1:A:119:LEU:CD2	2.85	0.42
1:A:78:LEU:HD21	1:A:123:ALA:HB2	2.02	0.42
1:A:361:ARG:O	1:A:361:ARG:CG	2.69	0.41
1:A:5:ARG:HD3	1:A:98:GLN:O	2.21	0.41
1:A:139:MET:HE3	1:A:188:ASN:ND2	2.32	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	338/384 (88%)	330 (98%)	8 (2%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	270/305 (88%)	251 (93%)	19 (7%)	15 12

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	8	LEU
1	A	53	ASP
1	A	78	LEU
1	A	88	ARG
1	A	110	LEU
1	A	165	THR
1	A	171	VAL
1	A	191	SER
1	A	192	SER
1	A	210	LEU
1	A	249	VAL
1	A	308	ASP
1	A	312	VAL
1	A	317	ASN
1	A	340	ARG
1	A	357	ASP
1	A	360	LEU
1	A	362	SER
1	A	364	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	70	HIS
1	A	85	GLN
1	A	188	ASN
1	A	197	GLN
1	A	279	GLN
1	A	284	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	317	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	ADP	A	368	3	24,29,29	1.26	3 (12%)	29,45,45	1.61	7 (24%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ADP	A	368	3	-	3/12/32/32	0/3/3/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	368	ADP	O4'-C4'	-2.50	1.39	1.45
2	A	368	ADP	C2'-C1'	-2.25	1.50	1.53
2	A	368	ADP	C5-N7	-2.04	1.32	1.39

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	368	ADP	PA-O3A-PB	-3.47	120.91	132.83
2	A	368	ADP	N3-C2-N1	-3.04	123.93	128.68
2	A	368	ADP	C4-C5-N7	-2.86	106.42	109.40
2	A	368	ADP	O3B-PB-O1B	2.69	121.19	110.68
2	A	368	ADP	C3'-C2'-C1'	2.56	104.83	100.98
2	A	368	ADP	O2'-C2'-C3'	-2.31	104.34	111.82
2	A	368	ADP	O3B-PB-O2B	2.02	115.35	107.64

There are no chirality outliers.


All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	368	ADP	O4'-C4'-C5'-O5'
2	A	368	ADP	PB-O3A-PA-O1A
2	A	368	ADP	PB-O3A-PA-O2A

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	344/384 (89%)	0.53	29 (8%) 11 14	16, 33, 55, 69	1 (0%)

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	248	VAL	4.5
1	A	362	SER	3.5
1	A	94	GLN	3.5
1	A	168	HIS	3.2
1	A	360	LEU	2.9
1	A	249	VAL	2.9
1	A	361	ARG	2.9
1	A	195	VAL	2.8
1	A	55	ASP	2.7
1	A	164	HIS	2.7
1	A	177	GLN	2.5
1	A	34	GLN	2.5
1	A	167	ALA	2.5
1	A	71	ARG	2.5
1	A	191	SER	2.5
1	A	194	GLY	2.5
1	A	180	LEU	2.4
1	A	275	ASP	2.4
1	A	116	LEU	2.3
1	A	190	GLY	2.3
1	A	59	LEU	2.2
1	A	120	LEU	2.2
1	A	280	GLN	2.1
1	A	133	LEU	2.1
1	A	359	LEU	2.1
1	A	132	VAL	2.1
1	A	119	LEU	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	203	ALA	2.0
1	A	276	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	MG	A	369	1/1	0.71	0.15	54,54,54,54	0
2	ADP	A	368	27/27	0.80	0.19	36,57,78,78	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.