



# Full wwPDB X-ray Structure Validation Report ⓘ

Jul 20, 2019 – 07:30 AM EDT

PDB ID : 1RGI  
Title : Crystal structure of gelsolin domains G1-G3 bound to actin  
Authors : Burtnick, L.D.; Urosev, D.; Irobi, E.; Narayan, K.; Robinson, R.C.  
Deposited on : 2003-11-12  
Resolution : 3.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.0 (224370), CSD as540be (2019)  
Xtriage (Phenix) : 1.13  
EDS : 2.3.2  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.3.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	1851 (3.00-3.00)
Clashscore	122126	2167 (3.00-3.00)
Ramachandran outliers	120053	2101 (3.00-3.00)
Sidechain outliers	120020	2104 (3.00-3.00)
RSRZ outliers	108989	1751 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	G	346	 2% 72% 25% . .
2	A	377	 69% 21% . 7%

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5541 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Gelsolin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	G	346	2740	1744	477	511	8	0	0	0

- Molecule 2 is a protein called Actin, alpha skeletal muscle.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	A	349	2740	1739	457	526	18	0	0	0

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	G	3	Total	Ca	0	0
			3	3		
3	A	1	Total	Ca	0	0
			1	1		


- Molecule 4 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: C<sub>10</sub>H<sub>16</sub>N<sub>5</sub>O<sub>13</sub>P<sub>3</sub>).




### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Gelsolin



- Molecule 2: Actin, alpha skeletal muscle



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	145.25Å 145.25Å 129.95Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	19.96 – 3.00 20.03 – 3.00	Depositor EDS
% Data completeness (in resolution range)	97.5 (19.96-3.00) 97.5 (20.03-3.00)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.06	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.52 (at 2.98Å)	Xtrriage
Refinement program	REFMAC 5.1.24, CNS	Depositor
R, $R_{free}$	0.224 , 0.258 0.194 , 0.225	Depositor DCC
$R_{free}$ test set	1570 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	80.0	Xtrriage
Anisotropy	0.254	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 49.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.016 for -h,-k,l	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	5541	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	31.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.16% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	G	0.64	0/2802	0.79	12/3784 (0.3%)
2	A	0.66	0/2797	0.86	14/3785 (0.4%)
All	All	0.65	0/5599	0.83	26/7569 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	G	0	1

There are no bond length outliers.

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
2	A	25	ASP	CB-CG-OD2	8.55	126.00	118.30
1	G	187	ASP	CB-CG-OD2	8.01	125.51	118.30
2	A	56	ASP	CB-CG-OD2	7.45	125.00	118.30
2	A	81	ASP	CB-CG-OD2	6.74	124.36	118.30
2	A	244	ASP	CB-CG-OD2	6.51	124.16	118.30
1	G	371	ASP	CB-CG-OD2	6.50	124.15	118.30
1	G	259	ASP	CB-CG-OD2	6.41	124.07	118.30
1	G	290	ASP	CB-CG-OD2	6.26	123.94	118.30
2	A	184	ASP	CB-CG-OD2	5.94	123.65	118.30
2	A	157	ASP	CB-CG-OD2	5.91	123.62	118.30
1	G	264	ASP	CB-CG-OD2	5.73	123.46	118.30
2	A	222	ASP	CB-CG-OD2	5.71	123.44	118.30
2	A	288	ASP	CB-CG-OD2	5.68	123.41	118.30
1	G	312	ASP	CB-CG-OD2	5.66	123.40	118.30

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	11	ASP	CB-CG-OD2	5.60	123.34	118.30
2	A	179	ASP	CB-CG-OD2	5.60	123.34	118.30
2	A	363	ASP	CB-CG-OD2	5.57	123.31	118.30
2	A	154	ASP	CB-CG-OD2	5.56	123.31	118.30
1	G	308	ASP	CB-CG-OD2	5.49	123.24	118.30
1	G	109	ASP	CB-CG-OD2	5.43	123.19	118.30
2	A	51	ASP	CB-CG-OD2	5.43	123.19	118.30
1	G	96	ASP	CB-CG-OD2	5.41	123.17	118.30
1	G	343	ASP	CB-CG-OD2	5.33	123.10	118.30
2	A	292	ASP	CB-CG-OD2	5.13	122.92	118.30
1	G	84	ASP	CB-CG-OD2	5.10	122.89	118.30
1	G	192	ASP	CB-CG-OD2	5.06	122.85	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	G	292	ASN	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	G	2740	0	2704	33	0
2	A	2740	0	2704	37	0
3	A	1	0	0	0	0
3	G	3	0	0	0	0
4	A	31	0	12	0	0
5	A	19	0	0	0	0
5	G	7	0	0	0	0
All	All	5541	0	5420	67	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:212:ILE:HD11	2:A:248:ILE:HD11	1.72	0.71
2:A:16:LEU:O	2:A:18:LYS:NZ	2.28	0.67
2:A:212:ILE:HD11	2:A:248:ILE:CD1	2.30	0.62
1:G:199:GLN:HE21	1:G:233:VAL:HG13	1.65	0.61
1:G:29:HIS:ND1	1:G:30:PRO:HD2	2.15	0.61
1:G:258:GLU:O	1:G:259:ASP:O	2.19	0.60
2:A:111:ASN:HD21	2:A:115:ASN:HD22	1.49	0.60
2:A:70:PRO:HG3	2:A:81:ASP:HB3	1.86	0.57
1:G:153:VAL:HG22	1:G:156:GLU:CB	2.35	0.57
2:A:282:ILE:HG22	2:A:290:ARG:HD3	1.86	0.57
1:G:153:VAL:HG22	1:G:156:GLU:HB3	1.87	0.56
1:G:110:ASP:OD2	2:A:169:TYR:OH	2.17	0.56
2:A:352:PHE:HE2	2:A:356:TRP:CH2	2.23	0.56
2:A:31:PHE:CE2	2:A:93:GLU:HG3	2.41	0.55
2:A:314:GLN:O	2:A:318:THR:CG2	2.55	0.55
2:A:194:THR:HA	2:A:198:TYR:O	2.07	0.55
1:G:367:LYS:O	1:G:368:ASN:C	2.46	0.54
1:G:270:LEU:HD22	1:G:309:HIS:CE1	2.44	0.53
1:G:206:ASN:ND2	1:G:209:GLU:OE1	2.43	0.51
1:G:270:LEU:HD22	1:G:309:HIS:HE1	1.75	0.51
2:A:218:TYR:OH	2:A:226:GLU:OE1	2.27	0.51
1:G:268:ARG:HA	1:G:292:ASN:HB2	1.93	0.49
1:G:221:ARG:HD3	1:G:231:VAL:HG23	1.93	0.49
2:A:207:GLU:CD	2:A:210:ARG:HH22	2.16	0.49
2:A:32:PRO:HB2	2:A:34:ILE:HG12	1.95	0.49
1:G:190:ILE:HD12	1:G:217:SER:OG	2.12	0.49
1:G:160:GLN:HE21	1:G:193:LEU:HD21	1.78	0.48
1:G:334:THR:O	1:G:337:ASP:HB2	2.13	0.48
1:G:153:VAL:CG2	1:G:156:GLU:HB2	2.43	0.48
1:G:80:ILE:HG22	1:G:81:LEU:H	1.78	0.48
1:G:283:MET:HE1	1:G:327:GLU:HG2	1.95	0.47
1:G:56:PRO:HA	1:G:59:TYR:CD1	2.49	0.47
2:A:314:GLN:O	2:A:318:THR:HG23	2.14	0.47
1:G:228:ARG:NH2	2:A:93:GLU:OE1	2.46	0.47
2:A:219:VAL:CG1	2:A:312:ARG:HG3	2.44	0.47
1:G:72:LYS:HG2	1:G:133:TYR:CE1	2.50	0.46
2:A:253:GLU:HA	2:A:256:ARG:HB2	1.97	0.46
2:A:314:GLN:O	2:A:318:THR:HG22	2.16	0.45
1:G:234:PHE:HB2	1:G:238:ALA:HB3	1.98	0.45
2:A:345:ILE:O	2:A:349:LEU:HG	2.16	0.45
1:G:292:ASN:HB3	1:G:293:PRO:CD	2.48	0.43
1:G:27:VAL:HG11	2:A:351:THR:HB	2.00	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:375:PHE:N	2:A:375:PHE:CD1	2.87	0.43
1:G:199:GLN:NE2	1:G:233:VAL:HG13	2.32	0.43
2:A:106:THR:HA	2:A:135:ALA:O	2.19	0.42
2:A:26:ALA:HB1	2:A:27:PRO:HD2	2.00	0.42
1:G:157:VAL:HG13	1:G:159:VAL:HG23	2.01	0.42
2:A:215:LYS:O	2:A:216:LEU:HD23	2.19	0.42
2:A:250:ILE:HG12	2:A:253:GLU:HG2	2.01	0.42
1:G:94:SER:OG	1:G:96:ASP:OD1	2.26	0.42
2:A:62:ARG:HH12	2:A:204:ALA:HB2	1.84	0.42
2:A:161:HIS:HA	2:A:176:MET:O	2.20	0.42
1:G:190:ILE:HG22	1:G:197:ILE:HG23	2.03	0.41
2:A:332:PRO:HA	2:A:333:PRO:HD3	1.93	0.41
2:A:219:VAL:HG12	2:A:312:ARG:HG3	2.03	0.41
2:A:240:TYR:HE1	2:A:242:LEU:CD1	2.32	0.41
1:G:56:PRO:HA	1:G:59:TYR:CE1	2.55	0.41
1:G:163:LEU:HD12	1:G:190:ILE:HG13	2.02	0.41
2:A:335:ARG:HA	2:A:338:SER:OG	2.21	0.41
1:G:275:LYS:HG3	1:G:305:PHE:CZ	2.55	0.41
2:A:211:ASP:O	2:A:215:LYS:HG3	2.21	0.41
2:A:299:MET:CE	2:A:309:ILE:HG23	2.51	0.41
1:G:185:ASN:HD21	1:G:253:LEU:HD22	1.85	0.41
2:A:240:TYR:CE1	2:A:242:LEU:HD13	2.56	0.40
1:G:275:LYS:HG3	1:G:305:PHE:CE2	2.56	0.40
2:A:190:MET:HG3	2:A:209:VAL:HG11	2.01	0.40
2:A:34:ILE:HD11	2:A:59:GLN:HG3	2.04	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	G	344/346 (99%)	314 (91%)	26 (8%)	4 (1%)	<b>14</b> 51

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	A	342/377 (91%)	323 (94%)	18 (5%)	1 (0%)	43	80
All	All	686/723 (95%)	637 (93%)	44 (6%)	5 (1%)	24	64

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	G	259	ASP
1	G	293	PRO
2	A	25	ASP
1	G	368	ASN
1	G	292	ASN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	G	291/291 (100%)	245 (84%)	46 (16%)	3	14
2	A	296/320 (92%)	266 (90%)	30 (10%)	8	31
All	All	587/611 (96%)	511 (87%)	76 (13%)	5	20

All (76) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	27	VAL
1	G	28	GLU
1	G	31	GLU
1	G	38	GLU
1	G	51	LEU
1	G	57	ASN
1	G	64	THR
1	G	74	VAL
1	G	78	ASN
1	G	82	GLN
1	G	84	ASP

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>
1	G	92	GLU
1	G	108	LEU
1	G	112	LEU
1	G	115	ARG
1	G	136	SER
1	G	149	PHE
1	G	150	LYS
1	G	152	VAL
1	G	153	VAL
1	G	166	LYS
1	G	168	ARG
1	G	169	ARG
1	G	170	VAL
1	G	190	ILE
1	G	232	SER
1	G	239	GLU
1	G	241	GLU
1	G	245	GLN
1	G	252	THR
1	G	253	LEU
1	G	292	ASN
1	G	299	LEU
1	G	300	ARG
1	G	307	LEU
1	G	311	LYS
1	G	314	LYS
1	G	336	SER
1	G	340	SER
1	G	341	LYS
1	G	342	MET
1	G	346	LYS
1	G	347	GLN
1	G	349	GLN
1	G	351	SER
1	G	371	ASP
2	A	12	ASN
2	A	18	LYS
2	A	25	ASP
2	A	28	ARG
2	A	33	SER
2	A	59	GLN
2	A	62	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	A	66	THR
2	A	118	LYS
2	A	126	THR
2	A	185	LEU
2	A	193	LEU
2	A	226	GLU
2	A	236	LEU
2	A	238	LYS
2	A	242	LEU
2	A	248	ILE
2	A	257	CYS
2	A	275	HIS
2	A	291	LYS
2	A	293	LEU
2	A	297	ASN
2	A	312	ARG
2	A	317	ILE
2	A	318	THR
2	A	324	THR
2	A	328	LYS
2	A	334	GLU
2	A	359	LYS
2	A	375	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	G	160	GLN
1	G	185	ASN
1	G	196	ASN
1	G	230	GLN
1	G	267	ASN
1	G	309	HIS
2	A	115	ASN

### 5.3.3 RNA

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 4 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	ATP	A	380	3	26,33,33	1.30	3 (11%)	27,52,52	1.72	4 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ATP	A	380	3	-	4/18/38/38	0/3/3/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	380	ATP	C2-N3	3.08	1.37	1.32
4	A	380	ATP	PG-O3B	2.91	1.64	1.60
4	A	380	ATP	C2-N1	2.16	1.38	1.33

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	380	ATP	N3-C2-N1	-6.73	117.84	128.68

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	380	ATP	PB-O3B-PG	-3.31	122.05	132.57
4	A	380	ATP	PA-O3A-PB	-2.72	123.92	132.57
4	A	380	ATP	C4-C5-N7	-2.10	107.21	109.40

There are no chirality outliers.


All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	380	ATP	PG-O3B-PB-O1B
4	A	380	ATP	PB-O3B-PG-O2G
4	A	380	ATP	PB-O3B-PG-O3G
4	A	380	ATP	PG-O3B-PB-O2B

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	G	346/346 (100%)	-0.35	6 (1%) 70 41	16, 31, 46, 72	0
2	A	349/377 (92%)	-0.55	1 (0%) 93 83	20, 30, 41, 73	0
All	All	695/723 (96%)	-0.45	7 (1%) 82 59	16, 31, 44, 73	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	257	THR	2.9
1	G	256	ALA	2.7
1	G	347	GLN	2.5
1	G	30	PRO	2.2
1	G	26	VAL	2.1
1	G	343	ASP	2.1
2	A	364	GLU	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	CA	G	406	1/1	0.92	0.17	47,47,47,47	0
3	CA	G	404	1/1	0.96	0.04	36,36,36,36	0
3	CA	A	401	1/1	0.97	0.42	31,31,31,31	0
4	ATP	A	380	31/31	0.98	0.11	17,23,30,31	0
3	CA	G	403	1/1	0.99	0.12	25,25,25,25	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.