



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 7, 2018 – 10:22 am GMT

PDB ID : 1RHG
Title : THE STRUCTURE OF GRANULOCYTE-COLON-STIMULATING FACTOR AND ITS RELATIONSHIP TO THOSE OF OTHER GROWTH FACTORS
Authors : Hill, C.P.; Osslund, T.D.; Eisenberg, D.
Deposited on : 1993-01-29
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : trunk30967
Percentile statistics : (not set)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30967


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	122126	5027 (2.20-2.20)
Ramachandran outliers	120053	4952 (2.20-2.20)
Sidechain outliers	120020	4953 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	174	
1	B	174	
1	C	174	

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 4078 atoms, of which 735 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GRANULOCYTE COLONY-STIMULATING FACTOR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	145	1320	716	215	182	200	7	0	0	0
1	B	144	1321	708	223	186	197	7	0	0	0
1	C	145	1317	711	217	184	198	7	0	0	0

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	VAL	DELETION	UNP P09919
A	?	-	SER	DELETION	UNP P09919
A	?	-	GLU	DELETION	UNP P09919
B	?	-	VAL	DELETION	UNP P09919
B	?	-	SER	DELETION	UNP P09919
B	?	-	GLU	DELETION	UNP P09919
C	?	-	VAL	DELETION	UNP P09919
C	?	-	SER	DELETION	UNP P09919
C	?	-	GLU	DELETION	UNP P09919

- Molecule 2 is water.


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	H	O		
2	A	13	39	26	13	0	0
2	B	13	39	26	13	0	0
2	C	14	42	28	14	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: GRANULOCYTE COLONY-STIMULATING FACTOR

Chain A: 




- Molecule 1: GRANULOCYTE COLONY-STIMULATING FACTOR

Chain B: 



- Molecule 1: GRANULOCYTE COLONY-STIMULATING FACTOR

Chain C: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	90.90Å 110.70Å 49.50Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	(Not available) – 2.20 47.28 – 2.04	Depositor EDS
% Data completeness (in resolution range)	(Not available) ((Not available)-2.20) 83.7 (47.28-2.04)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.66 (at 2.05Å)	Xtrriage
Refinement program	X-PLOR	Depositor
R, R_{free}	0.215 , (Not available) 0.281 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	36.9	Xtrriage
Anisotropy	0.172	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 97.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	4078	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.76% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.94	0/1127	1.59	17/1528 (1.1%)
1	B	0.94	1/1120 (0.1%)	1.70	18/1518 (1.2%)
1	C	0.95	1/1122 (0.1%)	1.70	21/1522 (1.4%)
All	All	0.94	2/3369 (0.1%)	1.67	56/4568 (1.2%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	98	GLU	CD-OE1	7.97	1.34	1.25
1	C	153	VAL	CA-CB	6.46	1.68	1.54

All (56) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	22	ARG	NE-CZ-NH1	11.30	125.95	120.30
1	C	147	ARG	NE-CZ-NH1	11.26	125.93	120.30
1	B	58	TRP	CD1-CG-CD2	10.51	114.71	106.30
1	A	146	ARG	NE-CZ-NH1	10.13	125.36	120.30
1	B	22	ARG	NE-CZ-NH2	8.97	124.78	120.30
1	C	118	TRP	CD1-CG-CD2	8.78	113.32	106.30
1	A	39	TYR	CB-CG-CD2	-8.58	115.85	121.00
1	B	58	TRP	CE2-CD2-CG	-8.43	100.55	107.30
1	C	22	ARG	NE-CZ-NH2	-8.21	116.20	120.30
1	A	118	TRP	CD1-CG-CD2	7.69	112.45	106.30
1	C	146	ARG	NE-CZ-NH2	-7.65	116.48	120.30
1	C	165	TYR	CB-CG-CD2	-7.53	116.48	121.00
1	A	58	TRP	CE2-CD2-CG	-7.52	101.28	107.30
1	A	146	ARG	NE-CZ-NH2	-7.43	116.58	120.30
1	B	17	CYS	CA-CB-SG	-7.41	100.67	114.00
1	A	58	TRP	CD1-CG-CD2	7.36	112.18	106.30
1	A	118	TRP	CE2-CD2-CG	-7.21	101.53	107.30
1	C	118	TRP	CE2-CD2-CG	-7.13	101.59	107.30
1	B	118	TRP	CE2-CD2-CG	-7.01	101.69	107.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	58	TRP	CE2-CD2-CG	-6.99	101.71	107.30
1	C	147	ARG	NE-CZ-NH2	-6.90	116.85	120.30
1	A	22	ARG	NE-CZ-NH1	-6.88	116.86	120.30
1	B	118	TRP	CD1-CG-CD2	6.76	111.71	106.30
1	A	137	MET	CG-SD-CE	6.72	110.96	100.20
1	C	58	TRP	CD1-CG-CD2	6.61	111.59	106.30
1	C	58	TRP	CG-CD2-CE3	6.60	139.84	133.90
1	B	145	GLN	CA-CB-CG	-6.53	99.03	113.40
1	A	19	GLU	CA-CB-CG	6.47	127.64	113.40
1	B	58	TRP	CG-CD1-NE1	-6.45	103.66	110.10
1	A	118	TRP	CG-CD2-CE3	6.29	139.56	133.90
1	A	118	TRP	CB-CG-CD1	-6.25	118.87	127.00
1	C	145	GLN	CA-CB-CG	-6.25	99.65	113.40
1	C	169	ARG	NE-CZ-NH1	6.20	123.40	120.30
1	B	37	ALA	CB-CA-C	-6.19	100.82	110.10
1	C	118	TRP	CG-CD1-NE1	-6.15	103.95	110.10
1	B	85	TYR	CB-CG-CD1	-6.10	117.34	121.00
1	C	155	SER	N-CA-CB	-6.08	101.38	110.50
1	A	140	PHE	N-CA-C	-6.05	94.67	111.00
1	C	58	TRP	CB-CG-CD1	-5.98	119.22	127.00
1	A	118	TRP	CG-CD1-NE1	-5.73	104.37	110.10
1	B	112	ASP	CB-CA-C	-5.65	99.10	110.40
1	A	124	LEU	CA-CB-CG	5.63	128.25	115.30
1	C	41	LEU	N-CA-C	-5.63	95.81	111.00
1	B	9	LEU	CA-CB-CG	5.55	128.07	115.30
1	A	152	LEU	CB-CG-CD2	-5.49	101.67	111.00
1	B	121	MET	CA-CB-CG	-5.48	103.99	113.30
1	B	141	ALA	CA-C-N	5.44	129.17	117.20
1	C	48	VAL	CG1-CB-CG2	-5.42	102.23	110.90
1	C	139	ALA	N-CA-CB	5.36	117.61	110.10
1	B	110	VAL	CG1-CB-CG2	-5.34	102.35	110.90
1	A	19	GLU	N-CA-CB	-5.33	101.00	110.60
1	C	118	TRP	CB-CG-CD1	-5.30	120.11	127.00
1	B	50	LEU	CB-CG-CD2	-5.26	102.06	111.00
1	B	95	ILE	CA-C-N	5.06	128.33	117.20
1	C	74	CYS	CA-CB-SG	-5.04	104.92	114.00
1	B	37	ALA	N-CA-CB	5.00	117.11	110.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1105	215	1111	21	0
1	B	1098	223	1099	23	0
1	C	1100	217	1098	16	0
2	A	13	26	0	1	0
2	B	13	26	0	0	0
2	C	14	28	0	0	0
All	All	3343	735	3308	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:61:LEU:HD12	1:C:163:VAL:HG12	1.46	0.96
1:B:137:MET:HG3	1:B:138:PRO:HD2	1.57	0.85
1:A:14:LEU:HD22	1:A:172:ALA:HB3	1.68	0.76
1:A:103:LEU:O	1:A:107:GLN:HG3	1.99	0.62
1:C:141:ALA:HB3	1:C:145:GLN:OE1	2.02	0.60
1:A:164:SER:HA	1:A:167:VAL:HG22	1.86	0.57
1:C:95:ILE:HG13	1:C:103:LEU:HD11	1.86	0.57
1:C:61:LEU:HD12	1:C:163:VAL:CG1	2.29	0.56
1:A:162:GLU:O	1:A:165:TYR:HB3	2.07	0.54
1:B:31:LEU:HD11	1:B:103:LEU:HD12	1.92	0.52
1:B:49:LEU:O	1:B:52:HIS:HB3	2.10	0.51
1:A:95:ILE:HD12	1:A:103:LEU:HD11	1.91	0.51
1:C:86:GLN:HE22	1:C:107:GLN:HG2	1.75	0.51
1:B:54:LEU:HD21	1:B:145:GLN:HG2	1.92	0.50
1:B:100:GLY:N	1:B:101:PRO:HD2	2.26	0.50
1:A:36:CYS:HA	1:A:41:LEU:O	2.12	0.50
1:A:155:SER:O	1:A:158:GLN:HB3	2.12	0.49
1:B:108:LEU:O	1:B:111:ALA:HB3	2.13	0.49
1:B:85:TYR:OH	1:B:156:HIS:HB3	2.13	0.49
1:B:49:LEU:H	1:B:49:LEU:HD12	1.77	0.48
1:A:162:GLU:HA	1:A:162:GLU:OE1	2.12	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:22:ARG:NH2	1:C:83:PHE:CE2	2.81	0.48
1:C:14:LEU:HD11	1:C:172:ALA:HA	1.95	0.48
1:B:38:THR:HG22	1:B:39:TYR:CE1	2.49	0.47
1:B:22:ARG:NH1	1:C:83:PHE:O	2.47	0.47
1:B:9:LEU:N	1:B:171:LEU:O	2.48	0.47
1:B:27:ASP:HB2	1:B:106:LEU:HD13	1.96	0.47
1:B:9:LEU:HD22	1:B:171:LEU:O	2.15	0.47
1:A:115:THR:O	1:A:119:GLN:HG3	2.14	0.46
1:A:167:VAL:O	1:A:171:LEU:HB2	2.14	0.46
1:B:22:ARG:NH1	1:C:87:GLY:N	2.62	0.46
1:C:45:GLU:O	1:C:48:VAL:HG12	2.16	0.46
1:A:35:LEU:HD13	1:A:151:VAL:HG23	1.98	0.46
2:A:619:HOH:O	1:C:162:GLU:HG2	2.16	0.46
1:B:92:LEU:HD23	1:B:140:PHE:CD2	2.51	0.45
1:B:11:GLN:O	1:B:15:LEU:HD12	2.16	0.45
1:A:78:LEU:HD23	1:A:82:LEU:HD11	1.98	0.45
1:A:14:LEU:HD21	1:A:168:LEU:O	2.17	0.45
1:B:31:LEU:HD23	1:B:154:ALA:HB2	1.99	0.44
1:A:39:TYR:HB2	1:A:41:LEU:HD12	1.99	0.44
1:A:71:LEU:HD12	1:A:71:LEU:HA	1.76	0.43
1:B:137:MET:CG	1:B:138:PRO:HD2	2.40	0.43
1:B:14:LEU:HD21	1:B:169:ARG:HD3	2.01	0.43
1:B:96:SER:HB2	1:B:97:PRO:HD2	2.01	0.43
1:B:22:ARG:NH1	1:C:87:GLY:CA	2.82	0.42
1:C:55:GLY:C	1:C:138:PRO:HG3	2.40	0.42
1:A:58:TRP:CE3	1:A:58:TRP:HA	2.54	0.42
1:A:71:LEU:HD23	1:A:124:LEU:HD21	2.02	0.41
1:C:72:ALA:HA	1:C:126:MET:SD	2.61	0.41
1:A:151:VAL:O	1:A:155:SER:HB3	2.20	0.41
1:A:27:ASP:HB2	1:A:106:LEU:HD13	2.02	0.41
1:C:90:GLN:HG3	1:C:107:GLN:NE2	2.36	0.41
1:A:88:LEU:HA	1:A:88:LEU:HD23	1.87	0.41
1:C:71:LEU:O	1:C:73:GLY:N	2.54	0.40
1:A:124:LEU:HG	1:A:126:MET:SD	2.60	0.40
1:B:21:VAL:HG13	1:B:161:LEU:HB3	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	139/174 (80%)	134 (96%)	4 (3%)	1 (1%)	24	23
1	B	138/174 (79%)	128 (93%)	6 (4%)	4 (3%)	5	2
1	C	139/174 (80%)	130 (94%)	8 (6%)	1 (1%)	24	23
All	All	416/522 (80%)	392 (94%)	18 (4%)	6 (1%)	12	9

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	72	ALA
1	B	141	ALA
1	C	72	ALA
1	A	11	GLN
1	B	95	ILE
1	B	138	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	117/141 (83%)	105 (90%)	12 (10%)	8	7
1	B	116/141 (82%)	102 (88%)	14 (12%)	5	4
1	C	115/141 (82%)	101 (88%)	14 (12%)	5	4
All	All	348/423 (82%)	308 (88%)	40 (12%)	6	5

All (40) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	11	GLN
1	A	12	SER
1	A	14	LEU
1	A	33	GLU
1	A	45	GLU
1	A	46	GLU
1	A	98	GLU
1	A	137	MET
1	A	142	SER
1	A	152	LEU
1	A	155	SER
1	A	159	SER
1	B	9	LEU
1	B	11	GLN
1	B	22	ARG
1	B	45	GLU
1	B	49	LEU
1	B	50	LEU
1	B	52	HIS
1	B	62	SER
1	B	74	CYS
1	B	77	GLN
1	B	78	LEU
1	B	83	PHE
1	B	103	LEU
1	B	142	SER
1	C	9	LEU
1	C	11	GLN
1	C	14	LEU
1	C	53	SER
1	C	61	LEU
1	C	90	GLN
1	C	93	GLU
1	C	98	GLU
1	C	126	MET
1	C	146	ARG
1	C	153	VAL
1	C	165	TYR
1	C	167	VAL
1	C	170	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	20	GLN
1	A	77	GLN
1	A	158	GLN
1	B	11	GLN
1	B	20	GLN
1	B	77	GLN
1	C	86	GLN
1	C	107	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.