

wwPDB X-ray Structure Validation Summary Report (i)

Sep 24, 2024 - 07:24 pm BST

PDB ID	:	8RIU
Title	:	Crystal structure of the F420-reducing carbon monoxide dehydrogenase com-
		ponent from the ethanotroph Candidatus Ethanoperedens thermophilum
Authors	:	Lemaire, O.N.; Wagner, T.
Deposited on	:	2023-12-19
Resolution	:	1.89 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	3.0
buster-report	:	1.1.7(2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.002 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.38.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 1.89 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Matria	Whole archive	Similar resolution
Metric	$(\# { m Entries})$	$(\# { m Entries}, { m resolution} { m range}({ m \AA}))$
R _{free}	164625	7293(1.90-1.90)
Clashscore	180529	8090 (1.90-1.90)
Ramachandran outliers	177936	8022 (1.90-1.90)
Sidechain outliers	177891	8022 (1.90-1.90)
RSRZ outliers	164620	7292 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	А	792	90%	6% •
1	D	792	90%	6% •
2	В	370	2% 95%	5%
2	Е	370	94%	6%
3	С	174	90%	10% •



Mol	Chain	Length	Quality of chain	
	I		%	
3	F,	174	95%	5%



2 Entry composition (i)

There are 12 unique types of molecules in this entry. The entry contains 21942 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Acetyl-CoA decarbonylase/synthase complex subunit alpha.

Mol	Chain	Residues		Α	toms			ZeroOcc	AltConf	Trace
1	А	759	Total 5859	C 3706	N 1011	O 1091	$\begin{array}{c} \mathrm{S} \\ 51 \end{array}$	0	0	0
1	D	760	Total 5902	C 3732	N 1023	O 1096	S 51	0	4	0

• Molecule 2 is a protein called Coenzyme F420 hydrogenase/dehydrogenase, beta subunit C terminus.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	В	370	Total	С	Ν	0	\mathbf{S}	0	0	0
			2843	1791	475	553	24	0		
2	F	370	Total	С	Ν	0	\mathbf{S}	0	1	0
	Ľ	570	2854	1797	479	554	24	0		0

• Molecule 3 is a protein called Acetyl-CoA decarbonylase/synthase complex subunit epsilon.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
3	C	174	Total	С	Ν	0	S	0	2	0
0		114	1352	859	233	253	7	0		
2	Б	174	Total	С	Ν	0	S	0	1	0
0	Ľ	1/4	1343	855	231	249	8			U

• Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 6 3 3 \end{array}$	0	0
4	А	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 6 3 3 \end{array}$	0	0
4	А	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 6 3 3 \end{array}$	0	0
4	А	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 6 3 3 \end{array}$	0	0
4	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 6 & 3 & 3 \end{array}$	0	0
4	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 6 & 3 & 3 \end{array}$	0	0
4	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 6 & 3 & 3 \end{array}$	0	0
4	В	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 6 & 3 & 3 \end{array}$	0	0
4	С	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 6 & 3 & 3 \end{array}$	0	0
4	С	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 6 3 3 \end{array}$	0	0
4	С	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 6 3 3 \end{array}$	0	0
4	D	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 6 & 3 & 3 \end{array}$	0	0
4	Е	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 6 & 3 & 3 \end{array}$	0	0
4	F	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 6 & 3 & 3 \end{array}$	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	F	1	Total 6	${ m C} { m 3}$	O 3	0	0

• Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0
5	А	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 4 2 2 \end{array}$	0	0
5	А	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 4 2 2 \end{array}$	0	0
5	А	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 4 2 2 \end{array}$	0	0
5	А	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 4 2 2 \end{array}$	0	0
5	А	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 4 2 2 \end{array}$	0	0
5	А	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 4 2 2 \end{array}$	0	0
5	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0
5	В	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0
5	D	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	F	1	Total 4	${ m C} 2$	O 2	0	0

• Molecule 6 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe_4S_4) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	А	1	TotalFeS844	0	0
6	А	1	TotalFeS844	0	0
6	А	1	TotalFeS844	0	0
6	А	1	TotalFeS844	0	0
6	В	1	TotalFeS844	0	0
6	В	1	TotalFeS844	0	0
6	В	1	TotalFeS844	0	0
6	D	1	TotalFeS844	0	0
6	D	1	TotalFeS844	0	0
6	D	1	TotalFeS844	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	Е	1	TotalFeS844	0	0
6	Е	1	TotalFeS844	0	0
6	Е	1	TotalFeS844	0	0

• Molecule 7 is FE(4)-NI(1)-S(4) CLUSTER (three-letter code: XCC) (formula: Fe_4NiS_4) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	А	1	Total 9	Fe 4	Ni 1	${S \atop 4}$	0	0
7	D	1	Total 9	Fe 4	Ni 1	$\frac{S}{4}$	0	0

• Molecule 8 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	А	1	Total K 1 1	0	0
8	D	1	Total K 1 1	0	0

• Molecule 9 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf		
0	В	1	Total	С	Ν	Ο	Р	0	0	
9	D	I	53	27	9	15	2	0	0	
0	F	1	Total	С	Ν	Ο	Р	0	0	
9	Ľ	L	53	27	9	15	2	0	0	

• Molecule 10 is 2-{2-[2-(2-{2-[2-(2-ETHOXY-ETHOXY)-ETHOXY]-ETHOXY}PA



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	
10	D	1	Total 6	$\begin{array}{c} \mathrm{C} \\ 4 \end{array}$	O 2	0	0



• Molecule 11 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
11	D	1	Total Cl 1 1	0	1

• Molecule 12 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
12	А	455	Total O 455 455	0	0
12	В	140	Total O 140 140	0	0
12	С	160	Total O 160 160	0	1
12	D	351	Total O 351 351	0	0
12	Ε	179	Total O 179 179	0	0
12	F	133	Total O 133 133	0	1



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Acetyl-CoA decarbonylase/synthase complex subunit alpha



• Molecule 2: Coenzyme F420 hydrogenase/dehydrogenase, beta subunit C terminus



Chain E:	94%	6%
M1 738 738 738 738 738 790 791 792 7131 7131	L195 L195 L213 E235 E235 E235 E236 1260 E236 C287 C287 C287 C287 C287 C287 C287 C287	0344 6370
• Molecule 3: Acety	l-CoA decarbonylase/synthase comple	ex subunit epsilon
Chain C:	90%	10% •
11 1641 1644 168 71 71 882	H86 192 192 192 192 193 1134 1157 1134 1157 1157 1157 1157 1157 1157 1157	
• Molecule 3: Acety	l-CoA decarbonylase/synthase comple	ex subunit epsilon
Chain F:	95%	5%
10 1945 192 192 192 192 192 192 192 192		



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	97.07Å 159.21Å 191.44Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Bosolution(A)	39.80 - 1.89	Depositor
Resolution (A)	39.80 - 1.89	EDS
% Data completeness	70.0(39.80-1.89)	Depositor
(in resolution range)	70.0(39.80-1.89)	EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$1.70 (at 1.90 \text{\AA})$	Xtriage
Refinement program	PHENIX (1.20.1_4487: ???)	Depositor
P. P.	0.165 , 0.187	Depositor
n, n_{free}	0.170 , 0.188	DCC
R_{free} test set	11819 reflections (5.03%)	wwPDB-VP
Wilson B-factor $(Å^2)$	37.6	Xtriage
Anisotropy	0.022	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.34 , 42.7	EDS
L-test for $twinning^2$	$ < L >=0.49, < L^2>=0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	21942	wwPDB-VP
Average B, all atoms $(Å^2)$	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 2.51% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: K, CL, PE4, GOL, FAD, EDO, XCC, SF4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond	lengths	Bond angles		
	Unam	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.43	0/5972	0.61	0/8088	
1	D	0.41	0/6021	0.59	0/8152	
2	В	0.42	0/2886	0.58	0/3885	
2	Е	0.39	0/2897	0.58	0/3899	
3	С	0.42	0/1379	0.60	0/1861	
3	F	0.40	0/1370	0.60	0/1848	
All	All	0.41	0/20525	0.60	0/27733	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	5859	0	5870	31	0
1	D	5902	0	5922	29	0
2	В	2843	0	2852	10	0
2	Е	2854	0	2864	15	0
3	С	1352	0	1370	9	0
3	F	1343	0	1368	3	0
4	А	42	0	56	3	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	В	6	0	8	0	0
4	С	18	0	24	0	0
4	D	6	0	8	1	0
4	Е	6	0	8	0	0
4	F	12	0	16	0	0
5	А	32	0	48	0	0
5	В	4	0	6	0	0
5	D	4	0	6	0	0
5	F	4	0	6	0	0
6	А	32	0	0	1	0
6	В	24	0	0	0	0
6	D	24	0	0	1	0
6	Ε	24	0	0	0	0
7	А	9	0	0	1	0
7	D	9	0	0	0	0
8	А	1	0	0	0	0
8	D	1	0	0	0	0
9	В	53	0	31	0	0
9	Е	53	0	31	1	0
10	D	6	0	9	0	0
11	D	1	0	0	0	0
12	А	455	0	0	2	0
12	В	140	0	0	0	0
12	С	160	0	0	1	0
12	D	351	0	0	0	0
12	Е	179	0	0	2	0
12	F	133	0	0	0	0
All	All	21942	0	20503	91	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 91 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:A:812:XCC:FE4	7:A:812:XCC:S2	1.86	0.66
1:D:292:LEU:O	1:D:296[B]:ARG:HG3	2.04	0.57
3:C:41:GLU:O	3:C:44:GLU:HG2	2.10	0.51
1:A:51:TYR:CD2	4:A:813:GOL:H31	2.45	0.51
1:A:83:LYS:HA	1:A:89:ASP:HA	1.96	0.47



8RIU

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	757/792~(96%)	739~(98%)	17 (2%)	1 (0%)	48	41
1	D	762/792~(96%)	743~(98%)	18 (2%)	1 (0%)	48	41
2	В	368/370~(100%)	359~(98%)	9(2%)	0	100	100
2	Ε	369/370~(100%)	362~(98%)	7 (2%)	0	100	100
3	С	174/174~(100%)	172 (99%)	2(1%)	0	100	100
3	F	173/174~(99%)	172 (99%)	1 (1%)	0	100	100
All	All	2603/2672~(97%)	2547 (98%)	54 (2%)	2(0%)	48	41

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	А	308	GLN
1	D	308	GLN

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentil	\mathbf{es}
1	А	628/657~(96%)	621~(99%)	7 (1%)	70 71]
1	D	633/657~(96%)	625~(99%)	8 (1%)	65 65	
2	В	308/308~(100%)	308 (100%)	0	100 10	0



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
2	Ε	309/308~(100%)	308 (100%)	1 (0%)	91 92
3	С	143/141~(101%)	141 (99%)	2(1%)	62 62
3	F	142/141~(101%)	139~(98%)	3~(2%)	48 45
All	All	2163/2212 (98%)	2142~(99%)	21 (1%)	73 74

Continued from previous page...

5 of 21 residues with a non-rotameric sidechain are listed below:

Mol	Chain	\mathbf{Res}	Type
1	D	405	ASP
2	Ε	256	ARG
3	F	124	ILE
3	F	86	HIS
1	D	479	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

Of 47 ligands modelled in this entry, 3 are monoatomic - leaving 44 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the



Mol	Tuno	Chain	Dog	Link	Bo	ond leng	gths Bond angles			les
WIOI	туре	Ullalli	nes	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2
4	GOL	A	814	-	5,5,5	1.01	0	5,5,5	0.85	0
5	EDO	F	402	-	3,3,3	0.13	0	2,2,2	0.25	0
5	EDO	А	818	-	3,3,3	0.06	0	2,2,2	0.23	0
4	GOL	А	801	-	$5,\!5,\!5$	0.86	0	$5,\!5,\!5$	1.02	0
5	EDO	А	803	-	3,3,3	0.09	0	2,2,2	0.18	0
6	SF4	А	809	1	0,12,12	-	-	-		
7	XCC	D	806	1	0,11,11	-	-	-		
4	GOL	А	815	-	$5,\!5,\!5$	0.10	0	$5,\!5,\!5$	0.35	0
6	SF4	В	405	2	0,12,12	-	-	-		
4	GOL	С	203	-	$5,\!5,\!5$	0.94	0	$5,\!5,\!5$	0.88	0
4	GOL	С	201	-	$5,\!5,\!5$	0.08	0	$5,\!5,\!5$	0.38	0
6	SF4	D	803	1	0,12,12	-	-	-		
4	GOL	С	202	-	5,5,5	0.82	0	$5,\!5,\!5$	0.97	0
6	SF4	D	804	1	0,12,12	-	-	-		
9	FAD	В	404	-	53,58,58	0.64	0	68,89,89	0.68	1 (1%)
4	GOL	А	816	-	$5,\!5,\!5$	0.09	0	$5,\!5,\!5$	0.35	0
4	GOL	Е	405	-	$5,\!5,\!5$	0.92	0	$5,\!5,\!5$	0.91	0
6	SF4	Е	401	2	0,12,12	-	_	-		
5	EDO	А	808	-	3,3,3	0.08	0	2,2,2	0.23	0
4	GOL	F	401	-	5,5,5	0.08	0	$5,\!5,\!5$	0.34	0
5	EDO	А	802	-	3,3,3	0.10	0	2,2,2	0.10	0
4	GOL	А	813	-	5,5,5	0.08	0	$5,\!5,\!5$	0.33	0
4	GOL	А	820	-	$5,\!5,\!5$	0.08	0	$5,\!5,\!5$	0.41	0
4	GOL	А	817	-	5,5,5	0.95	0	$5,\!5,\!5$	0.95	0
9	FAD	Е	403	-	53,58,58	0.66	0	68,89,89	0.68	1 (1%)
5	EDO	А	806	-	3,3,3	0.07	0	2,2,2	0.16	0
6	SF4	А	811	1	0,12,12	-	-	-		
4	GOL	F	403	-	5,5,5	0.99	0	$5,\!5,\!5$	0.84	0
6	SF4	В	403	2	0,12,12	-	-	-		
6	SF4	D	805	1	0,12,12	-	-	-		
5	EDO	А	804	-	3,3,3	0.12	0	2,2,2	0.15	0
6	SF4	В	402	2	0,12,12	-	-	-		
6	SF4	Е	402	2	0,12,12	-	-	-		
6	SF4	Е	404	2	0,12,12	-	-	-		
7	XCC	А	812	1	0,11,11	-	-	-		
6	SF4	А	819	1	0,12,12	-	-	-		
10	PE4	D	807	-	5,5,23	0.34	0	4,4,22	0.41	0
5	EDO	В	401	-	3,3,3	0.11	0	2,2,2	0.21	0
5	EDO	D	802	-	3,3,3	0.06	0	2,2,2	0.20	0
6	SF4	A	810	1	0,12,12	-	-	-		

expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mal	Turne	Chain	Dec	Tink	Bond lengths Bond angles			les		
WIOI	туре	Unain	nes	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2
5	EDO	A	807	-	$3,\!3,\!3$	0.14	0	2,2,2	0.27	0
4	GOL	В	406	-	$5,\!5,\!5$	0.08	0	$5,\!5,\!5$	0.36	0
4	GOL	D	801	-	$5,\!5,\!5$	0.09	0	$5,\!5,\!5$	0.32	0
5	EDO	A	805	-	3,3,3	0.16	0	2,2,2	0.23	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	\mathbf{Res}	Link	Chirals	Torsions	Rings
4	GOL	А	814	-	-	0/4/4/4	-
5	EDO	F	402	-	-	0/1/1/1	-
5	EDO	А	818	-	-	1/1/1/1	-
4	GOL	А	801	-	-	0/4/4/4	-
5	EDO	А	803	-	-	1/1/1/1	-
6	SF4	А	809	1	-	-	0/6/5/5
7	XCC	D	806	1	-	-	0/3/3/3
4	GOL	А	815	-	-	2/4/4/4	-
6	SF4	В	405	2	-	-	0/6/5/5
4	GOL	С	203	-	-	0/4/4/4	-
4	GOL	С	201	-	-	2/4/4/4	-
9	FAD	В	404	-	-	1/30/50/50	0/6/6/6
4	GOL	С	202	-	-	2/4/4/4	-
6	SF4	D	803	1	-	-	0/6/5/5
6	SF4	D	804	1	-	-	0/6/5/5
4	GOL	А	816	-	-	3/4/4/4	-
4	GOL	Е	405	-	-	3/4/4/4	-
6	SF4	Е	401	2	-	-	0/6/5/5
5	EDO	А	808	-	-	0/1/1/1	-
4	GOL	F	401	-	-	0/4/4/4	-
5	EDO	А	802	-	-	1/1/1/1	-
4	GOL	А	813	-	-	4/4/4/4	-
4	GOL	А	820	-	-	3/4/4/4	-
4	GOL	А	817	-	-	3/4/4/4	-
9	FAD	Е	403	-	-	1/30/50/50	0/6/6/6
5	EDO	А	806	-	-	1/1/1/1	-
6	SF4	А	811	1	-	-	0/6/5/5
4	GOL	F	403	-	-	3/4/4/4	-
6	SF4	В	403	2	-	-	0/6/5/5
6	SF4	D	805	1	-	-	0/6/5/5



8RIU

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings				
5	EDO	А	804	-	-	0/1/1/1	-				
6	SF4	В	402	2	-	-	0/6/5/5				
6	SF4	Е	402	2	-	-	0/6/5/5				
6	SF4	Е	404	2	-	-	0/6/5/5				
7	XCC	А	812	1	-	-	0/3/3/3				
6	SF4	А	819	1	-	-	0/6/5/5				
10	PE4	D	807	-	-	2/3/3/21	-				
5	EDO	В	401	-	-	0/1/1/1	-				
5	EDO	D	802	-	-	1/1/1/1	-				
6	SF4	А	810	1	-	-	0/6/5/5				
5	EDO	А	807	-	-	0/1/1/1	-				
4	GOL	В	406	-	-	2/4/4/4	-				
4	GOL	D	801	-	-	3/4/4/4	-				
5	EDO	А	805	-	-	0/1/1/1	-				

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
9	Е	403	FAD	C5A-C6A-N6A	2.33	123.90	120.35
9	В	404	FAD	C5A-C6A-N6A	2.29	123.84	120.35

There are no chirality outliers.

5 of 39 torsion outliers are listed below:

Mol	Chain	\mathbf{Res}	Type	Atoms
4	А	813	GOL	C1-C2-C3-O3
4	А	816	GOL	C1-C2-C3-O3
4	А	817	GOL	O1-C1-C2-C3
4	А	820	GOL	C1-C2-C3-O3
4	С	201	GOL	O1-C1-C2-C3

There are no ring outliers.

8 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	D	804	SF4	1	0
4	А	816	GOL	1	0
4	А	813	GOL	1	0
4	А	820	GOL	1	0



	5	1	1 5		
Mol	Chain	Res	Type	Clashes	Symm-Clashes
9	Ε	403	FAD	1	0
7	А	812	XCC	1	0
6	А	810	SF4	1	0
4	D	801	GOL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and sufficient that rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.













































































5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle RSRZ \rangle$	#RSRZ>2	$OWAB(Å^2)$	Q < 0.9
1	А	759/792~(95%)	-0.12	3 (0%) 89 90	28, 38, 56, 90	0
1	D	760/792~(95%)	0.15	3 (0%) 89 90	16, 43, 62, 92	4 (0%)
2	В	370/370~(100%)	0.48	8 (2%) 62 64	31, 53, 84, 114	0
2	Е	370/370~(100%)	0.09	0 100 100	21, 44, 70, 104	1 (0%)
3	С	174/174~(100%)	-0.28	0 100 100	22, 34, 45, 52	2 (1%)
3	F	174/174~(100%)	-0.19	1 (0%) 85 87	26, 38, 49, 62	1 (0%)
All	All	2607/2672~(97%)	0.06	15 (0%) 85 87	16, 41, 67, 114	8 (0%)

The worst 5 of 15 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	В	255	VAL	3.1
1	D	127	ARG	3.0
2	В	245	GLY	2.9
2	В	260	ILE	2.6
1	D	137	ILE	2.6

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.



6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(Å^2)$	Q<0.9
5	EDO	А	808	4/4	0.57	0.22	$51,\!58,\!59,\!63$	0
4	GOL	А	815	6/6	0.63	0.20	55,57,57,58	0
5	EDO	D	802	4/4	0.65	0.20	66,69,69,70	0
5	EDO	В	401	4/4	0.68	0.20	$51,\!53,\!55,\!57$	0
4	GOL	С	201	6/6	0.69	0.17	$50,\!57,\!62,\!65$	0
5	EDO	А	818	4/4	0.70	0.18	$53,\!55,\!57,\!59$	0
5	EDO	А	805	4/4	0.72	0.15	$57,\!64,\!67,\!67$	0
4	GOL	D	801	6/6	0.73	0.13	$50,\!55,\!57,\!62$	0
4	GOL	А	814	6/6	0.73	0.17	$51,\!55,\!57,\!58$	0
5	EDO	А	806	4/4	0.75	0.14	$63,\!66,\!67,\!67$	0
4	GOL	Е	405	6/6	0.76	0.14	47,50,53,58	0
5	EDO	А	803	4/4	0.77	0.15	$52,\!53,\!58,\!58$	0
5	EDO	А	807	4/4	0.77	0.16	$59,\!59,\!60,\!63$	0
4	GOL	F	403	6/6	0.79	0.15	$47,\!53,\!54,\!57$	0
4	GOL	А	816	6/6	0.79	0.13	52,57,60,65	0
4	GOL	А	801	6/6	0.79	0.13	$60,\!62,\!66,\!67$	0
10	PE4	D	807	6/24	0.79	0.17	$51,\!54,\!57,\!59$	0
5	EDO	А	802	4/4	0.81	0.13	49,49,52,56	0
4	GOL	А	817	6/6	0.82	0.14	57,62,65,67	0
4	GOL	С	202	6/6	0.83	0.13	44,54,59,60	0
4	GOL	С	203	6/6	0.84	0.12	40,52,55,55	0
4	GOL	F	401	6/6	0.85	0.13	44,49,50,53	0
11	CL	D	808[A]	1/1	0.85	0.36	55, 55, 55, 55	1
5	EDO	F	402	4/4	0.87	0.10	52,54,55,60	0
5	EDO	А	804	4/4	0.87	0.12	49,51,51,57	0
4	GOL	В	406	6/6	0.87	0.11	44,52,56,58	0
4	GOL	А	813	6/6	0.88	0.10	38,39,44,46	0
9	FAD	В	404	53/53	0.93	0.09	40,47,55,59	0
4	GOL	А	820	6/6	0.94	0.07	38,40,41,42	0
7	XCC	D	806	9/9	0.95	0.06	32,36,41,42	0
7	XCC	А	812	9/9	0.96	0.05	31,34,38,40	0
9	FAD	E	403	$\overline{53/53}$	0.96	0.06	33,37,42,44	0
8	Κ	А	821	1/1	0.97	0.09	35,35,35,35	0
6	SF4	В	405	8/8	0.98	0.04	36,38,39,40	0
8	Κ	D	809	1/1	0.98	0.08	36,36,36,36	0
6	SF4	D	805	8/8	0.98	0.03	31,32,34,34	0
6	SF4	Е	401	8/8	0.99	0.03	30,32,33,34	0



Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	SF4	Е	402	8/8	0.99	0.02	29,31,31,33	0
6	SF4	Е	404	8/8	0.99	0.03	33,34,35,37	0
6	SF4	А	811	8/8	0.99	0.03	28,30,31,31	0
6	SF4	А	819	8/8	0.99	0.03	28,29,30,30	0
6	SF4	В	402	8/8	0.99	0.03	30,33,34,34	0
6	SF4	В	403	8/8	0.99	0.03	31,31,32,33	0
6	SF4	А	809	8/8	0.99	0.03	28,29,31,32	0
6	SF4	D	803	8/8	0.99	0.03	26,28,29,30	0
6	SF4	D	804	8/8	0.99	0.03	30,31,32,33	0
6	SF4	А	810	8/8	0.99	0.02	27,29,29,30	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







































































6.5 Other polymers (i)

There are no such residues in this entry.

