

wwPDB X-ray Structure Validation Summary Report (i)

Dec 5, 2024 – 06:14 pm GMT

PDB ID : 8RLY

Title: E. coli endonuclease IV complexed with sulfate, catalytic Fe2+

Authors: Saper, M.A.; Paterson, N.G.; Kirillov, S.; Rouvinski, A.

Deposited on : 2024-01-04

Resolution : 1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.4, CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 3.0

buster-report : 1.1.7 (2018)

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

CCP4 : 9.0.003 (Gargrove)

Density-Fitness : 1.0.11

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.40

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	Similar resolution $(\#\text{Entries, resolution range}(\mathring{\mathbf{A}}))$
R_{free}	164625	7293 (1.90-1.90)
Clashscore	180529	8090 (1.90-1.90)
Ramachandran outliers	177936	8022 (1.90-1.90)
Sidechain outliers	177891	8022 (1.90-1.90)
RSRZ outliers	164620	7292 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	285	95%	5%
1	В	285	96%	
1	С	285	95%	
1	D	285	93%	7%
1	Е	285	96%	•



2 Entry composition (i)

There are 8 unique types of molecules in this entry. The entry contains 22932 atoms, of which 10736 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Endonuclease 4.

Mol	Chain	Residues			Atom	ıs			ZeroOcc	AltConf	Trace
1	A	284	Total	С	Н	N	О	S	0	2	0
1	Λ	204	4373	1393	2159	389	420	12	U	J.	0
1	В	279	Total	\mathbf{C}	Η	N	O	\mathbf{S}	0	3	0
1	D	219	4294	1368	2118	383	414	11	U	5	U
1	C	280	Total	С	Η	N	О	S	0	1	0
1		200	4298	1371	2117	384	415	11	O	1	U
1	D	285	Total	С	Η	N	O	S	0	3	0
1	D	200	4386	1395	2165	390	424	12	0	5	U
1	Е	285	Total	С	Н	N	О	S	0	2	0
1	ינו	200	4377	1394	2160	390	422	11		<u> </u>	U

• Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	2	Total Zn 2 2	0	0
2	В	2	Total Zn 2 2	0	0
2	С	2	Total Zn 2 2	0	0
2	D	2	$\begin{array}{ccc} \text{Total} & \text{Zn} \\ 2 & 2 \end{array}$	0	0
2	Е	2	Total Zn 2 2	0	0

• Molecule 3 is FE (II) ION (three-letter code: FE2) (formula: Fe) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Fe 1 1	0	0


Continued on next page...



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	В	1	Total Fe 1 1	0	0
3	С	1	Total Fe 1 1	0	0
3	D	1	Total Fe 1 1	0	0
3	E	1	Total Fe 1 1	0	0

• Molecule 4 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (three-letter code: EPE) (formula: $C_8H_{18}N_2O_4S$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf		
4	A	1	Total 32		H 17		O 4	S 1	0	0

• Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O₄S) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total O S 5 4 1	0	0
5	В	1	Total O S 5 4 1	0	0
5	С	1	Total O S 5 4 1	0	0
5	D	1	Total O S 5 4 1	0	0
5	Е	1	Total O S 5 4 1	0	0

• Molecule 6 is NICKEL (II) ION (three-letter code: NI) (formula: Ni) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total Ni 1 1	0	0
6	В	2	Total Ni 2 2	0	0
6	С	2	Total Ni 2 2	0	0
6	D	1	Total Ni 1 1	0	0
6	Ε	1	Total Ni 1 1	0	0

• Molecule 7 is CHLORIDE ION (three-letter code: CL) (formula: Cl).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	2	Total Cl 2 2	0	0
7	В	1	Total Cl 1 1	0	0
7	С	1	Total Cl 1 1	0	0
7	Ε	1	Total Cl 1 1	0	0

• Molecule 8 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	275	Total O 276 276	0	1
8	В	253	Total O 254 254	0	1
8	С	242	Total O 242 242	0	0
8	D	159	Total O 159 159	0	0
8	E	189	Total O 189 189	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Endonuclease 4





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	69.78Å 115.98Å 177.70Å	Donositon
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	97.12 - 1.90	Depositor
Resolution (A)	97.12 - 1.90	EDS
% Data completeness	87.8 (97.12-1.90)	Depositor
(in resolution range)	87.6 (97.12-1.90)	EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.33 (at 1.87Å)	Xtriage
Refinement program	PHENIX 1.21_5190	Depositor
D D.	0.157 , 0.207	Depositor
R, R_{free}	0.156 , 0.205	DCC
R_{free} test set	51039 reflections (50.22%)	wwPDB-VP
Wilson B-factor (Å ²)	22.6	Xtriage
Anisotropy	0.185	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.38, 35.8	EDS
L-test for twinning ²	$ < L >=0.51, < L^2>=0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	22932	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 2.93% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, FE2, NI, EPE, SO4, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
MIOI	Chain	RMSZ $ \# Z > 5$		RMSZ	# Z > 5	
1	A	0.51	0/2272	0.67	0/3074	
1	В	0.50	0/2239	0.65	0/3030	
1	С	0.50	0/2232	0.65	0/3021	
1	D	0.43	0/2282	0.61	0/3086	
1	Е	0.47	0/2275	0.62	0/3079	
All	All	0.48	0/11300	0.64	0/15290	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a maintain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	125	ARG	Sidechain
1	A	213	ARG	Sidechain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within



the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2214	2159	2151	9	2
1	В	2176	2118	2108	6	0
1	С	2181	2117	2115	5	0
1	D	2221	2165	2159	12	1
1	Ε	2217	2160	2152	5	1
2	A	2	0	0	0	0
2	В	2	0	0	0	0
2	С	2	0	0	0	0
2	D	2	0	0	0	0
2	Ε	2	0	0	0	0
3	A	1	0	0	0	0
3	В	1	0	0	0	0
3	С	1	0	0	0	0
3	D	1	0	0	0	0
3	Ε	1	0	0	0	0
4	A	15	17	17	0	0
5	A	5	0	0	0	0
5	В	5	0	0	0	0
5	С	5	0	0	0	0
5	D	5	0	0	1	0
5	${ m E}$	5	0	0	0	0
6	A	1	0	0	0	0
6	В	2	0	0	0	1
6	С	2	0	0	0	1
6	D	1	0	0	0	0
6	Ε	1	0	0	0	1
7	A	2	0	0	0	0
7	В	1	0	0	0	0
7	С	1	0	0	0	0
7	Ε	1	0	0	0	0
8	A	276	0	0	5	1
8	В	254	0	0	4	1
8	С	242	0	0	1	3
8	D	159	0	0	2	0
8	Ε	189	0	0	2	0
All	All	12196	10736	10702	36	7

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 36 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	$egin{array}{c} ext{Interatomic} \ ext{distance } (ext{Å}) \end{array}$	Clash overlap (Å)
1:B:152:SER:OG	8:B:401:HOH:O	1.72	1.08
1:E:23:GLU:OE2	8:E:401:HOH:O	1.82	0.96
1:B:198:LYS:NZ	8:B:403:HOH:O	2.16	0.77
1:B:40:ARG:NH2	8:B:402:HOH:O	1.97	0.74
1:C:168:VAL:O	1:C:171:LYS:NZ	2.30	0.64

The worst 5 of 7 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$egin{array}{ll} ext{Interatomic} \ ext{distance } (ext{Å}) \end{array}$	Clash overlap (Å)
8:A:615:HOH:O	8:C:601:HOH:O[4_555]	1.82	0.38
1:A:241:HIS:HE2	6:C:306:NI:NI[4_555]	1.23	0.37
8:C:545:HOH:O	8:C:622:HOH:O[4_545]	1.88	0.32
1:E:241:HIS:HE2	6:B:306:NI:NI[1_455]	1.29	0.31
1:D:60:HIS:HE2	6:E:305:NI:NI[4_545]	1.38	0.22

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	$284/285\ (100\%)$	282 (99%)	2 (1%)	0	100 100
1	В	280/285~(98%)	280 (100%)	0	0	100 100
1	C	$279/285\ (98\%)$	279 (100%)	0	0	100 100
1	D	$286/285\ (100\%)$	281 (98%)	5 (2%)	0	100 100
1	E	$285/285\ (100\%)$	284 (100%)	1 (0%)	0	100 100
All	All	$1414/1425 \ (99\%)$	1406 (99%)	8 (1%)	0	100 100

There are no Ramachandran outliers to report.



5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	A	$229/227 \ (101\%)$	227 (99%)	2 (1%)	75	77
1	В	226/227 (100%)	224 (99%)	2 (1%)	75	77
1	С	$225/227\ (99\%)$	225 (100%)	0	100	100
1	D	230/227 (101%)	229 (100%)	1 (0%)	89	90
1	E	229/227 (101%)	226 (99%)	3 (1%)	65	65
All	All	1139/1135 (100%)	1131 (99%)	8 (1%)	81	83

5 of 8 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	Е	232	HIS
1	Е	166	ASP
1	D	1	MET
1	В	222	SER
1	Е	1	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.



5.6 Ligand geometry (i)

Of 33 ligands modelled in this entry, 27 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trmo	Chain	Dag	Link	Во	ond leng	ths	В	ond ang	les
MIOI	Type	Chain	Res	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	EPE	A	304	-	15,15,15	0.82	0	18,20,20	1.62	1 (5%)
5	SO4	D	304	3,2	4,4,4	0.49	0	6,6,6	0.40	0
5	SO4	Е	304	3,2	4,4,4	0.69	0	6,6,6	0.65	0
5	SO4	В	304	3,2	4,4,4	0.75	0	6,6,6	0.45	0
5	SO4	С	304	3,2	4,4,4	0.77	0	6,6,6	0.46	0
5	SO4	A	305	3,2	4,4,4	0.55	0	6,6,6	0.38	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EPE	A	304	-	-	2/9/19/19	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
4	A	304	EPE	O1S-S-C10	-5.59	100.19	106.92

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	304	EPE	C8-C7-N4-C5
4	A	304	EPE	N4-C7-C8-O8

There are no ring outliers.




1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	D	304	SO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.
























5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\# \mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q<0.9
1	A	$284/285 \ (99\%)$	-0.76	1 (0%) 89 90	12, 21, 33, 49	2 (0%)
1	В	279/285 (97%)	-0.74	0 100 100	15, 21, 33, 50	3 (1%)
1	С	280/285~(98%)	-0.76	1 (0%) 89 90	15, 21, 32, 64	1 (0%)
1	D	285/285 (100%)	-0.24	9 (3%) 50 52	16, 29, 47, 81	3 (1%)
1	E	285/285 (100%)	-0.50	3 (1%) 77 79	17, 24, 42, 91	2 (0%)
All	All	1413/1425 (99%)	-0.60	14 (0%) 79 81	12, 23, 40, 91	11 (0%)

The worst 5 of 14 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	284	VAL	7.0
1	D	285	ALA	6.3
1	D	283	ALA	6.0
1	С	280	THR	4.8
1	Е	285	ALA	4.3

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,




median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
5	SO4	A	305	5/5	0.95	0.13	23,27,31,31	0
6	NI	С	305	1/1	0.95	0.07	59,59,59,59	0
4	EPE	A	304	15/15	0.96	0.08	22,39,51,52	0
5	SO4	С	304	5/5	0.97	0.12	21,26,28,30	0
5	SO4	В	304	5/5	0.97	0.11	21,22,27,27	0
5	SO4	Ε	304	5/5	0.98	0.07	18,27,30,31	0
6	NI	В	305	1/1	0.98	0.05	47,47,47,47	0
5	SO4	D	304	5/5	0.98	0.08	25,29,31,32	0
6	NI	D	305	1/1	0.98	0.08	58,58,58,58	0
2	ZN	D	303	1/1	0.99	0.04	25,25,25,25	0
6	NI	Ε	305	1/1	0.99	0.05	30,30,30,30	0
7	CL	A	307	1/1	0.99	0.04	18,18,18,18	0
7	CL	A	308	1/1	0.99	0.03	23,23,23,23	0
7	CL	С	307	1/1	0.99	0.02	19,19,19,19	0
7	CL	E	306	1/1	0.99	0.06	27,27,27,27	0
2	ZN	A	303	1/1	1.00	0.02	19,19,19,19	0
2	ZN	В	301	1/1	1.00	0.02	22,22,22,22	0
2	ZN	В	303	1/1	1.00	0.03	19,19,19,19	0
2	ZN	С	301	1/1	1.00	0.02	20,20,20,20	0
2	ZN	С	303	1/1	1.00	0.02	19,19,19,19	0
2	ZN	D	301	1/1	1.00	0.01	24,24,24,24	0
6	NI	A	306	1/1	1.00	0.06	25,25,25,25	0
2	ZN	A	301	1/1	1.00	0.01	20,20,20,20	0
6	NI	В	306	1/1	1.00	0.04	22,22,22,22	0
2	ZN	E	301	1/1	1.00	0.01	22,22,22,22	0
6	NI	С	306	1/1	1.00	0.05	23,23,23,23	0
2	ZN	Ε	303	1/1	1.00	0.03	20,20,20,20	0
3	FE2	A	302	1/1	1.00	0.09	20,20,20,20	0
3	FE2	В	302	1/1	1.00	0.09	21,21,21,21	0
3	FE2	С	302	1/1	1.00	0.09	20,20,20,20	0
7	CL	В	307	1/1	1.00	0.02	24,24,24,24	0
3	FE2	D	302	1/1	1.00	0.09	27,27,27,27	0
3	FE2	Е	302	1/1	1.00	0.11	23,23,23,23	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.










Electron density around NI C 305:


 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)














Electron density around SO4 B 304: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_{o}\text{-}\mathrm{DF}_{c}$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around NI B 305:


 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)

























Electron density around NI E 305: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)




Electron density around ZN A 303:

 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around ZN B 301:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around ZN B 303:

 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)







Electron density around ZN C 301: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_{o}\text{-}\mathrm{DF}_{c}$ (at 3 rmsd) in purple (negative) and green (positive)





Electron density around ZN C 303: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)





Electron density around ZN D 301:

 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)





















Electron density around ZN E 301: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_{o}\text{-}\mathrm{DF}_{c}$ (at 3 rmsd) in purple (negative) and green (positive)





Electron density around NI C 306: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_{o}\text{-}\mathrm{DF}_{c}$ (at 3 rmsd) in purple (negative) and green (positive)





Electron density around ZN E 303:

 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)





























6.5 Other polymers (i)

There are no such residues in this entry.

