



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 1, 2022 – 12:56 PM EDT

PDB ID : 5RM1
Title : PanDDA analysis group deposition – Crystal Structure of SARS-CoV-2 heli-
case in complex with Z426041412
Authors : Newman, J.A.; Yosaatmadja, Y.; Douangamath, A.; Aimon, A.; Powell, A.J.;
Dias, A.; Fearon, D.; Dunnett, L.; Brandao-Neto, J.; Krojer, T.; Skyner, R.;
Gorrie-Stone, T.; Thompson, W.; von Delft, F.; Arrowsmith, C.H.; Edwards,
A.; Bountra, C.; Gileadi, O.
Deposited on : 2020-09-16
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at <http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.29
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.29


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	601	
1	B	601	

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 9418 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Helicase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	572	4417	2816	737	832	32	0	0	0
1	B	585	4508	2875	750	848	35	0	1	0

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	3	Total	Zn	0	0
			3	3		
2	B	3	Total	Zn	0	0
			3	3		

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	P	0	0
			5	4	1		
3	A	1	Total	O	P	0	0
			5	4	1		
3	B	1	Total	O	P	0	0
			5	4	1		
3	B	1	Total	O	P	0	0
			5	4	1		

- Molecule 4 is N-[4-(aminomethyl)phenyl]methanesulfonamide (three-letter code: RY4) (formula: C₈H₁₂N₂O₂S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	B	1	Total	C	N	O	S	0	0
			13	8	2	2	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	204	Total	O	0	0
			204	204		
5	B	250	Total	O	0	0
			250	250		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Helicase



- Molecule 1: Helicase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	59.32Å 70.17Å 85.16Å 102.40° 96.07° 112.75°	Depositor
Resolution (Å)	81.33 – 1.90 81.33 – 1.90	Depositor EDS
% Data completeness (in resolution range)	93.7 (81.33-1.90) 93.7 (81.33-1.90)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.12 (at 1.90Å)	Xtriage
Refinement program	BUSTER 2.10.3 (20-MAY-2020)	Depositor
R, R_{free}	0.176 , 0.234 0.197 , 0.252	Depositor DCC
R_{free} test set	4614 reflections (5.13%)	wwPDB-VP
Wilson B-factor (Å ²)	35.3	Xtriage
Anisotropy	0.290	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 40.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9418	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.77% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, ZN, RY4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.52	0/4517	0.64	1/6156 (0.0%)
1	B	0.52	0/4610	0.65	0/6283
All	All	0.52	0/9127	0.65	1/12439 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	591	GLU	C-N-CA	5.08	134.40	121.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4417	0	4321	35	0
1	B	4508	0	4423	35	0
2	A	3	0	0	0	0
2	B	3	0	0	0	0
3	A	10	0	0	0	0
3	B	10	0	0	0	0
4	B	13	0	0	1	0
5	A	204	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	250	0	0	3	0
All	All	9418	0	8744	68	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:12:THR:CG2	1:B:26:CYS:HA	2.12	0.79
1:A:12:THR:CG2	1:A:26:CYS:HA	2.14	0.78
1:A:255:THR:HG23	1:A:300:PRO:HG3	1.69	0.74
1:B:12:THR:HG21	1:B:26:CYS:HA	1.72	0.71
1:A:12:THR:HG21	1:A:26:CYS:HA	1.73	0.70
1:B:293:ILE:HG13	1:B:320:LYS:HB3	1.76	0.68
1:B:13:SER:HB2	1:B:92:LEU:HB2	1.77	0.66
1:A:352:LEU:HD11	1:B:234:PRO:HD3	1.77	0.66
1:A:293:ILE:HG13	1:A:320:LYS:HB3	1.78	0.65
1:B:12:THR:HG21	1:B:25:LEU:O	1.96	0.65
1:A:12:THR:HG21	1:A:25:LEU:O	1.97	0.65
1:B:474[B]:MET:HG2	1:B:590:LEU:HB2	1.80	0.64
1:A:19:CYS:HB2	1:A:23:PRO:HD2	1.79	0.64
1:B:12:THR:HG22	5:B:811:HOH:O	2.00	0.61
1:A:158:LEU:HD11	1:A:164:HIS:CE1	2.37	0.59
1:A:215:THR:HG22	1:B:193:VAL:HG21	1.87	0.57
1:B:12:THR:HG23	1:B:26:CYS:HA	1.86	0.56
1:A:157:VAL:HG21	1:A:219:LEU:O	2.05	0.56
1:B:269:TYR:CD1	1:B:295:LEU:HD13	2.41	0.56
1:A:269:TYR:CD1	1:A:295:LEU:HD13	2.41	0.55
1:B:504:PRO:HB3	1:B:507:ARG:HH21	1.71	0.55
1:A:280:LEU:HD11	1:A:438:LEU:HG	1.89	0.54
1:B:103:VAL:HG12	1:B:103:VAL:O	2.09	0.53
1:B:156:GLU:HG2	1:B:158:LEU:HD12	1.91	0.53
1:B:271:LYS:NZ	5:B:806:HOH:O	2.40	0.53
1:A:442:ARG:NH1	1:A:464:HIS:CE1	2.77	0.53
1:B:445:PRO:HB3	1:B:468:SER:HB3	1.90	0.53
1:A:12:THR:HG23	1:A:26:CYS:HA	1.91	0.52
1:B:12:THR:CG2	5:B:811:HOH:O	2.58	0.52
1:A:445:PRO:HB3	1:A:468:SER:HB3	1.91	0.51
1:B:16:CYS:O	1:B:22:ARG:HA	2.10	0.51
1:A:7:LEU:HD22	1:A:103:VAL:HG22	1.93	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:6:VAL:HG11	4:B:701:RY4:C1	2.41	0.51
1:B:510:VAL:HG21	1:B:541:TYR:CD1	2.46	0.51
1:A:510:VAL:HG21	1:A:541:TYR:CD1	2.45	0.50
1:A:519:ASN:HB3	1:A:530:THR:CG2	2.40	0.50
1:B:93:TYR:HB3	1:B:96:THR:OG1	2.12	0.50
1:B:249:ILE:HG23	1:B:273:GLY:HA3	1.94	0.50
1:A:82:PRO:HG2	1:A:85:ALA:HB2	1.95	0.49
1:B:280:LEU:HD11	1:B:438:LEU:HG	1.94	0.49
1:A:158:LEU:HD11	1:A:164:HIS:ND1	2.28	0.49
1:A:140:ALA:O	1:A:144:THR:HG23	2.13	0.48
1:B:451:THR:HG21	1:B:585:LEU:HD23	1.96	0.47
1:B:158:LEU:CD1	1:B:164:HIS:ND1	2.77	0.47
1:A:7:LEU:HD11	1:A:106:PHE:CG	2.50	0.47
1:A:519:ASN:HB3	1:A:530:THR:HG23	1.97	0.47
1:B:252:LEU:HB3	1:B:299:TYR:CD1	2.51	0.46
1:A:163:LEU:HG	1:A:211:TYR:CD2	2.51	0.46
1:A:326:PRO:HG2	1:A:329:LYS:NZ	2.32	0.45
1:B:13:SER:O	1:B:44:SER:HA	2.18	0.43
1:A:249:ILE:HG23	1:A:273:GLY:HA3	1.99	0.43
1:A:377:SER:O	1:A:406:PRO:HA	2.19	0.43
1:B:377:SER:O	1:B:406:PRO:HA	2.18	0.43
1:A:13:SER:O	1:A:44:SER:HA	2.18	0.43
1:A:158:LEU:HG	1:A:163:LEU:HA	2.01	0.43
1:A:512:ILE:O	1:A:546:PHE:HA	2.18	0.43
1:B:512:ILE:O	1:B:546:PHE:HA	2.19	0.43
1:B:163:LEU:HD21	1:B:219:LEU:CD1	2.49	0.43
1:A:451:THR:HG21	1:A:585:LEU:HD23	2.01	0.42
1:B:19:CYS:HB2	1:B:23:PRO:HD2	2.01	0.42
1:B:149:TYR:HB3	1:B:174:PRO:HD3	2.02	0.42
1:B:158:LEU:HD11	1:B:164:HIS:ND1	2.33	0.42
1:A:350:SER:HB3	1:A:353:GLU:HG2	2.02	0.41
1:A:252:LEU:HB3	1:A:299:TYR:CD2	2.56	0.41
1:B:376:ILE:HG12	1:B:425:VAL:HG11	2.03	0.41
1:A:304:ILE:HG12	1:A:370:ILE:HB	2.02	0.40
1:A:376:ILE:HG12	1:A:425:VAL:HG11	2.02	0.40
1:B:103:VAL:O	1:B:103:VAL:CG1	2.69	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	564/601 (94%)	550 (98%)	13 (2%)	1 (0%)	47	38
1	B	580/601 (96%)	559 (96%)	20 (3%)	1 (0%)	47	38
All	All	1144/1202 (95%)	1109 (97%)	33 (3%)	2 (0%)	47	38

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	484	VAL
1	B	219	LEU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	485/523 (93%)	466 (96%)	19 (4%)	32	23
1	B	498/523 (95%)	479 (96%)	19 (4%)	33	24
All	All	983/1046 (94%)	945 (96%)	38 (4%)	32	23

All (38) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	12	THR
1	A	20	ILE
1	A	35	ILE
1	A	46	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	81	PHE
1	A	127	THR
1	A	129	ARG
1	A	173	ARG
1	A	207	ASP
1	A	209	VAL
1	A	231	THR
1	A	255	THR
1	A	257	ASN
1	A	373	PHE
1	A	390	ARG
1	A	392	ARG
1	A	530	THR
1	A	531	GLN
1	A	592	ILE
1	B	7	LEU
1	B	12	THR
1	B	51	ASN
1	B	69	SER
1	B	95	ASN
1	B	124	ASN
1	B	156	GLU
1	B	158	LEU
1	B	160	ASP
1	B	164	HIS
1	B	192	LYS
1	B	195	ILE
1	B	220	ASN
1	B	231	THR
1	B	247	VAL
1	B	353	GLU
1	B	373	PHE
1	B	458	ASP
1	B	516	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	86	ASN
1	A	179	ASN
1	A	464	HIS
1	A	531	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	268	ASN
1	B	516	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 6 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	PO4	B	705	-	4,4,4	2.48	1 (25%)	6,6,6	0.43	0
3	PO4	A	704	-	4,4,4	2.49	1 (25%)	6,6,6	0.93	0
3	PO4	A	705	-	4,4,4	2.45	2 (50%)	6,6,6	1.00	0
3	PO4	B	706	-	4,4,4	2.45	1 (25%)	6,6,6	0.23	0
4	RY4	B	701	-	13,13,13	0.35	0	18,18,18	0.48	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	RY4	B	701	-	-	0/7/7/7	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	704	PO4	P-O1	4.15	1.60	1.50
3	B	705	PO4	P-O1	4.05	1.60	1.50
3	B	706	PO4	P-O1	4.03	1.60	1.50
3	A	705	PO4	P-O1	3.94	1.60	1.50
3	A	705	PO4	P-O3	2.06	1.60	1.54

There are no bond angle outliers.

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	701	RY4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	572/601 (95%)	0.06	35 (6%) 21 24	28, 50, 99, 131	0
1	B	585/601 (97%)	-0.01	21 (3%) 42 45	30, 44, 87, 132	0
All	All	1157/1202 (96%)	0.03	56 (4%) 30 33	28, 48, 94, 132	0

All (56) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	217	TYR	9.4
1	A	215	THR	6.6
1	B	340	VAL	6.1
1	B	96	THR	5.9
1	B	193	VAL	5.6
1	A	339	ARG	5.5
1	A	229	SER	5.4
1	A	216	THR	5.2
1	A	169	VAL	4.5
1	A	228	THR	4.5
1	A	170	GLY	4.2
1	B	196	GLY	4.1
1	A	227	LEU	3.9
1	A	256	LEU	3.8
1	A	184	GLY	3.7
1	A	258	ILE	3.7
1	B	69	SER	3.6
1	A	176	LEU	3.5
1	B	218	LYS	3.4
1	A	195	ILE	3.4
1	B	67	GLY	3.2
1	A	214	THR	3.2
1	A	230	HIS	3.2
1	A	217	TYR	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	187	VAL	3.2
1	A	210	VAL	3.0
1	B	53	PRO	2.9
1	B	157	VAL	2.9
1	A	149	TYR	2.8
1	A	336	ALA	2.8
1	A	173	ARG	2.8
1	B	45	VAL	2.7
1	A	337	ARG	2.7
1	A	225	PHE	2.7
1	A	167	TRP	2.7
1	A	81	PHE	2.6
1	B	219	LEU	2.6
1	A	352	LEU	2.5
1	A	194	GLN	2.5
1	B	158	LEU	2.4
1	A	257	ASN	2.4
1	A	340	VAL	2.4
1	B	216	THR	2.4
1	A	219	LEU	2.3
1	A	148	SER	2.3
1	A	338	ALA	2.2
1	A	156	GLU	2.1
1	A	182	PHE	2.1
1	B	81	PHE	2.1
1	A	508	LYS	2.1
1	B	95	ASN	2.1
1	B	65	LEU	2.1
1	A	220	ASN	2.0
1	B	2	VAL	2.0
1	B	92	LEU	2.0
1	B	55	CYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	RY4	B	701	13/13	0.85	0.28	35,35,37,38	13
3	PO4	B	706	5/5	0.97	0.12	55,57,61,64	0
2	ZN	B	703	1/1	0.98	0.06	66,66,66,66	0
3	PO4	A	705	5/5	0.98	0.11	39,41,42,44	0
2	ZN	A	703	1/1	0.99	0.09	58,58,58,58	0
3	PO4	B	705	5/5	0.99	0.09	44,46,47,48	0
2	ZN	B	704	1/1	0.99	0.08	82,82,82,82	0
3	PO4	A	704	5/5	0.99	0.11	42,42,46,47	0
2	ZN	A	701	1/1	1.00	0.10	55,55,55,55	0
2	ZN	B	702	1/1	1.00	0.09	52,52,52,52	0
2	ZN	A	702	1/1	1.00	0.10	50,50,50,50	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.