



Full wwPDB X-ray Structure Validation Report ⓘ

May 21, 2020 – 05:22 am BST

PDB ID : 3RN3
Title : SEGMENTED ANISOTROPIC REFINEMENT OF BOVINE RIBONUCLEASE A BY THE APPLICATION OF THE RIGID-BODY TLS MODEL
Authors : Howlin, B.; Moss, D.S.; Harris, G.W.; Palmer, R.A.
Deposited on : 1991-10-30
Resolution : 1.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	1202 (1.46-1.46)
Ramachandran outliers	138981	1178 (1.46-1.46)
Sidechain outliers	138945	1178 (1.46-1.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	124	 62% 27% 9% .

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 1069 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RIBONUCLEASE A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	124	957	579	173	193	12	0	1	0

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	A	1	5	4	1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	107	107	107	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: RIBONUCLEASE A



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	30.45Å 38.37Å 53.22Å 90.00° 105.96° 90.00°	Depositor
Resolution (Å)	(Not available) – 1.45 6.28 – 1.46	Depositor EDS
% Data completeness (in resolution range)	(Not available) ((Not available)-1.45) 85.6 (6.28-1.46)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.57 (at 1.46Å)	Xtrriage
Refinement program	RESTRAIN	Depositor
R, R_{free}	0.223 , (Not available) 0.278 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	9.5	Xtrriage
Anisotropy	0.190	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.46 , 36.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.050 for h,-k,-h-l	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	1069	wwPDB-VP
Average B, all atoms (Å ²)	0.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 15.30% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.11	4/978 (0.4%)	2.19	37/1319 (2.8%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	69	GLN	CD-OE1	5.18	1.35	1.24
1	A	67	ASN	CG-OD1	5.17	1.35	1.24
1	A	28	GLN	CD-OE1	5.09	1.35	1.24
1	A	101	GLN	CD-OE1	5.04	1.35	1.24

All (37) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	10	ARG	NE-CZ-NH1	20.97	130.78	120.30
1	A	10	ARG	NE-CZ-NH2	-17.51	111.54	120.30
1	A	115	TYR	CB-CG-CD2	-16.12	111.33	121.00
1	A	85	ARG	NE-CZ-NH2	-12.29	114.16	120.30
1	A	92	TYR	CB-CG-CD2	-11.96	113.82	121.00
1	A	33	ARG	NE-CZ-NH1	10.47	125.53	120.30
1	A	25	TYR	CB-CG-CD1	-9.86	115.08	121.00
1	A	97	TYR	CG-CD2-CE2	-8.64	114.39	121.30
1	A	25	TYR	CG-CD1-CE1	-8.44	114.55	121.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	115	TYR	CB-CG-CD1	8.14	125.89	121.00
1	A	97	TYR	CB-CG-CD2	-7.98	116.21	121.00
1	A	97	TYR	O-C-N	7.58	134.82	122.70
1	A	34	ASN	CB-CG-OD1	-6.88	107.83	121.60
1	A	24	ASN	CA-CB-CG	-6.71	98.63	113.40
1	A	38	ASP	CB-CG-OD2	-6.43	112.51	118.30
1	A	34	ASN	OD1-CG-ND2	6.40	136.63	121.90
1	A	62	ASN	CB-CG-ND2	6.30	131.82	116.70
1	A	63	VAL	CG1-CB-CG2	6.28	120.95	110.90
1	A	62	ASN	CB-CG-OD1	-6.16	109.27	121.60
1	A	34	ASN	O-C-N	-6.15	112.85	122.70
1	A	73	TYR	O-C-N	5.92	132.18	122.70
1	A	124	VAL	CA-CB-CG1	5.75	119.52	110.90
1	A	97	TYR	CA-C-N	-5.69	104.67	117.20
1	A	92	TYR	CG-CD2-CE2	-5.68	116.76	121.30
1	A	25	TYR	CZ-CE2-CD2	-5.63	114.73	119.80
1	A	83	ASP	CB-CG-OD2	-5.59	113.27	118.30
1	A	4	ALA	O-C-N	-5.43	114.02	122.70
1	A	42	PRO	CA-N-CD	5.42	119.29	111.70
1	A	12	HIS	CA-C-O	-5.34	108.89	120.10
1	A	104	LYS	O-C-N	5.28	131.15	122.70
1	A	34	ASN	CA-CB-CG	-5.27	101.81	113.40
1	A	116	VAL	CA-CB-CG2	5.21	118.72	110.90
1	A	77	SER	O-C-N	5.16	130.95	122.70
1	A	60	GLN	CA-CB-CG	-5.09	102.19	113.40
1	A	121	ASP	CB-CG-OD1	-5.04	113.76	118.30
1	A	117	PRO	CA-N-CD	5.02	118.73	111.70
1	A	39	ARG	CD-NE-CZ	-5.02	116.58	123.60

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	39	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	957	0	908	35	0
2	A	5	0	0	1	0
3	A	107	0	0	10	1
All	All	1069	0	908	35	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:85:ARG:NH1	1:A:85:ARG:HB2	1.82	0.93
1:A:39:ARG:HH11	1:A:39:ARG:HG2	1.40	0.87
1:A:27:ASN:HD21	1:A:97:TYR:H	1.22	0.85
1:A:66:LYS:NZ	3:A:230:HOH:O	2.10	0.83
1:A:85:ARG:HB2	1:A:85:ARG:CZ	2.20	0.72
1:A:66:LYS:NZ	1:A:66:LYS:HB3	2.09	0.68
1:A:27:ASN:ND2	1:A:97:TYR:H	1.92	0.67
1:A:66:LYS:NZ	1:A:66:LYS:CB	2.58	0.66
1:A:34:ASN:OD1	1:A:37:LYS:HG3	1.96	0.65
1:A:39:ARG:HH11	1:A:39:ARG:CG	2.09	0.65
1:A:66:LYS:HZ3	1:A:66:LYS:HB3	1.60	0.65
1:A:119[B]:HIS:HE1	3:A:203:HOH:O	1.78	0.64
1:A:119[B]:HIS:CE1	3:A:203:HOH:O	2.50	0.64
1:A:85:ARG:HH11	1:A:85:ARG:HB2	1.61	0.63
1:A:85:ARG:HH11	1:A:85:ARG:CB	2.13	0.62
1:A:43:VAL:HG12	3:A:211:HOH:O	2.01	0.60
1:A:85:ARG:NH1	1:A:85:ARG:CB	2.62	0.60
1:A:67:ASN:CB	3:A:263:HOH:O	2.52	0.56
1:A:11:GLN:HG2	1:A:35:LEU:HD21	1.88	0.55
1:A:60:GLN:HG2	1:A:76:TYR:CE2	2.42	0.54
1:A:41:LYS:NZ	3:A:282:HOH:O	2.20	0.53
1:A:66:LYS:HG2	1:A:67:ASN:N	2.22	0.53
1:A:60:GLN:HG2	1:A:76:TYR:CD2	2.44	0.53
1:A:20:ALA:H	1:A:101:GLN:NE2	2.09	0.50
1:A:43:VAL:HG12	1:A:44:ASN:N	2.27	0.49
1:A:39:ARG:NH1	1:A:39:ARG:CG	2.71	0.49
1:A:67:ASN:HB3	3:A:263:HOH:O	2.10	0.49
1:A:114:PRO:HA	3:A:340:HOH:O	2.13	0.47
1:A:66:LYS:HB2	3:A:230:HOH:O	2.15	0.46
1:A:66:LYS:HZ3	1:A:66:LYS:CB	2.25	0.46
1:A:25:TYR:CZ	1:A:29:MET:HG3	2.53	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:27:ASN:HD21	1:A:97:TYR:N	2.03	0.43
1:A:72:CYS:HA	1:A:108:VAL:O	2.19	0.42
1:A:77:SER:HB2	3:A:299:HOH:O	2.20	0.41
1:A:119[A]:HIS:ND1	2:A:155:SO4:O4	2.53	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:205:HOH:O	3:A:231:HOH:O[1_455]	1.80	0.40

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	123/124 (99%)	119 (97%)	4 (3%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	110/109 (101%)	101 (92%)	9 (8%)	11 1

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	LYS
1	A	21	SER
1	A	28	GLN
1	A	34	ASN
1	A	37	LYS
1	A	39	ARG
1	A	66	LYS
1	A	83	ASP
1	A	91	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	27	ASN
1	A	69	GLN
1	A	101	GLN
1	A	103	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	155	-	4,4,4	0.53	0	6,6,6	0.81	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	155	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers

Unable to reproduce the depositors R factor - this section is therefore empty.