



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 16, 2024 – 04:19 pm BST

PDB ID : 8RSN
Title : macrodomain-fused SirTM (Mfs1) from *Fusarium oxysporum* f. sp. *cubense* race 1
Authors : Ariza, A.
Deposited on : 2024-01-24
Resolution : 2.22 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.002 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.38.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.22 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	7167 (2.24-2.20)
Clashscore	180529	8096 (2.24-2.20)
Ramachandran outliers	177936	8010 (2.24-2.20)
Sidechain outliers	177891	8011 (2.24-2.20)
RSRZ outliers	164620	7166 (2.24-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	599	
1	B	599	

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 8664 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ADP-ribose 1'-phosphate phosphatase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	538	Total	C	N	O	S	0	2	0
			4202	2655	732	793	22			
1	B	532	Total	C	N	O	S	0	2	0
			4147	2625	717	783	22			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	expression tag	UNP A0A559KX76
A	0	PRO	-	expression tag	UNP A0A559KX76
A	401	THR	ILE	variant	UNP A0A559KX76
A	419	ALA	ASN	engineered mutation	UNP A0A559KX76
B	-1	GLY	-	expression tag	UNP A0A559KX76
B	0	PRO	-	expression tag	UNP A0A559KX76
B	401	THR	ILE	variant	UNP A0A559KX76
B	419	ALA	ASN	engineered mutation	UNP A0A559KX76

- Molecule 2 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0

- Molecule 3 is THIOCYANATE ION (three-letter code: SCN) (formula: CNS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	S	0	0
			3	1	1	1		
3	A	1	Total	C	N	S	0	0
			3	1	1	1		
3	A	1	Total	C	N	S	0	0
			3	1	1	1		
3	A	1	Total	C	N	S	0	0
			3	1	1	1		
3	B	1	Total	C	N	S	0	0
			3	1	1	1		
3	B	1	Total	C	N	S	0	0
			3	1	1	1		
3	B	1	Total	C	N	S	0	0
			3	1	1	1		
3	B	1	Total	C	N	S	0	0
			3	1	1	1		

- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Zn	0	0
			1	1		
4	B	1	Total	Zn	0	0
			1	1		

- Molecule 5 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total K 1 1	0	0
5	B	1	Total K 1 1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	137	Total O 137 137	0	0
6	B	115	Total O 115 115	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: ADP-ribose 1''-phosphate phosphatase



- Molecule 1: ADP-ribose 1''-phosphate phosphatase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	61.74Å 135.09Å 147.62Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	99.86 – 2.22 99.66 – 2.22	Depositor EDS
% Data completeness (in resolution range)	99.8 (99.86-2.22) 99.4 (99.66-2.22)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.40 (at 2.22Å)	Xtrriage
Refinement program	REFMAC 5.8.0415	Depositor
R, R_{free}	0.203 , 0.253 0.205 , 0.251	Depositor DCC
R_{free} test set	3033 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å ²)	38.4	Xtrriage
Anisotropy	0.336	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 39.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8664	wwPDB-VP
Average B, all atoms (Å ²)	57.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.80% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, ZN, K, SCN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.56	3/4292 (0.1%)	0.88	1/5830 (0.0%)
1	B	0.54	2/4242 (0.0%)	0.87	9/5762 (0.2%)
All	All	0.55	5/8534 (0.1%)	0.87	10/11592 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3
1	B	0	6
All	All	0	9

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	574	GLU	CD-OE2	6.51	1.32	1.25
1	B	492	MET	CG-SD	-6.09	1.65	1.81
1	B	377	GLU	CD-OE1	5.60	1.31	1.25
1	A	226	GLU	CD-OE2	-5.48	1.19	1.25
1	A	48	GLU	CD-OE2	5.48	1.31	1.25

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	379	ARG	NE-CZ-NH1	8.38	124.49	120.30
1	B	451	ARG	NE-CZ-NH2	-8.36	116.12	120.30
1	B	492	MET	CG-SD-CE	6.89	111.23	100.20
1	B	379	ARG	NE-CZ-NH2	-6.88	116.86	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	407	ARG	NE-CZ-NH2	-6.73	116.93	120.30
1	B	492	MET	CA-CB-CG	-6.44	102.35	113.30
1	A	520	ARG	NE-CZ-NH1	6.28	123.44	120.30
1	B	486	ARG	CB-CG-CD	5.46	125.80	111.60
1	B	553	ARG	CB-CA-C	-5.11	100.19	110.40
1	B	451	ARG	NE-CZ-NH1	5.01	122.81	120.30

There are no chirality outliers.

All (9) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	170	ARG	Sidechain
1	A	379	ARG	Sidechain
1	A	523	ARG	Sidechain
1	B	158	ARG	Sidechain
1	B	211	ARG	Sidechain
1	B	379	ARG	Sidechain
1	B	486	ARG	Sidechain
1	B	497[A]	ARG	Sidechain
1	B	497[B]	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4202	0	4167	33	0
1	B	4147	0	4118	26	0
2	A	20	0	30	4	0
2	B	12	0	18	1	0
3	A	12	0	0	0	0
3	B	15	0	0	1	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	137	0	0	2	0
6	B	115	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	8664	0	8333	60	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:123:THR:HG21	1:B:311:GLN:HE22	1.27	0.99
1:B:420:ALA:HB1	2:B:607:EDO:H21	1.55	0.85
1:B:44:LEU:HA	1:B:47:LEU:HG	1.72	0.71
6:A:820:HOH:O	3:B:603:SCN:S	2.51	0.68
1:A:63:ARG:CG	1:A:63:ARG:HH11	2.08	0.66
1:A:13:ASP:O	1:A:17:THR:HG23	1.98	0.64
1:A:420:ALA:HB1	2:A:608:EDO:C1	2.29	0.63
1:B:123:THR:HG21	1:B:311:GLN:NE2	2.08	0.63
1:B:66:LEU:HB2	1:B:67:PRO:HD3	1.82	0.61
1:A:66:LEU:HB2	1:A:67:PRO:HD3	1.82	0.60
1:B:13:ASP:O	1:B:17:THR:HG23	2.01	0.59
1:A:498:ALA:HA	2:A:609:EDO:H21	1.85	0.58
1:A:451[A]:ARG:HD2	1:A:453:ASP:HB2	1.86	0.57
1:B:486:ARG:HD2	6:B:766:HOH:O	2.03	0.57
1:B:63:ARG:CZ	1:B:69:LEU:HD13	2.37	0.54
1:B:44:LEU:HD23	1:B:45:SER:N	2.22	0.54
1:B:157:PRO:O	1:B:161[A]:GLU:HG2	2.09	0.53
1:A:167:MET:HE3	1:A:180:LEU:HG	1.90	0.52
1:A:420:ALA:HB1	2:A:608:EDO:H12	1.91	0.52
1:B:74:LEU:HD22	1:B:158:ARG:HG3	1.90	0.52
1:A:167:MET:HE2	1:A:170:ARG:HD3	1.92	0.51
1:A:451[A]:ARG:HG3	1:A:453:ASP:H	1.75	0.51
1:B:40:TYR:O	1:B:44:LEU:HB3	2.11	0.51
1:B:172:LYS:HG2	1:B:173:GLU:O	2.11	0.51
1:B:225:LEU:HD22	1:B:238:ILE:HB	1.92	0.50
1:A:66:LEU:HB2	1:A:67:PRO:CD	2.42	0.50
1:A:63:ARG:CG	1:A:63:ARG:NH1	2.72	0.50
1:A:476:LEU:HD21	1:A:481:LYS:HB2	1.93	0.50
1:B:330:THR:HA	1:B:416:ARG:O	2.12	0.49
1:A:330:THR:HA	1:A:416:ARG:O	2.13	0.49
1:B:17:THR:HG22	1:B:44:LEU:HD11	1.95	0.49
1:B:101:LEU:HD21	1:B:109:ALA:HB1	1.95	0.47
1:B:226:GLU:OE2	1:B:273:THR:HA	2.14	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:63:ARG:HG2	1:A:160:ARG:HH22	1.79	0.47
1:A:485:CYS:SG	6:A:804:HOH:O	2.60	0.47
1:A:502:PHE:CE2	1:A:504:GLN:HB3	2.50	0.47
1:A:420:ALA:HB1	2:A:608:EDO:H11	1.97	0.46
1:A:225:LEU:HD22	1:A:238:ILE:HB	1.98	0.46
1:B:16:LEU:HD23	1:B:44:LEU:HD12	1.97	0.46
1:A:63:ARG:HH11	1:A:63:ARG:HG2	1.80	0.46
1:B:128:ILE:HD11	1:B:187:LEU:HD13	1.98	0.45
1:A:63:ARG:CG	1:A:160:ARG:HH22	2.29	0.45
1:A:167:MET:CE	1:A:180:LEU:HG	2.45	0.45
1:A:450:CYS:C	1:A:451[B]:ARG:HG3	2.37	0.45
1:B:544:ARG:NH2	1:B:568:GLU:O	2.50	0.45
1:A:123:THR:HG21	1:A:311:GLN:OE1	2.16	0.45
1:A:505:ALA:N	1:A:506:PRO:CD	2.80	0.44
1:A:66:LEU:CB	1:A:67:PRO:HD3	2.45	0.44
1:A:128:ILE:HD11	1:A:187:LEU:HD13	1.99	0.44
1:A:200:LEU:HD22	1:A:210:GLU:OE2	2.17	0.44
1:B:290:PHE:O	1:B:294:LEU:HD13	2.18	0.43
1:A:64:PRO:C	1:A:66:LEU:H	2.21	0.43
1:A:485:CYS:O	1:A:489:GLY:N	2.50	0.42
1:A:63:ARG:NH1	1:A:63:ARG:HG3	2.34	0.42
1:B:68:LEU:HD12	1:B:68:LEU:HA	1.83	0.42
1:A:17:THR:HG22	1:A:44:LEU:HD22	2.01	0.41
1:A:301:LYS:HB3	1:A:301:LYS:HE2	1.85	0.41
1:B:362:THR:HG23	1:B:365:TYR:H	1.85	0.41
1:B:550:LEU:HD23	1:B:550:LEU:HA	1.83	0.41
1:B:337:ALA:HA	1:B:341:LEU:O	2.22	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	526/599 (88%)	504 (96%)	19 (4%)	3 (1%)	22	22
1	B	520/599 (87%)	498 (96%)	21 (4%)	1 (0%)	44	51
All	All	1046/1198 (87%)	1002 (96%)	40 (4%)	4 (0%)	30	33

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	44	LEU
1	B	25	GLU
1	A	66	LEU
1	A	24	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	462/511 (90%)	435 (94%)	27 (6%)	17	19
1	B	456/511 (89%)	427 (94%)	29 (6%)	14	16
All	All	918/1022 (90%)	862 (94%)	56 (6%)	16	17

All (56) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	26	ASN
1	A	37	LEU
1	A	43	HIS
1	A	44	LEU
1	A	63	ARG
1	A	68	LEU
1	A	101	LEU
1	A	149	ASN
1	A	199	GLN
1	A	201	LYS
1	A	265	SER
1	A	293	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	294	LEU
1	A	300	THR
1	A	314	LEU
1	A	323	SER
1	A	325	ASP
1	A	342	ASP
1	A	361	LEU
1	A	362	THR
1	A	451[A]	ARG
1	A	451[B]	ARG
1	A	452	VAL
1	A	453	ASP
1	A	490	SER
1	A	552	MET
1	A	568	GLU
1	B	26	ASN
1	B	44	LEU
1	B	63	ARG
1	B	66	LEU
1	B	68	LEU
1	B	69	LEU
1	B	106	ASP
1	B	107	LEU
1	B	148	ASP
1	B	150	ILE
1	B	170	ARG
1	B	199	GLN
1	B	205	SER
1	B	211	ARG
1	B	231	ASP
1	B	265	SER
1	B	293	LYS
1	B	314	LEU
1	B	322	SER
1	B	362	THR
1	B	453	ASP
1	B	486	ARG
1	B	490	SER
1	B	497[A]	ARG
1	B	497[B]	ARG
1	B	525	LYS
1	B	552	MET

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	555	ASP
1	B	568	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	38	GLN
1	A	130	ASN
1	A	168	GLN
1	A	395	ASN
1	A	513	GLN
1	A	585	GLN
1	B	130	ASN
1	B	149	ASN
1	B	168	GLN
1	B	199	GLN
1	B	311	GLN
1	B	395	ASN
1	B	513	GLN
1	B	585	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 21 ligands modelled in this entry, 4 are monoatomic - leaving 17 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SCN	B	604	-	1,2,2	0.48	0	0,1,1	-	-
3	SCN	B	608	-	1,2,2	0.38	0	0,1,1	-	-
3	SCN	A	606	-	1,2,2	0.76	0	0,1,1	-	-
2	EDO	A	608	-	3,3,3	0.97	0	2,2,2	0.43	0
2	EDO	A	601	-	3,3,3	0.19	0	2,2,2	0.34	0
2	EDO	A	609	-	3,3,3	0.49	0	2,2,2	0.97	0
2	EDO	B	602	-	3,3,3	0.36	0	2,2,2	0.66	0
2	EDO	B	601	-	3,3,3	0.80	0	2,2,2	0.56	0
3	SCN	A	603	-	1,2,2	0.61	0	0,1,1	-	-
2	EDO	B	607	-	3,3,3	0.45	0	2,2,2	0.85	0
3	SCN	A	605	-	1,2,2	2.82	1 (100%)	0,1,1	-	-
3	SCN	B	606	-	1,2,2	0.65	0	0,1,1	-	-
3	SCN	B	605	-	1,2,2	2.18	1 (100%)	0,1,1	-	-
3	SCN	A	604	-	1,2,2	0.67	0	0,1,1	-	-
2	EDO	A	602	-	3,3,3	0.44	0	2,2,2	0.48	0
3	SCN	B	603	-	1,2,2	1.04	0	0,1,1	-	-
2	EDO	A	607	-	3,3,3	0.51	0	2,2,2	0.32	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	A	608	-	-	1/1/1/1	-
2	EDO	A	601	-	-	0/1/1/1	-
2	EDO	A	609	-	-	1/1/1/1	-
2	EDO	B	602	-	-	1/1/1/1	-
2	EDO	B	601	-	-	0/1/1/1	-
2	EDO	B	607	-	-	0/1/1/1	-
2	EDO	A	602	-	-	1/1/1/1	-
2	EDO	A	607	-	-	1/1/1/1	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	605	SCN	C-N	2.82	1.24	1.15
3	B	605	SCN	C-N	2.18	1.22	1.15

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	607	EDO	O1-C1-C2-O2
2	A	608	EDO	O1-C1-C2-O2
2	A	609	EDO	O1-C1-C2-O2
2	B	602	EDO	O1-C1-C2-O2
2	A	602	EDO	O1-C1-C2-O2

There are no ring outliers.

4 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	608	EDO	3	0
2	A	609	EDO	1	0
2	B	607	EDO	1	0
3	B	603	SCN	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	538/599 (89%)	0.34	43 (7%) 20 18	17, 45, 102, 133	2 (0%)
1	B	532/599 (88%)	0.70	67 (12%) 9 7	19, 55, 125, 159	2 (0%)
All	All	1070/1198 (89%)	0.52	110 (10%) 13 11	17, 49, 116, 159	4 (0%)

All (110) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	364	LEU	6.6
1	A	364	LEU	6.6
1	B	361	LEU	6.5
1	A	361	LEU	6.0
1	B	66	LEU	5.9
1	A	37	LEU	5.6
1	B	310	PRO	5.2
1	B	37	LEU	5.2
1	B	107	LEU	4.8
1	B	363	SER	4.7
1	B	249	ALA	4.7
1	A	40	TYR	4.5
1	B	30	PHE	4.4
1	A	66	LEU	4.4
1	A	249	ALA	4.3
1	B	174	LEU	4.2
1	A	30	PHE	4.1
1	B	201	LYS	4.1
1	B	132	ALA	4.0
1	B	29	SER	4.0
1	B	131	ALA	3.9
1	B	243	ILE	3.9
1	B	40	TYR	3.8
1	A	132	ALA	3.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	206	PRO	3.6
1	B	362	THR	3.5
1	B	200	LEU	3.4
1	A	201	LYS	3.4
1	A	362	THR	3.4
1	A	107	LEU	3.4
1	A	200	LEU	3.2
1	B	65	PRO	3.1
1	B	267	LEU	3.1
1	A	309	THR	3.0
1	A	24	ILE	2.9
1	B	101	LEU	2.9
1	B	150	ILE	2.9
1	A	46	HIS	2.9
1	B	283	PHE	2.8
1	A	8	THR	2.8
1	B	94	PHE	2.8
1	B	24	ILE	2.8
1	A	172	LYS	2.7
1	B	109	ALA	2.7
1	B	8	THR	2.7
1	B	205	SER	2.7
1	B	229	PRO	2.7
1	B	227	LEU	2.6
1	B	169	ALA	2.6
1	B	273	THR	2.6
1	B	28	SER	2.6
1	B	164	PHE	2.6
1	B	224	ALA	2.6
1	B	241	CYS	2.6
1	A	65	PRO	2.6
1	A	43	HIS	2.6
1	A	489	GLY	2.6
1	B	67	PRO	2.5
1	B	148	ASP	2.5
1	B	242	CYS	2.5
1	A	487	PHE	2.5
1	B	39	HIS	2.5
1	B	250	PHE	2.5
1	B	228	LEU	2.5
1	A	64	PRO	2.5
1	B	206	PRO	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	302	SER	2.5
1	B	290	PHE	2.5
1	B	38	GLN	2.5
1	B	297	PRO	2.5
1	B	295	LEU	2.4
1	A	451[A]	ARG	2.4
1	A	205	SER	2.4
1	B	106	ASP	2.4
1	B	103	LYS	2.4
1	A	39	HIS	2.4
1	B	167	MET	2.4
1	A	148	ASP	2.4
1	A	452	VAL	2.4
1	A	298	SER	2.4
1	B	112	ILE	2.4
1	B	279	ILE	2.4
1	A	363	SER	2.3
1	A	300	THR	2.3
1	A	490	SER	2.3
1	B	27	GLY	2.3
1	A	308	ASN	2.3
1	B	26	ASN	2.2
1	A	242	CYS	2.2
1	A	28	SER	2.2
1	B	272	SER	2.2
1	B	20	ILE	2.2
1	B	214	LEU	2.2
1	A	94	PHE	2.2
1	B	225	LEU	2.2
1	A	42	GLN	2.2
1	A	199	GLN	2.2
1	B	62	VAL	2.2
1	A	283	PHE	2.2
1	A	9	ALA	2.1
1	A	243	ILE	2.1
1	B	269	LYS	2.1
1	B	102	ILE	2.1
1	B	64	PRO	2.1
1	B	222	LEU	2.1
1	B	171	GLY	2.0
1	B	42	GLN	2.0
1	A	310	PRO	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	108	GLY	2.0
1	B	93	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	EDO	A	607	4/4	0.70	0.22	62,63,78,82	0
2	EDO	B	602	4/4	0.75	0.23	62,70,74,74	0
2	EDO	B	601	4/4	0.80	0.19	42,56,58,71	0
5	K	B	610	1/1	0.80	0.18	75,75,75,75	0
3	SCN	A	604	3/3	0.81	0.20	71,71,75,90	0
2	EDO	A	601	4/4	0.83	0.20	52,74,75,77	0
2	EDO	A	602	4/4	0.84	0.23	60,62,64,71	0
3	SCN	A	606	3/3	0.85	0.16	62,62,81,83	0
3	SCN	A	603	3/3	0.87	0.18	49,49,61,74	0
2	EDO	A	608	4/4	0.87	0.14	38,42,48,50	0
3	SCN	B	606	3/3	0.89	0.22	45,45,61,66	0
3	SCN	B	608	3/3	0.90	0.18	42,42,55,59	0
3	SCN	B	603	3/3	0.90	0.14	36,36,57,82	0
3	SCN	B	604	3/3	0.92	0.18	44,44,53,58	0
5	K	A	611	1/1	0.92	0.14	54,54,54,54	0
2	EDO	A	609	4/4	0.92	0.12	42,45,46,57	0
2	EDO	B	607	4/4	0.93	0.11	37,43,51,53	0
4	ZN	A	610	1/1	0.93	0.07	67,67,67,67	0
3	SCN	A	605	3/3	0.96	0.15	25,25,33,47	0
3	SCN	B	605	3/3	0.97	0.20	29,29,35,58	0
4	ZN	B	609	1/1	0.99	0.03	33,33,33,33	0

6.5 Other polymers [i](#)

There are no such residues in this entry.