



# Full wwPDB X-ray Structure Validation Report

Mar 14, 2018 – 01:43 pm GMT

PDB ID : 2RTC  
Title : APOSTREPTAVIDIN, PH 3.60, SPACE GROUP I222  
Authors : Katz, B.A.  
Deposited on : 1997-09-11  
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

---

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.3 (157068), CSD as539be (2018)  
Xtriage (Phenix) : 1.13  
EDS : trunk31020  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk31020


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	122126	2727 (1.50-1.50)
Ramachandran outliers	120053	2661 (1.50-1.50)
Sidechain outliers	120020	2659 (1.50-1.50)
RSRZ outliers	108989	2481 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	135	
1	D	135	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SO4	B	403[A]	-	-	-	X
2	SO4	B	404[A]	-	-	-	X

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Type</b>	<b>Chain</b>	<b>Res</b>	<b>Chirality</b>	<b>Geometry</b>	<b>Clashes</b>	<b>Electron density</b>
2	SO4	D	400[A]	-	-	-	X

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 4197 atoms, of which 2073 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called STREPTAVIDIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	H	N	O			
1	B	123	Total 1943	C 628	H 939	N 171	O 205	45	13	0
1	D	121	Total 1902	C 613	H 918	N 169	O 202	73	13	0

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	B	1	Total 5	O 4	S 1	0	1
2	B	1	Total 5	O 4	S 1	0	1
2	D	1	Total 5	O 4	S 1	0	1
2	D	1	Total 5	O 4	S 1	0	1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	D	1	Total	O	S	0	1
			5	4	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	53	Total	H	O	0	8
			155	102	53		
3	D	58	Total	H	O	0	2
			172	114	58		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: STREPTAVIDIN



- Molecule 1: STREPTAVIDIN



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	95.18Å 105.39Å 47.89Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	7.50 – 1.50 23.95 – 1.32	Depositor EDS
% Data completeness (in resolution range)	60.0 (7.50-1.50) 65.4 (23.95-1.32)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.38 (at 1.32Å)	Xtrriage
Refinement program	X-PLOR	Depositor
R, $R_{free}$	0.210 , 0.265 0.239 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	13.4	Xtrriage
Anisotropy	0.075	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 118.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	4197	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	31.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.31% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	B	1.46	5/1043 (0.5%)	1.82	33/1428 (2.3%)
1	D	1.40	2/1021 (0.2%)	1.81	31/1399 (2.2%)
All	All	1.43	7/2064 (0.3%)	1.82	64/2827 (2.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	3
1	D	0	3
All	All	0	6

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	45	SER	CA-CB	6.53	1.62	1.52
1	B	92	TRP	NE1-CE2	-5.71	1.30	1.37
1	B	79	TRP	CG-CD2	-5.26	1.34	1.43
1	B	108	TRP	CB-CG	5.25	1.59	1.50
1	B	45	SER	CA-CB	5.20	1.60	1.52
1	D	21	TRP	CG-CD2	-5.16	1.34	1.43
1	B	108	TRP	NE1-CE2	-5.03	1.31	1.37

All (64) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	92	TRP	CD1-NE1-CE2	10.75	118.68	109.00
1	D	108	TRP	CD1-NE1-CE2	10.26	118.24	109.00

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	92	TRP	CD1-NE1-CE2	10.02	118.02	109.00
1	B	108	TRP	CD1-NE1-CE2	10.00	118.00	109.00
1	D	75	TRP	CD1-NE1-CE2	9.98	117.98	109.00
1	B	21	TRP	CD1-NE1-CE2	9.93	117.94	109.00
1	B	75	TRP	CD1-NE1-CE2	9.85	117.87	109.00
1	D	120	TRP	CD1-NE1-CE2	9.72	117.75	109.00
1	D	21	TRP	CD1-NE1-CE2	8.98	117.08	109.00
1	B	120	TRP	CD1-NE1-CE2	8.94	117.04	109.00
1	D	92	TRP	CG-CD1-NE1	-8.39	101.71	110.10
1	B	92	TRP	CG-CD1-NE1	-8.38	101.72	110.10
1	B	108	TRP	NE1-CE2-CZ2	8.37	139.60	130.40
1	B	79	TRP	CD1-NE1-CE2	8.25	116.42	109.00
1	D	108	TRP	CG-CD1-NE1	-8.19	101.91	110.10
1	B	75	TRP	CG-CD1-NE1	-8.13	101.97	110.10
1	D	75	TRP	CG-CD1-NE1	-8.10	102.00	110.10
1	D	59	ARG	NE-CZ-NH2	-8.02	116.29	120.30
1	D	79	TRP	CD1-NE1-CE2	7.90	116.11	109.00
1	B	108	TRP	CG-CD2-CE3	-7.71	126.96	133.90
1	B	21	TRP	NE1-CE2-CZ2	7.70	138.87	130.40
1	B	120	TRP	NE1-CE2-CZ2	7.55	138.70	130.40
1	B	79	TRP	CG-CD1-NE1	-7.50	102.60	110.10
1	D	92	TRP	NE1-CE2-CZ2	7.50	138.65	130.40
1	D	75	TRP	NE1-CE2-CZ2	7.45	138.59	130.40
1	D	120	TRP	CG-CD1-NE1	-7.40	102.70	110.10
1	D	120	TRP	NE1-CE2-CZ2	7.38	138.52	130.40
1	B	108	TRP	NE1-CE2-CD2	-7.13	100.17	107.30
1	D	21	TRP	NE1-CE2-CZ2	7.09	138.20	130.40
1	B	120	TRP	CG-CD1-NE1	-7.08	103.02	110.10
1	D	21	TRP	CG-CD1-NE1	-7.01	103.08	110.10
1	D	108	TRP	NE1-CE2-CZ2	6.92	138.02	130.40
1	D	84	ARG	NE-CZ-NH2	-6.90	116.85	120.30
1	B	21	TRP	CG-CD1-NE1	-6.84	103.26	110.10
1	B	108	TRP	CB-CG-CD2	6.72	135.34	126.60
1	B	78	ALA	N-CA-C	-6.70	92.92	111.00
1	D	78	ALA	N-CA-C	-6.68	92.95	111.00
1	D	79	TRP	NE1-CE2-CZ2	6.61	137.68	130.40
1	B	79	TRP	NE1-CE2-CZ2	6.59	137.65	130.40
1	B	21	TRP	NE1-CE2-CD2	-6.59	100.71	107.30
1	D	120	TRP	NE1-CE2-CD2	-6.50	100.80	107.30
1	D	79	TRP	CG-CD1-NE1	-6.50	103.61	110.10
1	D	75	TRP	NE1-CE2-CD2	-6.41	100.89	107.30
1	B	75	TRP	NE1-CE2-CZ2	6.38	137.41	130.40

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	108	TRP	CE2-CD2-CG	6.29	112.33	107.30
1	D	92	TRP	NE1-CE2-CD2	-6.21	101.08	107.30
1	B	107	GLN	CA-CB-CG	-6.21	99.74	113.40
1	B	108	TRP	CG-CD1-NE1	-6.05	104.05	110.10
1	D	107[A]	GLN	N-CA-C	-6.02	94.75	111.00
1	D	107[B]	GLN	N-CA-C	-6.02	94.75	111.00
1	D	21	TRP	NE1-CE2-CD2	-5.99	101.31	107.30
1	B	108	TRP	CB-CG-CD1	-5.99	119.22	127.00
1	D	108	TRP	NE1-CE2-CD2	-5.97	101.33	107.30
1	B	120	TRP	NE1-CE2-CD2	-5.89	101.41	107.30
1	B	75	TRP	NE1-CE2-CD2	-5.56	101.74	107.30
1	B	114	THR	OG1-CB-CG2	-5.47	97.41	110.00
1	D	83	TYR	CA-CB-CG	-5.29	103.35	113.40
1	D	31	VAL	CB-CA-C	-5.26	101.40	111.40
1	B	104	ILE	N-CA-C	-5.17	97.03	111.00
1	B	107	GLN	N-CA-C	-5.15	97.09	111.00
1	B	113	GLY	N-CA-C	-5.05	100.46	113.10
1	B	67[A]	ASP	CB-CG-OD2	-5.03	113.77	118.30
1	B	67[B]	ASP	CB-CG-OD2	-5.03	113.77	118.30
1	D	115	THR	N-CA-C	-5.02	97.44	111.00

There are no chirality outliers.

All (6) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	103	ARG	Sidechain
1	B	53	ARG	Sidechain
1	B	63[A]	ALA	Peptide
1	D	103	ARG	Sidechain
1	D	53	ARG	Sidechain
1	D	63[A]	ALA	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1004	939	927	15	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	984	918	906	18	0
2	B	10	0	0	0	0
2	D	15	0	0	0	0
3	B	53	102	0	0	4
3	D	58	114	0	1	3
All	All	2124	2073	1833	27	5

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:22[B]:TYR:CE1	1:B:131:THR:HB	2.21	0.76
1:B:114:THR:H	1:D:95:GLN:HE22	1.32	0.75
1:B:95:GLN:HE22	1:D:114:THR:H	1.37	0.71
1:D:51:GLU:O	1:D:52:SER:HB2	2.01	0.60
1:B:51:GLU:O	1:B:52:SER:CB	2.52	0.58
1:B:114:THR:H	1:D:95:GLN:NE2	2.04	0.52
1:B:51:GLU:O	1:B:52:SER:HB3	2.12	0.49
1:D:24:GLN:HG2	1:D:25:LEU:N	2.28	0.49
1:D:51:GLU:O	1:D:52:SER:CB	2.61	0.49
1:B:22[A]:TYR:CE1	1:B:28:THR:HG23	2.48	0.48
1:D:51:GLU:HB2	1:D:84:ARG:NE	2.28	0.48
1:B:114:THR:N	1:D:95:GLN:HE22	2.05	0.47
1:D:23:ASN:HB3	1:D:130:PHE:CE1	2.49	0.47
1:B:61[A]:ASP:OD2	1:D:85:ASN:OD1	2.33	0.47
1:D:13:ALA:N	3:D:1174:HOH:O	2.48	0.47
1:B:133:VAL:HG22	1:B:134:LYS:HG3	1.97	0.45
1:D:110:LEU:C	1:D:110:LEU:HD23	2.38	0.45
1:B:115:THR:HA	1:D:68[B]:GLY:O	2.18	0.44
1:D:120:TRP:CZ2	1:D:121:LYS:HD2	2.53	0.43
1:D:30:ILE:O	1:D:41:GLY:HA3	2.18	0.42
1:B:37:GLY:HA2	1:B:60[B]:TYR:CE1	2.55	0.41
1:B:37:GLY:HA2	1:B:60[A]:TYR:CE1	2.56	0.41
1:B:19:GLY:HA2	1:B:135:PRO:HA	2.02	0.41
1:B:39:LEU:HD13	1:B:60[B]:TYR:CD1	2.56	0.41
1:D:21:TRP:CZ2	1:D:132:LYS:HE3	2.56	0.41
1:D:121:LYS:H	1:D:121:LYS:HG2	1.73	0.41

All (5) symmetry-related close contacts are listed below. The label for Atom-2 includes the sym-

metry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:491[C]:HOH:O	3:B:491[C]:HOH:O[3_655]	0.51	1.69
3:D:1465[C]:HOH:O	3:D:1465[C]:HOH:O[3_556]	0.82	1.38
3:B:491[C]:HOH:O	3:D:1465[C]:HOH:O[8_555]	1.39	0.81
3:B:1460[C]:HOH:O	3:B:1460[C]:HOH:O[3_655]	1.42	0.78
3:B:1460[C]:HOH:O	3:D:1465[C]:HOH:O[8_555]	1.99	0.21

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	134/135 (99%)	123 (92%)	8 (6%)	3 (2%)	7	1
1	D	132/135 (98%)	127 (96%)	2 (2%)	3 (2%)	7	1
All	All	266/270 (98%)	250 (94%)	10 (4%)	6 (2%)	8	1

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	52	SER
1	D	15	ALA
1	D	52	SER
1	D	16	GLY
1	B	64[A]	PRO
1	B	64[B]	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	100/100 (100%)	95 (95%)	5 (5%)	27	4
1	D	98/100 (98%)	92 (94%)	6 (6%)	20	2
All	All	198/200 (99%)	187 (94%)	11 (6%)	26	3

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	39	LEU
1	B	62[A]	SER
1	B	62[B]	SER
1	B	73	LEU
1	B	133	VAL
1	D	31	VAL
1	D	36	ASP
1	D	52	SER
1	D	53	ARG
1	D	73[A]	LEU
1	D	73[B]	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	95	GLN
1	D	95	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	SO4	B	403[A]	-	4,4,4	0.54	0	6,6,6	0.26	0
2	SO4	B	404[A]	-	4,4,4	0.52	0	6,6,6	0.16	0
2	SO4	D	400[A]	-	4,4,4	0.36	0	6,6,6	0.26	0
2	SO4	D	401[B]	-	4,4,4	0.51	0	6,6,6	0.20	0
2	SO4	D	402[A]	-	4,4,4	0.64	0	6,6,6	0.43	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SO4	B	403[A]	-	-	0/0/0/0	0/0/0/0
2	SO4	B	404[A]	-	-	0/0/0/0	0/0/0/0
2	SO4	D	400[A]	-	-	0/0/0/0	0/0/0/0
2	SO4	D	401[B]	-	-	0/0/0/0	0/0/0/0
2	SO4	D	402[A]	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	B	122/135 (90%)	0.41	17 (13%) <b>2</b>   <b>3</b>	10, 25, 48, 55	20 (16%)
1	D	120/135 (88%)	1.23	18 (15%) <b>2</b>   <b>2</b>	13, 28, 61, 72	18 (15%)
All	All	242/270 (89%)	0.82	35 (14%) <b>2</b>   <b>2</b>	10, 27, 53, 72	38 (15%)

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	65[A]	ALA	27.8
1	D	64[A]	PRO	24.7
1	D	66[A]	THR	17.6
1	D	68[A]	GLY	11.2
1	D	13	ALA	9.1
1	B	15	ALA	9.0
1	B	100	ALA	7.8
1	D	63[A]	ALA	6.1
1	D	100	ALA	5.5
1	D	14	GLU	4.5
1	B	13	ALA	4.3
1	D	67[A]	ASP	4.2
1	B	22[A]	TYR	4.1
1	B	66[A]	THR	3.5
1	D	83	TYR	3.2
1	B	65[A]	ALA	3.0
1	B	64[A]	PRO	2.9
1	D	73[A]	LEU	2.7
1	B	29	PHE	2.7
1	B	135	PRO	2.6
1	B	67[A]	ASP	2.5
1	D	51	GLU	2.5
1	B	68[A]	GLY	2.4
1	B	99	GLY	2.4

*Continued on next page...*



Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	49	ASN	2.4
1	D	133	VAL	2.3
1	D	69[A]	SER	2.3
1	D	25	LEU	2.3
1	B	62[A]	SER	2.3
1	B	63[A]	ALA	2.2
1	B	101	GLU	2.1
1	B	35	ALA	2.1
1	D	36	ASP	2.0
1	B	115	THR	2.0
1	D	116	GLU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	SO4	D	401[B]	5/5	0.55	0.29	69,70,72,72	5
2	SO4	D	400[A]	5/5	0.60	0.81	65,65,67,67	5
2	SO4	B	404[A]	5/5	0.72	0.73	55,56,58,59	5
2	SO4	B	403[A]	5/5	0.76	0.43	61,62,63,65	5
2	SO4	D	402[A]	5/5	0.82	0.19	57,57,58,60	5

## 6.5 Other polymers [i](#)

There are no such residues in this entry.