



# wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 9, 2024 – 10:13 am BST

PDB ID : 8RTX  
Title : Crystal Structure of an Anti-idiotypic Fab Fragment  
Authors : Eronen, V.; Parkkinen, T.; Hakulinen, N.; Rouvinen, J.  
Deposited on : 2024-01-29  
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.002 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.38.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	3333 (2.70-2.70)
Clashscore	180529	3684 (2.70-2.70)
Ramachandran outliers	177936	3633 (2.70-2.70)
Sidechain outliers	177891	3633 (2.70-2.70)
RSRZ outliers	164620	3333 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	214	 86% 11%
1	C	214	 79% 21%
1	E	214	 83% 16%
2	B	227	 77% 17%
2	D	227	 78% 18%

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	F	227	 79% 16% ..

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 9738 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Anti-IC Fab fragment B12 light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	214	Total 1625	C 1016	N 273	O 330	S 6	0	0	0
1	C	214	Total 1625	C 1016	N 273	O 330	S 6	0	0	0
1	E	214	Total 1625	C 1016	N 273	O 330	S 6	0	0	0


- Molecule 2 is a protein called Anti-IC Fab fragment B12 heavy chain.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	221	Total 1621	C 1013	N 277	O 324	S 7	0	0	0
2	D	221	Total 1621	C 1013	N 277	O 324	S 7	0	0	0
2	F	221	Total 1621	C 1013	N 277	O 324	S 7	0	0	0

### 3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Anti-IC Fab fragment B12 light chain

Chain A:  86% 11%





- Molecule 1: Anti-IC Fab fragment B12 light chain

Chain C:  79% 21%





- Molecule 1: Anti-IC Fab fragment B12 light chain

Chain E:  83% 16%





- Molecule 2: Anti-IC Fab fragment B12 heavy chain

Chain B:  77% 17%





- Molecule 2: Anti-IC Fab fragment B12 heavy chain

Chain D:  78% 18% ..



- Molecule 2: Anti-IC Fab fragment B12 heavy chain

Chain F:  79% 16% ..



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 31	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	81.86Å 81.86Å 181.93Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	40.93 – 2.70 40.93 – 2.70	Depositor EDS
% Data completeness (in resolution range)	94.5 (40.93-2.70) 85.0 (40.93-2.70)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.54 (at 2.69Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, $R_{free}$	0.247 , 0.269 0.250 , 0.276	Depositor DCC
$R_{free}$ test set	30009 reflections (5.78%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	69.9	Xtrriage
Anisotropy	0.508	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 60.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.016 for -h,-k,l 0.459 for h,-h-k,-l 0.023 for -k,-h,-l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	9738	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	93.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.85% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.46	0/1660	0.86	5/2254 (0.2%)
1	C	0.50	0/1660	0.83	4/2254 (0.2%)
1	E	0.45	0/1660	0.82	2/2254 (0.1%)
2	B	0.54	0/1658	1.04	11/2255 (0.5%)
2	D	0.47	0/1658	0.93	6/2255 (0.3%)
2	F	0.47	0/1658	0.90	3/2255 (0.1%)
All	All	0.48	0/9954	0.90	31/13527 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	D	0	1

There are no bond length outliers.

The worst 5 of 31 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	106	ASP	CB-CG-OD1	11.37	128.53	118.30
2	B	29	PHE	CB-CG-CD1	9.12	127.19	120.80
2	F	79	LEU	CA-CB-CG	8.97	135.93	115.30
1	A	29	ILE	CG1-CB-CG2	-8.71	92.23	111.40
2	B	29	PHE	CB-CG-CD2	-8.30	114.99	120.80

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	D	104	ARG	Sidechain



## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1625	0	1579	17	0
1	C	1625	0	1579	26	0
1	E	1625	0	1580	18	0
2	B	1621	0	1585	29	0
2	D	1621	0	1585	21	0
2	F	1621	0	1586	20	0
All	All	9738	0	9494	127	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 127 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:130:ALA:HB1	2:B:218:PRO:HA	1.74	0.68
2:D:11:LEU:HA	2:D:115:THR:O	1.95	0.67
1:C:6:GLN:HG2	1:C:21:ILE:HD11	1.76	0.66
2:F:17:SER:HA	2:F:83:MET:O	1.95	0.65
2:F:13:GLN:HG3	2:F:14:PRO:HD2	1.81	0.63

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	212/214 (99%)	206 (97%)	6 (3%)	0	<b>100</b> <b>100</b>

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	212/214 (99%)	206 (97%)	6 (3%)	0	100	100
1	E	212/214 (99%)	205 (97%)	7 (3%)	0	100	100
2	B	219/227 (96%)	214 (98%)	5 (2%)	0	100	100
2	D	219/227 (96%)	213 (97%)	6 (3%)	0	100	100
2	F	219/227 (96%)	215 (98%)	3 (1%)	1 (0%)	25	49
All	All	1293/1323 (98%)	1259 (97%)	33 (3%)	1 (0%)	48	73

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	F	131	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	185/185 (100%)	178 (96%)	7 (4%)	28	56
1	C	185/185 (100%)	182 (98%)	3 (2%)	58	82
1	E	185/185 (100%)	182 (98%)	3 (2%)	58	82
2	B	182/188 (97%)	178 (98%)	4 (2%)	47	76
2	D	182/188 (97%)	179 (98%)	3 (2%)	58	82
2	F	182/188 (97%)	177 (97%)	5 (3%)	40	69
All	All	1101/1119 (98%)	1076 (98%)	25 (2%)	45	74

5 of 25 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	D	12	VAL
1	E	126	LYS
2	F	183	LEU
2	D	48	VAL
1	E	185	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	214/214 (100%)	-1.20	0 100 100	67, 87, 108, 136	0
1	C	214/214 (100%)	-1.22	0 100 100	68, 88, 107, 164	0
1	E	214/214 (100%)	-1.19	0 100 100	65, 89, 110, 154	0
2	B	221/227 (97%)	-1.05	0 100 100	77, 101, 133, 169	0
2	D	221/227 (97%)	-1.10	0 100 100	75, 95, 116, 188	0
2	F	221/227 (97%)	-1.09	0 100 100	69, 93, 119, 214	0
All	All	1305/1323 (98%)	-1.14	0 100 100	65, 92, 117, 214	0

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.