



Full wwPDB X-ray Structure Validation Report ⓘ

May 23, 2020 – 07:11 am BST

PDB ID : 3RVG
Title : Crystals structure of Jak2 with a 1-amino-5H-pyrido[4,3-b]indol-4-carboxamide inhibitor
Authors : Lim, J.; Taoka, B.; Otte, R.D.; Spencer, K.; Dinsmore, C.J.; Altman, M.D.; Chan, G.; Rosenstein, C.; Sharma, S.; Su, H.P.; Szewczak, A.A.; Xu, L.; Yin, H.; Zugay-Murphy, J.; Marshall, C.G.; Young, J.R.
Deposited on : 2011-05-06
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	303	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2399 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Tyrosine-protein kinase JAK2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	N	O	P				S
1	A	287	2363	1500	410	437	2	14	0	0	0

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	834	MET	-	EXPRESSION TAG	UNP O60674
A	1133	LEU	-	EXPRESSION TAG	UNP O60674
A	1134	VAL	-	EXPRESSION TAG	UNP O60674
A	1135	PRO	-	EXPRESSION TAG	UNP O60674
A	1136	ARG	-	EXPRESSION TAG	UNP O60674

- Molecule 2 is 1-(cyclohexylamino)-7-(1-methyl-1H-pyrazol-4-yl)-5H-pyrido[4,3-b]indole-4-carboxamide (three-letter code: 17P) (formula: C₂₂H₂₄N₆O).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	29	22	6	1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	7	7	7	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tyrosine-protein kinase JAK2



4 Data and refinement statistics

Property	Value	Source
Space group	P 65	Depositor
Cell constants a, b, c, α , β , γ	124.43 Å 124.43 Å 36.73 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	31.63 – 2.50 34.77 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.7 (31.63-2.50) 99.7 (34.77-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.53 (at 2.51 Å)	Xtrriage
Refinement program	PHENIX 1.6.4_486	Depositor
R, R_{free}	0.221 , 0.300 0.214 , 0.296	Depositor DCC
R_{free} test set	606 reflections (5.24%)	wwPDB-VP
Wilson B-factor (Å ²)	43.2	Xtrriage
Anisotropy	0.076	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 30.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.047 for h,-h-k,-l	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	2399	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.76% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PTR, 17P

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.45	0/2379	0.62	0/3200

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2363	0	2329	69	0
2	A	29	0	24	2	0
3	A	7	0	0	0	0
All	All	2399	0	2353	69	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All (69) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1016:SER:HB3	1:A:1017:PRO:HD3	1.32	1.06

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1016:SER:HB3	1:A:1017:PRO:CD	1.88	1.03
1:A:901:ILE:O	1:A:904:SER:HB2	1.75	0.86
1:A:890:GLU:H	1:A:890:GLU:CD	1.89	0.74
1:A:1122:ARG:O	1:A:1126:ILE:HD13	1.93	0.69
1:A:949:ASP:OD2	1:A:951:ILE:HB	1.93	0.69
1:A:873:ASP:O	1:A:874:ASN:HB2	1.98	0.63
1:A:1119:LEU:O	1:A:1123:VAL:HG23	2.02	0.60
1:A:978:ALA:HA	1:A:1043:VAL:HG22	1.83	0.59
1:A:1123:VAL:O	1:A:1127:ARG:HG3	2.03	0.59
1:A:976:ASP:O	1:A:981:ASN:ND2	2.36	0.58
1:A:1045:TYR:O	1:A:1049:THR:HG23	2.03	0.58
1:A:1115:SER:O	1:A:1119:LEU:HG	2.03	0.58
1:A:867:ARG:HG2	1:A:869:ASP:HB2	1.86	0.58
1:A:1086:GLY:O	1:A:1087:ARG:HD3	2.04	0.57
1:A:976:ASP:HB2	1:A:997:LEU:HD12	1.84	0.57
1:A:934:TYR:CE1	1:A:986:ASN:HA	2.40	0.57
1:A:847:ARG:HD3	1:A:848:HIS:CD2	2.39	0.56
1:A:1109:ASN:HB3	1:A:1112:GLN:HE21	1.71	0.55
1:A:1102:MET:HE3	1:A:1106:TRP:CH2	2.44	0.53
1:A:1104:GLU:CD	1:A:1122:ARG:HH12	2.12	0.52
1:A:1008:PTR:O3P	1:A:1008:PTR:HE1	2.10	0.51
1:A:1124:ASP:HA	1:A:1127:ARG:HD3	1.91	0.51
1:A:893:ARG:HG2	1:A:897:ARG:HH21	1.76	0.51
1:A:1009:LYS:HE3	1:A:1028:GLU:O	2.11	0.51
1:A:892:LEU:O	1:A:896:GLU:HG3	2.12	0.50
1:A:1045:TYR:CE2	1:A:1049:THR:HG21	2.47	0.50
1:A:952:LYS:HE2	1:A:987:GLU:O	2.12	0.49
1:A:841:PRO:HG2	1:A:842:THR:H	1.76	0.49
1:A:890:GLU:N	1:A:890:GLU:CD	2.61	0.49
1:A:883:LYS:HB3	1:A:926:LYS:HG2	1.93	0.49
1:A:855:LEU:HD21	1:A:865:MET:HB2	1.94	0.49
1:A:1058:PRO:O	1:A:1062:MET:HG2	2.12	0.49
1:A:967:LEU:HD13	1:A:972:TYR:O	2.13	0.49
1:A:938:ARG:HB3	1:A:938:ARG:HH11	1.78	0.49
1:A:1045:TYR:HB2	1:A:1102:MET:HE3	1.95	0.48
1:A:1121:LEU:O	1:A:1125:GLN:HG3	2.12	0.48
1:A:931:TYR:CD1	2:A:2000:17P:O	2.67	0.48
1:A:1045:TYR:HD1	1:A:1102:MET:HE1	1.78	0.47
1:A:974:HIS:O	1:A:975:ARG:HB2	2.15	0.46
1:A:938:ARG:HH11	1:A:938:ARG:CB	2.28	0.46
1:A:1049:THR:HG22	1:A:1091:PRO:HB3	1.97	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:884:LEU:HD23	1:A:884:LEU:HA	1.74	0.46
1:A:945:LYS:HE3	1:A:1050:TYR:HB3	1.98	0.45
1:A:932:LEU:HA	1:A:933:PRO:HD3	1.75	0.45
1:A:1016:SER:CB	1:A:1017:PRO:CD	2.75	0.45
1:A:1053:LYS:H	1:A:1053:LYS:HD2	1.82	0.45
1:A:851:PHE:O	1:A:852:LEU:HD23	2.16	0.45
1:A:1032:SER:N	1:A:1035:SER:OG	2.40	0.45
1:A:909:ASN:ND2	1:A:959:GLN:HB3	2.32	0.44
1:A:941:LEU:HD23	1:A:1051:ILE:HG12	1.99	0.44
1:A:1102:MET:CE	1:A:1106:TRP:CH2	3.01	0.44
1:A:931:TYR:HA	2:A:2000:17P:O	2.17	0.44
1:A:938:ARG:O	1:A:942:GLN:HB2	2.18	0.44
1:A:1109:ASN:CG	1:A:1112:GLN:NE2	2.71	0.44
1:A:973:ILE:HD11	1:A:1001:LEU:HD22	2.00	0.44
1:A:1073:MET:O	1:A:1077:HIS:HD2	2.02	0.43
1:A:925:LEU:HD23	1:A:925:LEU:HA	1.78	0.42
1:A:1028:GLU:O	1:A:1029:SER:HB3	2.19	0.42
1:A:1086:GLY:O	1:A:1087:ARG:CD	2.67	0.42
1:A:979:THR:HA	1:A:982:ILE:HD12	2.02	0.42
1:A:1023:PRO:HD3	1:A:1038:TRP:CH2	2.55	0.42
1:A:966:TYR:CZ	1:A:970:LYS:HE3	2.54	0.42
1:A:1025:SER:O	1:A:1029:SER:HA	2.20	0.42
1:A:1089:PRO:O	1:A:1091:PRO:HD3	2.20	0.41
1:A:841:PRO:CG	1:A:842:THR:H	2.33	0.41
1:A:966:TYR:OH	1:A:970:LYS:HE3	2.22	0.40
1:A:1045:TYR:HB2	1:A:1102:MET:CE	2.51	0.40
1:A:1094:CYS:HA	1:A:1095:PRO:HD3	1.88	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	281/303 (93%)	257 (92%)	21 (8%)	3 (1%)	14 26

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	925	LEU
1	A	1070	GLN
1	A	1068	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	256/274 (93%)	233 (91%)	23 (9%)	9 19

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	847	ARG
1	A	890	GLU
1	A	904	SER
1	A	908	ASP
1	A	922	ARG
1	A	925	LEU
1	A	938	ARG
1	A	949	ASP
1	A	952	LYS
1	A	967	LEU
1	A	977	LEU
1	A	980	ARG
1	A	1001	LEU
1	A	1005	LYS
1	A	1016	SER
1	A	1026	LEU
1	A	1052	GLU
1	A	1053	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	1081	LEU
1	A	1103	THR
1	A	1109	ASN
1	A	1118	ASP
1	A	1130	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	874	ASN
1	A	944	HIS
1	A	1077	HIS
1	A	1109	ASN
1	A	1112	GLN
1	A	1125	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
1	PTR	A	1008	1	15,16,17	1.95	1 (6%)	19,22,24	0.87	1 (5%)
1	PTR	A	1007	1	15,16,17	1.95	2 (13%)	19,22,24	0.87	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PTR	A	1008	1	-	2/10/11/13	0/1/1/1
1	PTR	A	1007	1	-	0/10/11/13	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1008	PTR	OH-CZ	-7.12	1.24	1.40
1	A	1007	PTR	OH-CZ	-7.09	1.24	1.40
1	A	1007	PTR	P-OH	2.19	1.62	1.59

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1007	PTR	O3P-P-OH	3.17	115.14	105.24
1	A	1008	PTR	O3P-P-OH	2.18	112.06	105.24

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	1008	PTR	CZ-OH-P-O1P
1	A	1008	PTR	CZ-OH-P-O2P

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	1008	PTR	1	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	17P	A	2000	-	27,33,33	5.87	18 (66%)	37,48,48	2.48	16 (43%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	17P	A	2000	-	-	2/12/20/20	0/5/5/5

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	2000	17P	C22-N23	13.06	1.48	1.35
2	A	2000	17P	C7-N6	10.36	1.48	1.34
2	A	2000	17P	O-C12	9.74	1.42	1.24
2	A	2000	17P	C12-N20	8.32	1.48	1.33
2	A	2000	17P	C8-C9	7.49	1.52	1.36
2	A	2000	17P	C22-C21	7.35	1.52	1.38
2	A	2000	17P	C11-C1	7.31	1.53	1.41
2	A	2000	17P	C5-C4	6.83	1.51	1.39
2	A	2000	17P	C9-C10	6.60	1.53	1.39
2	A	2000	17P	C5-N6	5.99	1.47	1.34
2	A	2000	17P	C7-C3	5.92	1.51	1.44
2	A	2000	17P	C8-C	5.89	1.53	1.41
2	A	2000	17P	C11-C10	5.87	1.53	1.38
2	A	2000	17P	C25-N24	5.64	1.49	1.33
2	A	2000	17P	C4-C2	5.06	1.51	1.43
2	A	2000	17P	C-C1	4.26	1.54	1.42
2	A	2000	17P	C1-N	2.99	1.48	1.38
2	A	2000	17P	C2-N	2.32	1.48	1.39

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	2000	17P	C5-C4-C2	-6.63	114.30	118.42
2	A	2000	17P	C10-C11-C1	-6.30	115.93	121.44
2	A	2000	17P	O-C12-C4	-4.08	115.41	120.22

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	2000	17P	C3-C7-N6	-4.04	118.02	121.35
2	A	2000	17P	C22-C21-C10	-4.01	122.49	127.74
2	A	2000	17P	C9-C8-C	-3.95	115.94	121.63
2	A	2000	17P	C25-C21-C10	3.64	132.52	127.74
2	A	2000	17P	C5-C4-C12	2.94	123.93	117.86
2	A	2000	17P	C7-N13-C14	-2.90	118.64	124.16
2	A	2000	17P	C4-C5-N6	-2.63	120.36	124.49
2	A	2000	17P	C5-N6-C7	2.34	120.75	116.05
2	A	2000	17P	C11-C1-C	2.32	124.15	121.16
2	A	2000	17P	C4-C12-N20	-2.27	114.83	118.28
2	A	2000	17P	O-C12-N20	2.18	125.67	122.58
2	A	2000	17P	C19-C14-C15	2.02	114.32	110.82
2	A	2000	17P	C9-C10-C11	-2.02	114.92	118.09

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	2000	17P	C11-C10-C21-C22
2	A	2000	17P	C11-C10-C21-C25

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	2000	17P	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	285/303 (94%)	-0.19	5 (1%) 68 71	25, 41, 65, 86	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1053	LYS	3.8
1	A	943	LYS	2.6
1	A	946	GLU	2.6
1	A	1016	SER	2.4
1	A	944	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	PTR	A	1007	16/17	0.88	0.17	41,59,75,76	0
1	PTR	A	1008	16/17	0.94	0.16	40,49,74,82	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	17P	A	2000	29/29	0.94	0.14	21,29,39,42	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.