



Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2020 – 10:33 pm BST

PDB ID : 3RWX
Title : Crystal structure of a putative outer membrane protein (BF2706) from *Bacteroides fragilis* NCTC 9343 at 2.40 Å resolution
Authors : Joint Center for Structural Genomics (JCSG)
Deposited on : 2011-05-09
Resolution : 2.40 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	261	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 2162 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Hypothetical bacterial outer membrane protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	254	1996	1261	309	416	2	8	0	14	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	leader sequence	UNP Q5LBW6

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	A	1	5	4	1	0	0
2	A	1	5	4	1	0	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	113	Total O 114 114	0	1

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Hypothetical bacterial outer membrane protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	154.15Å 154.15Å 40.16Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	29.82 – 2.40 29.82 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.8 (29.82-2.40) 99.8 (29.82-2.40)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.04 (at 2.39Å)	Xtrriage
Refinement program	REFMAC 5.5.0110	Depositor
R, R_{free}	0.198 , 0.242 0.191 , 0.227	Depositor DCC
R_{free} test set	1107 reflections (5.13%)	wwPDB-VP
Wilson B-factor (Å ²)	38.9	Xtrriage
Anisotropy	0.424	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 36.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.049 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2162	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.25% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MLY, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.64	0/1829	0.69	0/2492

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1996	0	2003	20	0
2	A	10	0	0	1	0
3	A	42	0	56	1	0
4	A	114	0	0	2	0
All	All	2162	0	2059	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:181:MLY:HH23	2:A:284:SO4:O3	1.81	0.80
1:A:211[A]:MSE:SE	1:A:275:ILE:HD11	2.33	0.77
1:A:32:MLY:HH12	1:A:92:SER:H	1.50	0.74
1:A:211[B]:MSE:SE	1:A:271:MSE:HE1	2.48	0.63
1:A:56:MSE:HE2	1:A:142:ILE:HD11	1.82	0.61
1:A:56:MSE:HE2	1:A:142:ILE:CD1	2.32	0.59
1:A:161:MLY:HH23	1:A:283:MLY:OXT	2.04	0.57
1:A:129:LEU:HB3	1:A:144:PHE:HB3	1.87	0.56
1:A:106:ASN:HA	4:A:389:HOH:O	2.10	0.50
1:A:121:ILE:HD12	1:A:121:ILE:N	2.26	0.50
1:A:131:VAL:HG22	1:A:142:ILE:HB	1.93	0.50
1:A:211[A]:MSE:SE	1:A:275:ILE:CD1	3.09	0.47
1:A:32:MLY:HH23	1:A:92:SER:HA	1.97	0.46
1:A:89[A]:MLY:HH22	1:A:89[A]:MLY:HD3	1.68	0.44
1:A:123[A]:GLU:CD	1:A:123[A]:GLU:H	2.21	0.43
1:A:118[B]:THR:HG23	4:A:351:HOH:O	2.18	0.43
1:A:211[B]:MSE:SE	1:A:271:MSE:CE	3.15	0.42
1:A:91:THR:O	1:A:94:THR:HB	2.21	0.41
1:A:278:MSE:SE	3:A:287:GOL:O2	2.89	0.40
1:A:117:LEU:C	1:A:117:LEU:HD23	2.42	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	246/261 (94%)	240 (98%)	5 (2%)	1 (0%)	34 48

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	109	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	211/198 (107%)	209 (99%)	2 (1%)	78 90

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	124	GLN
1	A	205	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

21 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	MLY	A	84	1	9,10,11	0.72	0	6,11,13	0.56	0
1	MLY	A	93	1	9,10,11	0.61	0	6,11,13	0.71	0
1	MLY	A	222	1	9,10,11	0.85	0	6,11,13	0.69	0
1	MLY	A	161	1	9,10,11	0.68	0	6,11,13	0.35	0
1	MLY	A	197	1	9,10,11	0.79	0	6,11,13	0.79	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	MLY	A	280	1	9,10,11	0.70	0	6,11,13	0.18	0
1	MLY	A	39	1	9,10,11	0.64	0	6,11,13	0.61	0
1	MLY	A	267	1	9,10,11	0.80	0	6,11,13	0.38	0
1	MLY	A	108	1	9,10,11	0.68	0	6,11,13	0.35	0
1	MLY	A	185	1	9,10,11	1.01	0	6,11,13	0.47	0
1	MLY	A	283	1	7,11,11	0.51	0	7,13,13	0.92	0
1	MLY	A	130	1	9,10,11	0.66	0	6,11,13	0.98	0
1	MLY	A	89[A]	-	9,10,11	0.55	0	6,11,13	0.20	0
1	MLY	A	126	1	9,10,11	0.73	0	6,11,13	0.45	0
1	MLY	A	89[B]	-	9,10,11	0.47	0	6,11,13	0.51	0
1	MLY	A	181	1	9,10,11	0.67	0	6,11,13	1.98	1 (16%)
1	MLY	A	32	1	9,10,11	0.89	0	6,11,13	0.55	0
1	MLY	A	113	1	9,10,11	0.69	0	6,11,13	0.82	0
1	MLY	A	282	1	9,10,11	0.76	0	6,11,13	0.74	0
1	MLY	A	59	1	5,6,11	0.67	0	2,6,13	0.36	0
1	MLY	A	232	1	9,10,11	0.65	0	6,11,13	0.61	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	MLY	A	84	1	-	0/8/9/11	-
1	MLY	A	93	1	-	2/8/9/11	-
1	MLY	A	222	1	-	3/8/9/11	-
1	MLY	A	161	1	-	1/8/9/11	-
1	MLY	A	197	1	-	0/8/9/11	-
1	MLY	A	280	1	-	0/8/9/11	-
1	MLY	A	39	1	-	0/8/9/11	-
1	MLY	A	267	1	-	4/8/9/11	-
1	MLY	A	108	1	-	2/8/9/11	-
1	MLY	A	185	1	-	2/8/9/11	-
1	MLY	A	283	1	-	2/7/11/11	-
1	MLY	A	130	1	-	5/8/9/11	-
1	MLY	A	89[A]	-	-	4/8/9/11	-
1	MLY	A	126	1	-	3/8/9/11	-
1	MLY	A	89[B]	-	-	1/8/9/11	-
1	MLY	A	181	1	-	2/8/9/11	-
1	MLY	A	32	1	-	5/8/9/11	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	MLY	A	113	1	-	4/8/9/11	-
1	MLY	A	282	1	-	2/8/9/11	-
1	MLY	A	59	1	-	1/4/5/11	-
1	MLY	A	232	1	-	6/8/9/11	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	181	MLY	CD-CE-NZ	-3.95	103.10	113.79

There are no chirality outliers.

All (49) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	161	MLY	O-C-CA-CB
1	A	222	MLY	O-C-CA-CB
1	A	267	MLY	C-CA-CB-CG
1	A	130	MLY	N-CA-CB-CG
1	A	130	MLY	C-CA-CB-CG
1	A	126	MLY	O-C-CA-CB
1	A	32	MLY	O-C-CA-CB
1	A	232	MLY	N-CA-CB-CG
1	A	232	MLY	C-CA-CB-CG
1	A	232	MLY	CD-CE-NZ-CH1
1	A	113	MLY	CD-CE-NZ-CH1
1	A	113	MLY	CD-CE-NZ-CH2
1	A	130	MLY	CD-CE-NZ-CH1
1	A	89[A]	MLY	CD-CE-NZ-CH2
1	A	181	MLY	CD-CE-NZ-CH1
1	A	181	MLY	CD-CE-NZ-CH2
1	A	232	MLY	CD-CE-NZ-CH2
1	A	59	MLY	CA-CB-CG-CD
1	A	113	MLY	CG-CD-CE-NZ
1	A	222	MLY	CG-CD-CE-NZ
1	A	89[A]	MLY	CG-CD-CE-NZ
1	A	267	MLY	CD-CE-NZ-CH2
1	A	89[A]	MLY	CD-CE-NZ-CH1
1	A	89[B]	MLY	CD-CE-NZ-CH2
1	A	93	MLY	CD-CE-NZ-CH2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
1	A	130	MLY	CD-CE-NZ-CH2
1	A	93	MLY	CD-CE-NZ-CH1
1	A	32	MLY	CE-CD-CG-CB
1	A	89[A]	MLY	CA-CB-CG-CD
1	A	32	MLY	C-CA-CB-CG
1	A	108	MLY	CE-CD-CG-CB
1	A	232	MLY	CE-CD-CG-CB
1	A	283	MLY	CD-CE-NZ-CH1
1	A	126	MLY	CD-CE-NZ-CH2
1	A	232	MLY	CA-CB-CG-CD
1	A	283	MLY	CD-CE-NZ-CH2
1	A	185	MLY	CD-CE-NZ-CH1
1	A	126	MLY	CD-CE-NZ-CH1
1	A	32	MLY	CG-CD-CE-NZ
1	A	282	MLY	CD-CE-NZ-CH2
1	A	267	MLY	N-CA-CB-CG
1	A	130	MLY	CG-CD-CE-NZ
1	A	222	MLY	CD-CE-NZ-CH1
1	A	108	MLY	CG-CD-CE-NZ
1	A	267	MLY	CG-CD-CE-NZ
1	A	32	MLY	CD-CE-NZ-CH1
1	A	185	MLY	CG-CD-CE-NZ
1	A	282	MLY	CG-CD-CE-NZ
1	A	113	MLY	CE-CD-CG-CB

There are no ring outliers.

5 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	161	MLY	1	0
1	A	283	MLY	1	0
1	A	89[A]	MLY	1	0
1	A	181	MLY	1	0
1	A	32	MLY	2	0

5.5 Carbohydrates

There are no carbohydrates in this entry.

5.6 Ligand geometry

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GOL	A	292	-	5,5,5	0.48	0	5,5,5	0.51	0
3	GOL	A	288	-	5,5,5	0.36	0	5,5,5	0.47	0
3	GOL	A	286	-	5,5,5	0.39	0	5,5,5	0.28	0
3	GOL	A	290	-	5,5,5	0.37	0	5,5,5	0.36	0
2	SO4	A	284	-	4,4,4	0.17	0	6,6,6	0.56	0
3	GOL	A	291	-	5,5,5	0.41	0	5,5,5	0.23	0
2	SO4	A	285	-	4,4,4	0.22	0	6,6,6	0.19	0
3	GOL	A	287	-	5,5,5	0.42	0	5,5,5	0.60	0
3	GOL	A	289	-	5,5,5	0.46	0	5,5,5	0.37	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	292	-	-	1/4/4/4	-
3	GOL	A	288	-	-	3/4/4/4	-
3	GOL	A	286	-	-	2/4/4/4	-
3	GOL	A	290	-	-	1/4/4/4	-
3	GOL	A	291	-	-	2/4/4/4	-
3	GOL	A	289	-	-	4/4/4/4	-
3	GOL	A	287	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (13) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	286	GOL	C1-C2-C3-O3
3	A	291	GOL	C1-C2-C3-O3
3	A	289	GOL	C1-C2-C3-O3
3	A	286	GOL	O2-C2-C3-O3
3	A	288	GOL	C1-C2-C3-O3
3	A	289	GOL	O1-C1-C2-C3
3	A	291	GOL	O2-C2-C3-O3
3	A	289	GOL	O1-C1-C2-O2
3	A	289	GOL	O2-C2-C3-O3
3	A	288	GOL	O1-C1-C2-C3
3	A	292	GOL	O2-C2-C3-O3
3	A	288	GOL	O2-C2-C3-O3
3	A	290	GOL	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	284	SO4	1	0
3	A	287	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	227/261 (86%)	-0.30	5 (2%) 62 60	20, 30, 45, 66	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	109	PRO	6.0
1	A	110	ALA	3.6
1	A	107	ASP	3.5
1	A	123[A]	GLU	3.0
1	A	244	ASN	2.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	MLY	A	108	11/12	0.81	0.27	60,62,63,64	0
1	MLY	A	185	11/12	0.83	0.24	29,32,56,56	0
1	MLY	A	32	11/12	0.88	0.19	40,45,62,63	0
1	MLY	A	222	11/12	0.88	0.25	27,31,54,55	0
1	MLY	A	282	11/12	0.89	0.20	28,30,50,52	0
1	MLY	A	130	11/12	0.90	0.25	32,34,54,55	0
1	MLY	A	84	11/12	0.91	0.20	31,32,51,52	0
1	MLY	A	126	11/12	0.92	0.17	36,42,54,55	0
1	MLY	A	283	12/12	0.92	0.14	28,30,32,32	0
1	MLY	A	93	11/12	0.93	0.15	36,40,52,54	0
1	MLY	A	267	11/12	0.93	0.20	30,33,53,53	0
1	MLY	A	232	11/12	0.93	0.19	27,30,51,53	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	MLY	A	113	11/12	0.94	0.14	39,44,55,55	0
1	MLY	A	89[A]	11/12	0.95	0.18	34,35,38,38	8
1	MLY	A	89[B]	11/12	0.95	0.18	34,34,35,36	8
1	MLY	A	197	11/12	0.96	0.13	23,26,41,42	0
1	MLY	A	280	11/12	0.96	0.15	25,27,44,46	0
1	MLY	A	39	11/12	0.96	0.11	24,25,37,37	0
1	MLY	A	161	11/12	0.96	0.13	25,27,38,39	0
1	MLY	A	59	7/12	0.97	0.21	31,32,37,40	0
1	MLY	A	181	11/12	0.97	0.14	23,25,29,30	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	A	289	6/6	0.56	0.35	67,68,68,69	0
3	GOL	A	292	6/6	0.68	0.34	73,74,75,75	0
3	GOL	A	287	6/6	0.69	0.25	66,67,68,68	0
3	GOL	A	291	6/6	0.70	0.33	89,91,92,92	0
3	GOL	A	288	6/6	0.79	0.22	75,75,76,76	0
3	GOL	A	286	6/6	0.93	0.18	48,53,55,56	0
2	SO4	A	285	5/5	0.93	0.21	75,75,76,77	0
3	GOL	A	290	6/6	0.96	0.26	57,58,60,60	0
2	SO4	A	284	5/5	0.99	0.11	42,43,44,44	0

6.5 Other polymers [i](#)

There are no such residues in this entry.