



Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2020 – 03:58 am BST

PDB ID : 3RXZ
Title : Crystal structure of putative polysaccharide deacetylase from *Mycobacterium smegmatis*
Authors : Michalska, K.; Tesar, C.; Bearden, J.; Joachimiak, A.; Midwest Center for Structural Genomics (MCSG)
Deposited on : 2011-05-10
Resolution : 2.01 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.01 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	300	 6% 85% 10% 5%
1	B	300	 5% 90% 5% 5%
1	C	300	 5% 90% 6% •
1	D	300	 6% 88% 8% •

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 9319 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Polysaccharide deacetylase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	286	2224	1422	380	411	3	8	0	0	0
1	B	286	2226	1423	380	412	3	8	0	0	0
1	C	287	2242	1432	382	417	3	8	0	1	0
1	D	287	2253	1438	387	417	3	8	0	2	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	EXPRESSION TAG	UNP A0R0G0
A	-1	ASN	-	EXPRESSION TAG	UNP A0R0G0
A	0	ALA	-	EXPRESSION TAG	UNP A0R0G0
B	-2	SER	-	EXPRESSION TAG	UNP A0R0G0
B	-1	ASN	-	EXPRESSION TAG	UNP A0R0G0
B	0	ALA	-	EXPRESSION TAG	UNP A0R0G0
C	-2	SER	-	EXPRESSION TAG	UNP A0R0G0
C	-1	ASN	-	EXPRESSION TAG	UNP A0R0G0
C	0	ALA	-	EXPRESSION TAG	UNP A0R0G0
D	-2	SER	-	EXPRESSION TAG	UNP A0R0G0
D	-1	ASN	-	EXPRESSION TAG	UNP A0R0G0
D	0	ALA	-	EXPRESSION TAG	UNP A0R0G0

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Cl	0	0
			1	1		
2	A	1	Total	Cl	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	D	1	Total Cl 1 1	0	0
2	C	1	Total Cl 1 1	0	0

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total Zn 1 1	0	0
3	A	1	Total Zn 1 1	0	0
3	D	1	Total Zn 1 1	0	0
3	C	1	Total Zn 1 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	102	Total O 102 102	0	0
4	B	79	Total O 79 79	0	4
4	C	88	Total O 88 88	0	2
4	D	97	Total O 97 97	0	3


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Polysaccharide deacetylase




- Molecule 1: Polysaccharide deacetylase



- Molecule 1: Polysaccharide deacetylase



- Molecule 1: Polysaccharide deacetylase





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	80.21Å 59.59Å 130.39Å 90.00° 95.35° 90.00°	Depositor
Resolution (Å)	29.13 – 2.01 29.13 – 2.01	Depositor EDS
% Data completeness (in resolution range)	(Not available) (29.13-2.01) 99.1 (29.13-2.01)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.18 (at 2.01Å)	Xtrriage
Refinement program	BUSTER 2.8.0	Depositor
R, R_{free}	0.175 , 0.211 0.180 , 0.230	Depositor DCC
R_{free} test set	1216 reflections (1.50%)	wwPDB-VP
Wilson B-factor (Å ²)	29.0	Xtrriage
Anisotropy	0.797	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 48.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	9319	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.92% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.76	1/2284 (0.0%)	0.72	0/3110
1	B	0.69	0/2286	0.70	0/3112
1	C	0.72	1/2302 (0.0%)	0.72	0/3134
1	D	0.70	0/2313	0.66	0/3148
All	All	0.72	2/9185 (0.0%)	0.70	0/12504

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	168	MSE	SE-CE	-5.50	1.63	1.95
1	C	148	MSE	SE-CE	-5.07	1.65	1.95

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2224	0	2141	25	0
1	B	2226	0	2140	10	0
1	C	2242	0	2152	11	0
1	D	2253	0	2166	13	0
2	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	102	0	0	0	0
4	B	79	0	0	0	0
4	C	88	0	0	0	0
4	D	97	0	0	0	0
All	All	9319	0	8599	50	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:168:MSE:HE3	1:A:189:SER:HA	1.15	1.15
1:A:168:MSE:CE	1:A:189:SER:HA	2.02	0.87
1:A:168:MSE:HE3	1:A:189:SER:CA	2.09	0.73
1:C:8:LEU:HD22	1:C:287:LEU:HD23	1.79	0.65
1:A:178:VAL:O	1:A:178:VAL:HG12	1.98	0.64
1:A:173:PRO:HB3	1:A:186:LEU:HD23	1.79	0.63
1:A:37:LEU:HD13	1:B:292:LEU:HD11	1.82	0.61
1:D:89:GLU:OE2	1:D:96:ARG:NH2	2.33	0.60
1:A:47:MSE:HE2	1:B:167:LEU:HD21	1.85	0.59
1:A:47:MSE:CE	1:B:167:LEU:HD11	2.34	0.58
1:A:148:MSE:HE3	1:A:167:LEU:HD21	1.86	0.57
1:A:8:LEU:HD23	1:A:287:LEU:HD23	1.89	0.55
1:D:148:MSE:HE2	1:D:167:LEU:HD21	1.90	0.54
1:A:154:LYS:HA	1:A:178:VAL:HG13	1.91	0.52
1:B:21:THR:O	1:B:231:GLY:HA2	2.10	0.52
1:D:36:LEU:HD21	1:D:201:VAL:HG11	1.91	0.52
1:A:47:MSE:HE3	1:B:167:LEU:HD11	1.92	0.51
1:C:36:LEU:HD21	1:C:201:VAL:HG11	1.94	0.50
1:A:219:LEU:HD12	1:D:212:THR:HA	1.94	0.50
1:A:154:LYS:HA	1:A:178:VAL:CG1	2.42	0.50
1:A:168:MSE:HE2	1:D:52:HIS:HD2	1.77	0.50
1:A:198:TYR:CZ	1:A:248:ARG:HG2	2.48	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:140:GLY:HA3	1:B:163:TYR:CZ	2.47	0.49
1:A:292:LEU:HD11	1:D:37:LEU:HD13	1.93	0.49
1:C:8:LEU:HD21	1:C:285:ARG:NH2	2.29	0.48
1:C:8:LEU:CD2	1:C:287:LEU:HD23	2.43	0.47
1:D:140:GLY:HA3	1:D:163:TYR:CZ	2.50	0.47
1:A:14:VAL:HG22	1:A:282:LEU:HD11	1.97	0.46
1:D:151:HIS:O	1:D:154:LYS:HG2	2.17	0.45
1:A:28:PHE:HA	1:A:238:ASN:O	2.15	0.45
1:C:84:ALA:HA	1:C:91:ILE:HG13	1.98	0.45
1:C:131:GLU:OE2	1:C:137:HIS:ND1	2.44	0.45
1:A:148:MSE:CE	1:A:167:LEU:HD21	2.46	0.44
1:D:21:THR:O	1:D:231:GLY:HA2	2.17	0.43
1:A:157:ALA:CB	1:A:178:VAL:HG11	2.49	0.43
1:C:89:GLU:H	1:C:89:GLU:CD	2.22	0.42
1:C:8:LEU:HD13	1:C:286:THR:O	2.19	0.42
1:B:11:ALA:HB3	1:B:170:SER:OG	2.19	0.42
1:B:8:LEU:HD22	1:B:287:LEU:HD23	2.02	0.42
1:D:227:MSE:HG2	1:D:234:TRP:HB2	2.01	0.42
1:A:21:THR:O	1:A:231:GLY:HA2	2.20	0.41
1:A:230:ILE:HD12	1:D:246:PRO:HB3	2.01	0.41
1:B:176:LEU:HD23	1:B:183:LEU:HD23	2.01	0.41
1:C:140:GLY:HA3	1:C:163:TYR:CZ	2.56	0.41
1:B:36:LEU:HD21	1:B:201:VAL:HG11	2.01	0.41
1:D:28:PHE:HA	1:D:238:ASN:O	2.20	0.41
1:C:28:PHE:HA	1:C:238:ASN:O	2.21	0.41
1:C:208:GLY:O	1:D:196:GLN:HB3	2.19	0.41
1:A:40:ASP:HA	1:A:41:PRO:HD2	1.95	0.41
1:A:36:LEU:HD21	1:A:201:VAL:HG11	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	284/300 (95%)	271 (95%)	13 (5%)	0	100	100
1	B	284/300 (95%)	269 (95%)	15 (5%)	0	100	100
1	C	286/300 (95%)	275 (96%)	11 (4%)	0	100	100
1	D	287/300 (96%)	274 (96%)	13 (4%)	0	100	100
All	All	1141/1200 (95%)	1089 (95%)	52 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	231/234 (99%)	227 (98%)	4 (2%)	60	65
1	B	231/234 (99%)	229 (99%)	2 (1%)	78	83
1	C	233/234 (100%)	232 (100%)	1 (0%)	91	93
1	D	234/234 (100%)	232 (99%)	2 (1%)	78	83
All	All	929/936 (99%)	920 (99%)	9 (1%)	76	81

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	22	CYS
1	A	117	ASP
1	A	125	ARG
1	A	155	LEU
1	B	96	ARG
1	B	117	ASP
1	C	9	THR
1	D	132	GLU
1	D	291	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 8 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	278/300 (92%)	0.04	18 (6%) 18 18	25, 35, 58, 81	0
1	B	278/300 (92%)	0.05	16 (5%) 23 22	27, 40, 63, 90	0
1	C	279/300 (93%)	-0.02	15 (5%) 25 24	26, 36, 57, 83	0
1	D	279/300 (93%)	0.07	17 (6%) 21 20	26, 39, 60, 99	0
All	All	1114/1200 (92%)	0.03	66 (5%) 22 21	25, 38, 60, 99	0

All (66) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	8	LEU	4.9
1	B	207	THR	4.9
1	D	207	THR	4.5
1	B	180	ASP	3.9
1	A	293	THR	3.8
1	D	235	VAL	3.8
1	D	188	VAL	3.7
1	D	165	SER	3.6
1	A	180	ASP	3.5
1	A	203	ASP	3.3
1	C	206	GLY	3.3
1	C	15	SER	3.2
1	A	181	GLY	3.2
1	D	166	THR	3.2
1	A	207	THR	3.1
1	B	188	VAL	3.1
1	B	202	PRO	3.1
1	D	181	GLY	3.1
1	C	207	THR	3.1
1	C	203	ASP	3.0
1	B	206	GLY	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	237	THR	2.9
1	A	204	PHE	2.9
1	D	204	PHE	2.9
1	D	180	ASP	2.8
1	D	7	GLU	2.8
1	B	165	SER	2.8
1	B	205	SER	2.7
1	D	186	LEU	2.7
1	B	203	ASP	2.7
1	A	209	LEU	2.7
1	C	165	SER	2.7
1	A	188	VAL	2.6
1	B	209	LEU	2.6
1	B	7	GLU	2.6
1	C	204	PHE	2.6
1	B	204	PHE	2.6
1	D	293	THR	2.6
1	B	208	GLY	2.6
1	C	27	THR	2.5
1	A	208	GLY	2.5
1	A	235	VAL	2.5
1	D	203	ASP	2.5
1	A	236	LEU	2.4
1	C	205	SER	2.4
1	B	237	THR	2.4
1	C	186	LEU	2.4
1	D	167	LEU	2.4
1	C	237	THR	2.4
1	C	236	LEU	2.4
1	B	199	CYS	2.4
1	C	188	VAL	2.2
1	C	261	ALA	2.2
1	C	143	ALA	2.2
1	D	279	ALA	2.2
1	A	186	LEU	2.2
1	D	192	LEU	2.1
1	A	78	PHE	2.1
1	D	292	LEU	2.1
1	C	208	GLY	2.1
1	B	27	THR	2.1
1	B	236	LEU	2.1
1	D	78	PHE	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	27	THR	2.1
1	A	202	PRO	2.0
1	A	142	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CL	A	298	1/1	0.98	0.12	27,27,27,27	0
2	CL	D	298	1/1	0.99	0.07	27,27,27,27	0
3	ZN	B	299	1/1	0.99	0.08	36,36,36,36	1
3	ZN	A	299	1/1	0.99	0.08	33,33,33,33	1
3	ZN	C	299	1/1	1.00	0.04	38,38,38,38	0
3	ZN	D	299	1/1	1.00	0.06	35,35,35,35	0
2	CL	C	298	1/1	1.00	0.10	28,28,28,28	0
2	CL	B	298	1/1	1.00	0.10	29,29,29,29	0

6.5 Other polymers [i](#)

There are no such residues in this entry.