

wwPDB X-ray Structure Validation Summary Report (i)

Mar 21, 2024 – 12:42 pm GMT

PDB ID	:	8RXU
Title	:	Crystal structure of octaheme nitrite reductase from Trichlorobacter ammonif-
		icans in space group P21
Authors	:	Polyakov, K.M.; Safonova, T.N.; Osipov, E.; Popov, A.N.; Tikhonova, T.V.;
		Popov, V.O.
Deposited on	:	2024-02-08
Resolution	:	1.74 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY\;DIFFRACTION$

The reported resolution of this entry is 1.74 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
	$(\# { m Entries})$	$(\# { m Entries}, { m resolution} { m range}({ m \AA}))$
R _{free}	130704	3764(1.76-1.72)
Clashscore	141614	3923 (1.76-1.72)
Ramachandran outliers	138981	3878(1.76-1.72)
Sidechain outliers	138945	3878 (1.76-1.72)
RSRZ outliers	127900	3705 (1.76-1.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	AAA	495	% 69%	28%	•
1	MMM	495	% 71%	25%	•
1	YYY	495	68%	30%	·

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard



residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	PO4	YYY	610	-	-	Х	-



$8 \mathrm{RXU}$

2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 13690 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	A A A	405	Total	С	Ν	0	\mathbf{S}	0	2	0
	AAA	495	3895	2437	714	717	27	0		
1	ммл	405	Total	С	Ν	0	S	0	3	0
		495	3894	2438	714	715	27	0		0
1	VVV	405	Total	С	Ν	0	S	0	2	0
		495	3898	2440	712	717	29	0		0

• Molecule 1 is a protein called Octaheme nitrite reductase.

• Molecule 2 is HEME C (three-letter code: HEC) (formula: C₃₄H₃₄FeN₄O₄) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues		Ate	oms			ZeroOcc	AltConf
9	ΛΛΛ	1	Total	С	Fe	Ν	Ο	0	0
Z AAA	1	43	34	1	4	4	0	0	
0	ΛΛΛ	1	Total	С	Fe	Ν	Ο	0	0
	AAA	1	43	34	1	4	4	0	0
0	ΛΛΛ	1	Total	С	Fe	Ν	Ο	0	0
	AAA	1	43	34	1	4	4	0	0



Continued from previous page...

Mol	Chain	Residues		Ate	oms			ZeroOcc	AltConf
0		1	Total	С	Fe	Ν	Ο	0	0
2	AAA	1	43	34	1	4	4	0	0
0		1	Total	С	Fe	Ν	Ο	0	0
	AAA	1	43	34	1	4	4	0	0
0	ΛΛΛ	1	Total	С	Fe	Ν	Ο	0	0
	AAA	1	43	34	1	4	4	0	0
9	ΛΛΛ	1	Total	С	Fe	Ν	0	0	1
2	ллл	1	46	35	1	4	6	0	T
2	ΔΔΔ	1	Total	С	Fe	Ν	Ο	0	0
	11111	1	43	34	1	4	4	0	0
2	MMM	1	Total	С	Fe	Ν	Ο	0	0
		1	43	34	1	4	4	0	0
2	MMM	1	Total	С	Fe	Ν	Ο	0	0
		-	43	34	1	4	4		
2	MMM	1	Total	С	Fe	Ν	Ο	0	0
		-	43	34	1	4	4	Ŭ	
2	MMM	1	Total	С	Fe	Ν	O	0	0
		_	43	34	1	4	4		
2	MMM	1	Total	С	Fe	N	O	0	0
			43	34	1	4	4		
2	MMM	1	Total	C	Fe	N	O 1	0	0
			43	34		4	4		
2	MMM	1	Total	C	Fe	N	0	0	0
			43	$\frac{34}{C}$		4 	4		
2	MMM	1		0	re 1	IN 4	0	0	0
			43 Total	$\frac{34}{C}$		4 N	4		
2	YYY	1	10tai 42	24	ге 1	IN A		0	0
			40 Total	$\frac{54}{C}$		4 N	4		
2	YYY	1	10tai 43	24	те 1	1N - /		0	0
			Total	-04 	Fo	-4 N	- 		
2	YYY	1	10tai /13	34	те 1	1	1	0	0
			Total	01 C	Fe	N	<u>г</u> О		
2	YYY	1	43	34	1	4	$\frac{0}{4}$	0	0
			Total	$\frac{01}{C}$	Fe	N	$\overline{0}$		
2	YYY	1	43	34	1	4	4	0	0
			Total	\overline{C}	Fe	N	0		_
2	YYY	1	43	34	1	4	4	0	0
	.		Total	C	Fe	N	0		_
2	YYY	1	43	34	1	4	4	0	0
	.		Total	C	Fe	N	0		-
2	YYY	1	43	34	1	4	4	0	0



• Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	AAA	1	Total Ca 1 1	0	0
3	MMM	1	Total Ca 1 1	0	0
3	YYY	1	Total Ca 1 1	0	0

• Molecule 4 is PHOSPHATE ION (three-letter code: PO4) (formula: O_4P) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	AAA	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{P} \\ 5 & 4 & 1 \end{array}$	0	0
4	MMM	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{P} \\ 5 & 4 & 1 \end{array}$	0	0
4	YYY	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{P} \\ 5 & 4 & 1 \end{array}$	0	0

• Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	AAA	323	Total O 323 323	0	0
5	MMM	308	Total O 308 308	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	YYY	319	Total O 319 319	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Octaheme nitrite reductase





M528 K531 Q532

• Molecule 1: Octaheme nitrite reductase

С	h	ai	n	Y	Y	Υ	:														6	589	%																		3	0%	, ว					•		
638	R39	E40 T41	C42	-	T47	0	K53 F54	H	K57	H58		161	A62	C63	104	H67	D68	K69	T70	K71	E72	H/3	K76		D79	<u>V80</u>	K81	P82 Vro 2		26N	86 <mark>0</mark>	66A	D112	OTTU	L125	-	E138	H145		M148	V149	V150		F153	-	R157	2011	R162	02 77	Y172
	T178	1181		R185		L191			C205	K206	S207	S208		Q213	W214 K015	H216	L217	G218	E219	K220		K223	r 224 D225	R226	-	N229	V230	N231	V24	V 203 K 235		G242		U240 H247	-	G260		12/3 F274	4	T280	D281	L282	K283	F287		F290	G294		E297	K298
T299	D300	8301 8302	M303	M304	C305		H309 V310	E311	Y312	N313	C314	N315		Q319	K320 T201	1321 D322		F328	A329	D330	A331	H332	1333 N334	1	P337		H347	Y348	6350 6350	1351	D352	F353	F354	T361	<mark>G362</mark>	A363	K364	T365	<mark>0369</mark>	H370		Y376	N386	2	0390	C391	0395		K399	D400
G401	K402 11102	K403 Y404	S405	T406	H407	G408	V409 V410	K411	-	0414	M415	V416	Q417	E418		0750	C423		K431	A432	K433	E427	E401	K444	G445	K446	M447	R448	U456	457 L457		F461	AA6A		S470	P471	T472	14/3 1474	D475	-	R478		E482	H485	A486	L487	w488 E489	Y490	W491	-
N495		A 505 R 506	Q507	S508		V520	1 500	L523	-	1527	M528	A529	V530	K531	7.59h																																			



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	114.20Å 64.80 Å 114.20 Å	Depositor
a, b, c, α , β , γ	90.00° 119.90° 90.00°	Depositor
$\mathbf{B}_{\mathrm{ascolution}}\left(\mathring{A}\right)$	42.88 - 1.74	Depositor
Resolution (A)	42.88 - 1.74	EDS
% Data completeness	97.8(42.88-1.74)	Depositor
(in resolution range)	97.8(42.88-1.74)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$1.53 (at 1.74 \text{\AA})$	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
B B c	0.191 , 0.254	Depositor
It, Itfree	0.198 , 0.264	DCC
R_{free} test set	7348 reflections (5.04%)	wwPDB-VP
Wilson B-factor $(Å^2)$	22.0	Xtriage
Anisotropy	0.333	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.28 , 14.8	EDS
L-test for $twinning^2$	$< L > = 0.51, < L^2 > = 0.35$	Xtriage
	0.457 for -h-l,k,h	
	0.457 for l,k,-h-l	
Estimated twinning fraction	0.023 for h,-k,-h-l	Xtriage
	0.029 for -h-l,-k,l	
	0.024 for l,-k,h	
	0.400 for H, K, L	
Reported twinning fraction	0.290 for L, K, -H-L	Depositor
	0.310 for -H-L, K, H	
Outliers	0 of 145920 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	13690	wwPDB-VP
Average B, all atoms $(Å^2)$	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.16% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: HEC, CA, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bo	nd lengths	Bond angles				
MIOI	Unain	RMSZ	# Z > 5	RMSZ	# Z > 5			
1	AAA	0.86	2/4002~(0.0%)	1.06	2/5390~(0.0%)			
1	MMM	0.88	4/4003~(0.1%)	1.05	3/5391~(0.1%)			
1	YYY	0.87	3/4000~(0.1%)	1.05	2/5388~(0.0%)			
All	All	0.87	9/12005~(0.1%)	1.05	7/16169~(0.0%)			

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	YYY	0	1

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms		Observed(Å)	Ideal(Å)
1	MMM	494	GLU	CD-OE1	7.06	1.33	1.25
1	YYY	423	CYS	CB-SG	-5.83	1.72	1.81
1	YYY	391	CYS	CB-SG	-5.64	1.72	1.81
1	AAA	451	GLU	CD-OE2	-5.60	1.19	1.25
1	MMM	423	CYS	CB-SG	-5.29	1.73	1.81

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
1	YYY	406	THR	CA-CB-OG1	-6.67	95.00	109.00
1	AAA	143	ARG	NE-CZ-NH1	-6.21	117.19	120.30
1	MMM	231	ASN	CB-CA-C	5.27	120.94	110.40
1	AAA	188	ASN	CB-CA-C	5.18	120.76	110.40
1	MMM	97	ASN	CB-CA-C	5.08	120.56	110.40



There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	YYY	310	VAL	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	3895	0	3804	124	0
1	MMM	3894	0	3811	104	0
1	YYY	3898	0	3812	122	0
2	AAA	347	0	214	39	0
2	MMM	344	0	240	31	0
2	YYY	344	0	240	31	0
3	AAA	1	0	0	0	0
3	MMM	1	0	0	0	0
3	YYY	1	0	0	0	0
4	AAA	5	0	0	1	0
4	MMM	5	0	0	0	0
4	YYY	5	0	0	2	0
5	AAA	323	0	0	14	1
5	MMM	308	0	0	8	0
5	YYY	319	0	0	13	1
All	All	13690	0	12121	374	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 374 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
$2 \cdot \Lambda \Lambda \Lambda \cdot 602 \cdot \text{HEC} \cdot \text{HBC}^2$	5· A A A · 1015· HOH·O	1.56	1.05
2.AAA.005.IIEC.IIDC5	J.AAA.1015.11011.0	1.50	1.05
1:YYY:319:GLN:NE2	1:YYY:322:ASP:OD2	1.98	0.97
1:MMM:456:GLN:HE21	1:MMM:456:GLN:HA	1.30	0.95
2:MMM:606:HEC:HBB3	2:MMM:606:HEC:HMB1	1.45	0.94
1:MMM:312:TYR:CE2	2:MMM:604:HEC:HMC2	2.08	0.89



All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:AAA:1019:HOH:O	5:YYY:1013:HOH:O[2_546]	2.12	0.08

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntile	\mathbf{s}
1	AAA	495/495~(100%)	459~(93%)	33~(7%)	3 (1%)	25	10	
1	MMM	496/495~(100%)	453 (91%)	41 (8%)	2 (0%)	34	17	
1	YYY	495/495~(100%)	448 (90%)	45 (9%)	2 (0%)	34	17	
All	All	1486/1485~(100%)	1360 (92%)	119 (8%)	7 (0%)	29	12	

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	AAA	138	GLU
1	AAA	241	VAL
1	AAA	282	LEU
1	YYY	475	ASP
1	MMM	241	VAL

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric Outliers		Percentiles		
1	AAA	416/414~(100%)	393~(94%)	23~(6%)	21	4	
1	MMM	415/414~(100%)	390~(94%)	25~(6%)	19	4	
1	YYY	416/414~(100%)	395~(95%)	21 (5%)	24	6	
All	All	1247/1242~(100%)	1178 (94%)	69~(6%)	22	4	

5 of 69 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	YYY	283	LYS
1	YYY	304	MET
1	YYY	489	GLU
1	MMM	108	ASP
1	MMM	85	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 31 ligands modelled in this entry, 3 are monoatomic - leaving 28 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mol	Type	Chain	Bos	Link	Bond lengths			Bond angles		
	Type	Chain	nes		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2
2	HEC	YYY	601	1	32,50,50	2.09	8 (25%)	24,82,82	3.26	11 (45%)
4	PO4	MMM	610	2	$4,\!4,\!4$	1.76	2 (50%)	$6,\!6,\!6$	1.27	0
2	HEC	AAA	605	1	32,50,50	2.07	12 (37%)	24,82,82	2.56	9 (37%)
2	HEC	MMM	606	1	32,50,50	2.13	10 (31%)	24,82,82	<mark>3.06</mark>	10 (41%)
2	HEC	YYY	603	1	32,50,50	2.11	10 (31%)	24,82,82	2.21	7 (29%)
2	HEC	MMM	607	1	32,50,50	1.76	9 (28%)	24,82,82	2.93	14 (58%)
2	HEC	YYY	607	1	32,50,50	1.94	8 (25%)	24,82,82	2.74	11 (45%)
4	PO4	YYY	610	2	4,4,4	0.45	0	6,6,6	0.91	0
2	HEC	MMM	608	1	32,50,50	1.92	8 (25%)	24,82,82	<mark>3.35</mark>	7 (29%)
2	HEC	AAA	602	1	32,50,50	1.99	9 (28%)	24,82,82	2.59	12 (50%)
2	HEC	AAA	607[A]	-	32,50,50	2.15	10 (31%)	24,82,82	<mark>3.11</mark>	12 (50%)
2	HEC	AAA	603	1	32,50,50	2.34	11 (34%)	24,82,82	2.57	7 (29%)
2	HEC	AAA	604	1,4	32,50,50	1.66	5 (15%)	24,82,82	1.57	4 (16%)
2	HEC	AAA	601	1	32,50,50	2.13	9 (28%)	24,82,82	3.45	11 (45%)
2	HEC	AAA	608	1	32,50,50	2.04	7 (21%)	24,82,82	3.18	7 (29%)
2	HEC	MMM	603	1	32,50,50	1.95	8 (25%)	24,82,82	3.02	8 (33%)
2	HEC	AAA	606	1	32,50,50	2.08	10 (31%)	24,82,82	2.60	10 (41%)
2	HEC	YYY	604	1,4	32,50,50	2.09	9 (28%)	24,82,82	2.06	8 (33%)
2	HEC	YYY	608	1	32,50,50	1.94	5 (15%)	24,82,82	<mark>2.93</mark>	7 (29%)
2	HEC	YYY	606	1	32,50,50	2.20	17 (53%)	24,82,82	2.91	9 (37%)
2	HEC	MMM	602	1	32,50,50	1.83	9 (28%)	24,82,82	2.03	8 (33%)
2	HEC	YYY	602	1	32,50,50	1.88	10 (31%)	24,82,82	2.08	9 (37%)
2	HEC	MMM	604	1,4	32,50,50	1.68	7 (21%)	24,82,82	2.21	7 (29%)
2	HEC	MMM	605	1	32,50,50	1.96	7 (21%)	24,82,82	2.52	7 (29%)
4	PO4	AAA	610	2	4,4,4	1.94	2(50%)	6,6,6	0.93	0
2	HEC	AAA	607[B]	-	32,50,50	2.13	10 (31%)	24,82,82	2.92	10 (41%)
2	HEC	YYY	605	1	32,50,50	1.94	10 (31%)	24,82,82	2.62	8 (33%)
2	HEC	MMM	601	1	32,50,50	2.24	9 (28%)	24,82,82	<mark>3.31</mark>	13 (54%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEC	YYY	601	1	-	7/10/54/54	-



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEC	AAA	605	1	-	2/10/54/54	-
2	HEC	MMM	606	1	-	0/10/54/54	-
2	HEC	YYY	603	1	-	4/10/54/54	-
2	HEC	MMM	607	1	-	5/10/54/54	-
2	HEC	YYY	607	1	-	3/10/54/54	-
2	HEC	MMM	608	1	-	3/10/54/54	-
2	HEC	AAA	602	1	-	2/10/54/54	-
2	HEC	AAA	607[A]	-	-	4/10/54/54	-
2	HEC	AAA	603	1	-	4/10/54/54	-
2	HEC	AAA	604	1,4	-	2/10/54/54	-
2	HEC	AAA	601	1	-	3/10/54/54	-
2	HEC	AAA	608	1	_	1/10/54/54	-
2	HEC	MMM	603	1	-	2/10/54/54	-
2	HEC	AAA	606	1	-	0/10/54/54	-
2	HEC	YYY	604	1,4	-	2/10/54/54	-
2	HEC	YYY	608	1	-	2/10/54/54	-
2	HEC	YYY	606	1	-	0/10/54/54	-
2	HEC	MMM	602	1	-	3/10/54/54	-
2	HEC	YYY	602	1	-	5/10/54/54	-
2	HEC	MMM	604	1,4	-	2/10/54/54	-
2	HEC	MMM	605	1	-	4/10/54/54	-
2	HEC	AAA	607[B]	-	_	5/10/54/54	-
2	HEC	YYY	605	1	-	2/10/54/54	-
2	HEC	MMM	601	1	-	2/10/54/54	-

Continued from previous page...

The worst 5 of 231 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	AAA	603	HEC	C3C-C2C	7.80	1.48	1.40
2	AAA	608	HEC	C3C-C2C	7.73	1.48	1.40
2	YYY	604	HEC	C2B-C3B	7.10	1.48	1.40
2	MMM	601	HEC	C3C-C2C	7.04	1.48	1.40
2	YYY	603	HEC	C3C-C2C	6.64	1.47	1.40

The worst 5 of 226 bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$\mathbf{Ideal}(^{o})$
2	AAA	608	HEC	CBA-CAA-C2A	-11.46	93.29	112.60
2	MMM	606	HEC	CMC-C2C-C3C	10.34	137.98	125.82
2	YYY	608	HEC	CBA-CAA-C2A	-10.19	95.43	112.60
2	MMM	608	HEC	CBA-CAA-C2A	-10.06	95.64	112.60
2	MMM	605	HEC	C1D-C2D-C3D	-9.54	100.36	107.00

There are no chirality outliers.

5 of 69 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	YYY	601	HEC	C2D-C3D-CAD-CBD
2	YYY	601	HEC	C4D-C3D-CAD-CBD
2	YYY	601	HEC	C3D-CAD-CBD-CGD
2	AAA	608	HEC	C3D-CAD-CBD-CGD
2	AAA	607[B]	HEC	C2A-CAA-CBA-CGA

There are no ring outliers.

26 monomers are involved in 104 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	YYY	601	HEC	10	0
2	AAA	605	HEC	7	0
2	MMM	606	HEC	5	0
2	YYY	603	HEC	1	0
2	MMM	607	HEC	8	0
2	YYY	607	HEC	4	0
4	YYY	610	PO4	2	0
2	MMM	608	HEC	1	0
2	AAA	602	HEC	8	0
2	AAA	607[A]	HEC	1	0
2	AAA	603	HEC	8	0
2	AAA	604	HEC	7	0
2	AAA	601	HEC	6	0
2	AAA	608	HEC	3	0
2	MMM	603	HEC	3	0
2	AAA	606	HEC	4	0
2	YYY	604	HEC	4	0
2	YYY	608	HEC	3	0
2	YYY	606	HEC	2	0
2	MMM	602	HEC	7	0
2	YYY	602	HEC	1	0
2	MMM	604	HEC	3	0



contract from proceeder pagem											
Mol	Chain	Res	Type	Clashes	Symm-Clashes						
2	MMM	605	HEC	3	0						
4	AAA	610	PO4	1	0						
2	YYY	605	HEC	6	0						
2	MMM	601	HEC	2	0						

Continued from previous page...

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.
















































































































5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ>2			$OWAB(Å^2)$	Q<0.9
1	AAA	495/495~(100%)	-0.33	3 (0%)	89	92	12, 22, 36, 55	0
1	MMM	495/495~(100%)	-0.29	3 (0%)	89	92	11, 21, 37, 69	0
1	YYY	495/495~(100%)	-0.32	1 (0%)	95	96	12, 22, 38, 53	0
All	All	1485/1485~(100%)	-0.31	7 (0%)	91	93	11, 22, 38, 69	0

The worst 5 of 7 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	AAA	530	VAL	3.7
1	MMM	401	GLY	2.8
1	AAA	529	ALA	2.3
1	MMM	400	ASP	2.3
1	MMM	468	GLY	2.2

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(Å^2)$	Q<0.9
2	HEC	YYY	602	43/43	0.94	0.10	17,24,27,29	0
2	HEC	AAA	602	43/43	0.95	0.09	19,22,25,32	0
2	HEC	MMM	602	43/43	0.95	0.10	16,21,23,26	0
2	HEC	YYY	601	43/43	0.95	0.10	20,25,31,40	0
2	HEC	AAA	601	43/43	0.95	0.10	16,23,32,38	0
4	PO4	MMM	610	5/5	0.95	0.08	19,21,23,24	0
2	HEC	MMM	603	43/43	0.96	0.10	24,26,35,42	0
2	HEC	MMM	605	43/43	0.96	0.09	14,21,29,30	0
2	HEC	AAA	605	43/43	0.96	0.09	17,22,25,27	0
2	HEC	MMM	601	43/43	0.96	0.10	22,26,30,39	0
2	HEC	YYY	603	43/43	0.96	0.10	20,27,29,31	0
2	HEC	AAA	603	43/43	0.96	0.10	22,28,32,33	0
4	PO4	YYY	610	5/5	0.96	0.07	16,20,23,23	0
2	HEC	AAA	604	43/43	0.97	0.07	10,15,18,19	0
2	HEC	AAA	607[A]	43/43	0.97	0.08	11,14,18,18	3
2	HEC	AAA	607[B]	43/43	0.97	0.08	11,14,18,18	3
2	HEC	YYY	604	43/43	0.97	0.08	12,16,20,23	0
2	HEC	YYY	605	43/43	0.97	0.08	14,22,27,36	0
2	HEC	YYY	607	43/43	0.97	0.08	13,16,20,22	0
2	HEC	YYY	608	43/43	0.97	0.09	14,22,28,31	0
4	PO4	AAA	610	5/5	0.97	0.06	17,18,20,21	0
2	HEC	AAA	608	43/43	0.97	0.08	13,18,27,34	0
2	HEC	MMM	608	43/43	0.97	0.09	13,17,27,35	0
2	HEC	MMM	606	43/43	0.98	0.07	11,14,16,21	0
2	HEC	MMM	607	43/43	0.98	0.07	12,15,20,23	0
2	HEC	MMM	604	43/43	0.98	0.07	11,15,17,22	0
2	HEC	AAA	606	43/43	0.98	0.07	12,15,18,21	0
2	HEC	YYY	606	43/43	0.98	0.07	13,16,18,20	0
3	CA	AAA	609	1/1	0.99	0.07	21,21,21,21	0
3	CA	MMM	609	1/1	0.99	0.04	21,21,21,21	0
3	CA	YYY	609	1/1	0.99	0.04	18,18,18,18	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




























































































































6.5 Other polymers (i)

There are no such residues in this entry.

