



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 27, 2024 – 07:15 pm BST

PDB ID : 8RYN
Title : Structure of S2 TCR in complex with HLA-A*11:01 bound to ELFSYLIEK peptide
Authors : Karuppiah, V.
Deposited on : 2024-02-09
Resolution : 1.97 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.002 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.38.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 1.97 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1356 (1.98-1.98)
Clashscore	180529	1437 (1.98-1.98)
Ramachandran outliers	177936	1426 (1.98-1.98)
Sidechain outliers	177891	1426 (1.98-1.98)
RSRZ outliers	164620	1356 (1.98-1.98)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	276	
2	B	100	
3	C	9	
4	D	198	
5	E	245	

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 6767 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MHC class I antigen.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	275	2244	1394	408	433	9	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	276	PRO	-	expression tag	UNP A0A583ZB34

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	97	812	517	138	155	2	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	MET	-	initiating methionine	UNP P61769

- Molecule 3 is a protein called ELFSYLIEK peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	C	9	81	55	10	16	0	0	0

- Molecule 4 is a protein called TCR alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	D	186	1411	867	246	290	8	0	0	0

- Molecule 5 is a protein called TCR beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
5	E	242	1918	1210	329	370	9	0	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	104	Total 104	O 104	0	0
6	B	53	Total 53	O 53	0	0
6	C	8	Total 8	O 8	0	0
6	D	30	Total 30	O 30	0	0
6	E	106	Total 106	O 106	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: MHC class I antigen




- Molecule 2: Beta-2-microglobulin



- Molecule 3: ELFSYLIEK peptide




- Molecule 4: TCR alpha



- Molecule 5: TCR beta





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	86.75Å 48.69Å 115.30Å 90.00° 106.39° 90.00°	Depositor
Resolution (Å)	77.90 – 1.97 77.90 – 1.97	Depositor EDS
% Data completeness (in resolution range)	99.9 (77.90-1.97) 99.9 (77.90-1.97)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.60 (at 1.97Å)	Xtrriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.220 , 0.251 0.229 , 0.258	Depositor DCC
R_{free} test set	3242 reflections (4.92%)	wwPDB-VP
Wilson B-factor (Å ²)	34.1	Xtrriage
Anisotropy	0.557	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 33.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6767	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.51% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.63	0/2305	0.71	0/3129
2	B	0.61	0/835	0.70	0/1131
3	C	0.66	0/82	0.65	0/107
4	D	0.66	0/1432	0.73	0/1938
5	E	0.63	0/1971	0.72	0/2685
All	All	0.63	0/6625	0.71	0/8990

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2244	0	2091	4	0
2	B	812	0	771	3	0
3	C	81	0	83	1	0
4	D	1411	0	1356	1	0
5	E	1918	0	1820	0	0
6	A	104	0	0	0	0
6	B	53	0	0	0	0
6	C	8	0	0	0	0
6	D	30	0	0	0	0
6	E	106	0	0	0	0
All	All	6767	0	6121	6	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All (6) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:96:GLN:OE1	2:B:31:HIS:HE1	2.02	0.43
1:A:182:THR:HG23	1:A:182:THR:O	2.18	0.42
2:B:51:HIS:HA	2:B:65:LEU:O	2.19	0.42
1:A:117:ALA:HB2	2:B:60:TRP:CE2	2.54	0.42
1:A:45:MET:CE	3:C:2:LEU:HD11	2.50	0.41
4:D:162:LEU:HB3	4:D:171:SER:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	273/276 (99%)	271 (99%)	2 (1%)	0	100	100
2	B	95/100 (95%)	94 (99%)	1 (1%)	0	100	100
3	C	7/9 (78%)	7 (100%)	0	0	100	100
4	D	180/198 (91%)	173 (96%)	7 (4%)	0	100	100
5	E	240/245 (98%)	238 (99%)	2 (1%)	0	100	100
All	All	795/828 (96%)	783 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	232/233 (100%)	231 (100%)	1 (0%)	89	89
2	B	92/95 (97%)	91 (99%)	1 (1%)	70	68
3	C	9/9 (100%)	9 (100%)	0	100	100
4	D	160/172 (93%)	159 (99%)	1 (1%)	84	84
5	E	209/211 (99%)	208 (100%)	1 (0%)	86	86
All	All	702/720 (98%)	698 (99%)	4 (1%)	84	84

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	116	ASP
2	B	70	PHE
4	D	21	VAL
5	E	154	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	87	GLN
1	A	127	ASN
1	A	155	GLN
1	A	188	HIS
2	B	31	HIS
4	D	58	GLN
5	E	221	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	275/276 (99%)	0.70	32 (11%) 11 16	25, 40, 72, 83	0
2	B	97/100 (97%)	0.35	3 (3%) 51 62	25, 35, 54, 69	0
3	C	9/9 (100%)	-0.15	0 100 100	29, 30, 36, 36	0
4	D	186/198 (93%)	1.55	61 (32%) 1 1	30, 52, 75, 89	0
5	E	242/245 (98%)	0.43	17 (7%) 24 33	26, 36, 57, 78	0
All	All	809/828 (97%)	0.77	113 (13%) 7 11	25, 40, 72, 89	0

All (113) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	D	80	ALA	5.3
1	A	194	ILE	4.6
4	D	61	GLY	4.5
4	D	162	LEU	4.0
4	D	54	SER	4.0
4	D	17	GLY	3.8
1	A	267	PRO	3.7
4	D	77	TYR	3.7
1	A	248	VAL	3.7
4	D	190	ASN	3.5
5	E	245	ASP	3.5
5	E	102	TYR	3.5
4	D	55	GLN	3.5
4	D	109	MET	3.4
4	D	59	THR	3.3
5	E	29	ASN	3.3
1	A	197	HIS	3.3
1	A	225	THR	3.3
4	D	79	ALA	3.3
4	D	66	SER	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	182	THR	3.2
4	D	75	THR	3.2
2	B	48	LYS	3.1
4	D	165	ARG	3.1
2	B	98	ASP	3.1
1	A	219	ARG	3.0
4	D	186	ALA	2.9
4	D	19	ASN	2.9
1	A	250	PRO	2.9
1	A	195	SER	2.9
4	D	56	ARG	2.9
4	D	129	SER	2.9
4	D	65	ALA	2.9
5	E	225	THR	2.8
4	D	107	ARG	2.8
4	D	49	LEU	2.8
4	D	63	LEU	2.8
4	D	185	CYS	2.8
5	E	241	TRP	2.8
1	A	220	ASP	2.8
4	D	163	ASP	2.8
5	E	155	HIS	2.7
1	A	221	GLY	2.7
1	A	230	LEU	2.7
4	D	68	ASP	2.7
1	A	256	ARG	2.7
1	A	226	GLN	2.7
4	D	20	LEU	2.7
4	D	147	SER	2.6
1	A	105	PRO	2.6
1	A	193	PRO	2.6
4	D	18	GLU	2.6
4	D	71	SER	2.6
5	E	226	GLN	2.6
1	A	269	PRO	2.6
4	D	191	ASN	2.6
4	D	164	MET	2.6
4	D	15	PRO	2.6
4	D	192	SER	2.6
1	A	199	ALA	2.6
5	E	27	ASP	2.6
4	D	110	VAL	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	274	TRP	2.5
4	D	67	LEU	2.5
4	D	160	CYS	2.5
4	D	50	LEU	2.5
1	A	1	GLY	2.5
4	D	113	HIS	2.5
4	D	144	THR	2.5
4	D	64	ASN	2.5
1	A	249	VAL	2.5
4	D	21	VAL	2.4
4	D	81	SER	2.4
4	D	189	PHE	2.4
1	A	200	THR	2.4
4	D	62	ARG	2.4
4	D	148	GLN	2.4
1	A	251	SER	2.3
4	D	83	PRO	2.3
1	A	17	ARG	2.3
4	D	57	GLU	2.3
5	E	208	HIS	2.3
2	B	2	GLN	2.3
4	D	143	GLN	2.3
1	A	223	ASP	2.3
4	D	51	ILE	2.2
5	E	183	ALA	2.2
4	D	69	LYS	2.2
1	A	128	GLU	2.2
4	D	14	VAL	2.2
1	A	257	TYR	2.2
4	D	78	ILE	2.2
5	E	4	GLY	2.2
1	A	86	ASN	2.2
1	A	255	GLN	2.2
1	A	138	MET	2.2
4	D	171	SER	2.2
4	D	70	SER	2.1
5	E	220	GLU	2.1
5	E	112	LEU	2.1
4	D	16	GLU	2.1
4	D	179	ASN	2.1
5	E	221	ASN	2.1
1	A	252	GLY	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	192	HIS	2.1
4	D	2	ALA	2.1
4	D	184	ALA	2.1
4	D	187	ASN	2.1
4	D	48	LEU	2.0
4	D	82	GLN	2.0
5	E	239	GLU	2.0
5	E	244	ALA	2.0
5	E	184	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.