



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 22, 2024 – 08:12 PM EDT

PDB ID : 6RZ3
Title : Crystal structure of a complex between the DNA-binding domain of p53 and the carboxyl-terminal conserved region of iASPP
Authors : Chen, S.; Ren, J.; Jones, E.Y.; Lu, X.
Deposited on : 2019-06-12
Resolution : 4.23 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.20.1
EDS : 2.37.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 4.23 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1015 (4.70-3.78)
Clashscore	141614	1051 (4.68-3.80)
Ramachandran outliers	138981	1007 (4.68-3.80)
Sidechain outliers	138945	1016 (4.70-3.78)
RSRZ outliers	127900	1065 (4.76-3.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	232	
2	B	238	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2867 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cellular tumor antigen p53.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	201	1584	979	293	296	16	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	61	MET	-	initiating methionine	UNP P04637

- Molecule 2 is a protein called RelA-associated inhibitor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	167	1282	808	213	252	9	0	0	0

There are 34 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	624	MET	-	initiating methionine	UNP Q8WUF5
B	829	LEU	-	expression tag	UNP Q8WUF5
B	830	GLU	-	expression tag	UNP Q8WUF5
B	831	GLY	-	expression tag	UNP Q8WUF5
B	832	GLY	-	expression tag	UNP Q8WUF5
B	833	SER	-	expression tag	UNP Q8WUF5
B	834	GLY	-	expression tag	UNP Q8WUF5
B	835	GLY	-	expression tag	UNP Q8WUF5
B	836	SER	-	expression tag	UNP Q8WUF5
B	837	GLY	-	expression tag	UNP Q8WUF5
B	838	LEU	-	expression tag	UNP Q8WUF5
B	839	ASN	-	expression tag	UNP Q8WUF5
B	840	ASP	-	expression tag	UNP Q8WUF5
B	841	ILE	-	expression tag	UNP Q8WUF5
B	842	PHE	-	expression tag	UNP Q8WUF5

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	843	GLU	-	expression tag	UNP Q8WUF5
B	844	ALA	-	expression tag	UNP Q8WUF5
B	845	GLN	-	expression tag	UNP Q8WUF5
B	846	LYS	-	expression tag	UNP Q8WUF5
B	847	ILE	-	expression tag	UNP Q8WUF5
B	848	GLU	-	expression tag	UNP Q8WUF5
B	849	TRP	-	expression tag	UNP Q8WUF5
B	850	HIS	-	expression tag	UNP Q8WUF5
B	851	GLU	-	expression tag	UNP Q8WUF5
B	852	GLY	-	expression tag	UNP Q8WUF5
B	853	ARG	-	expression tag	UNP Q8WUF5
B	854	THR	-	expression tag	UNP Q8WUF5
B	855	LYS	-	expression tag	UNP Q8WUF5
B	856	HIS	-	expression tag	UNP Q8WUF5
B	857	HIS	-	expression tag	UNP Q8WUF5
B	858	HIS	-	expression tag	UNP Q8WUF5
B	859	HIS	-	expression tag	UNP Q8WUF5
B	860	HIS	-	expression tag	UNP Q8WUF5
B	861	HIS	-	expression tag	UNP Q8WUF5


- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Zn 1 1	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Cellular tumor antigen p53


Chain A: 



LYS

- Molecule 2: RelA-associated inhibitor

Chain B: 



D768
Y769
L777
R780
A801
A802
L803
Q806
E807
L817
F818
R823
GLN
ARG
SER
LYS
VAL
LEU
GLY
GLY
GLY
SER
GLY
SER
GLY
GLY
LEU
ASN
ASP
ILE
PHE
GLU
ALA
GLN
LYS
ILE
GLU
TRP
HIS
GLU
GLY
ARG
THR
LYS
HIS
HIS
HIS
HIS
HIS

4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	71.09Å 71.09Å 255.59Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.80 – 4.23 29.79 – 4.23	Depositor EDS
% Data completeness (in resolution range)	93.2 (29.80-4.23) 93.6 (29.79-4.23)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.66 (at 4.26Å)	Xtrriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.243 , 0.292 0.255 , 0.313	Depositor DCC
R_{free} test set	473 reflections (9.81%)	wwPDB-VP
Wilson B-factor (Å ²)	160.5	Xtrriage
Anisotropy	0.535	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 124.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.42$, $\langle L^2 \rangle = 0.25$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	2867	wwPDB-VP
Average B, all atoms (Å ²)	198.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.51% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section:
ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.37	0/1622	0.60	0/2200
2	B	0.40	0/1318	0.60	0/1800
All	All	0.39	0/2940	0.60	0/4000

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1584	0	1544	10	0
2	B	1282	0	1175	17	0
3	A	1	0	0	0	0
All	All	2867	0	2719	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:766:LEU:O	2:B:767:TRP:HD1	1.40	1.05
2:B:766:LEU:O	2:B:767:TRP:CD1	2.23	0.90
1:A:123:THR:O	1:A:278:PRO:HG3	1.94	0.68
2:B:687:ASN:HD21	2:B:718:ALA:H	1.43	0.66
1:A:241:SER:HB2	1:A:248:ARG:NH2	2.20	0.56
2:B:693:GLY:HA3	2:B:724:LEU:HG	1.88	0.56
1:A:283:ARG:CZ	2:B:673:TYR:HB2	2.40	0.52
1:A:114:LEU:HG	1:A:115:HIS:N	2.25	0.52
1:A:117:GLY:HA2	1:A:123:THR:H	1.75	0.52
2:B:766:LEU:C	2:B:767:TRP:CD1	2.87	0.48
2:B:769:TYR:O	2:B:780:ARG:NH2	2.48	0.47
2:B:755:MET:HG2	2:B:817:LEU:HD13	1.97	0.47
1:A:241:SER:HB2	1:A:248:ARG:CZ	2.44	0.47
1:A:283:ARG:NH2	2:B:673:TYR:HB2	2.30	0.46
1:A:100:GLN:N	1:A:100:GLN:OE1	2.49	0.45
1:A:283:ARG:HG3	2:B:707:VAL:HG11	1.98	0.45
2:B:672:ASN:HB3	2:B:675:ILE:HD13	1.99	0.44
2:B:802:ALA:HA	2:B:806:GLN:O	2.17	0.44
2:B:801:ALA:O	2:B:807:GLU:HA	2.20	0.42
1:A:98:PRO:HG2	1:A:162:ILE:HG21	2.01	0.42
2:B:777:LEU:HD21	2:B:803:LEU:HB3	2.02	0.41
2:B:817:LEU:HD12	2:B:817:LEU:HA	1.95	0.41
2:B:714:GLN:HA	2:B:818:PHE:CD2	2.55	0.41
2:B:711:ALA:O	2:B:715:HIS:ND1	2.46	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	199/232 (86%)	194 (98%)	4 (2%)	1 (0%)	29 68
2	B	165/238 (69%)	160 (97%)	5 (3%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	364/470 (77%)	354 (97%)	9 (2%)	1 (0%)	41 76

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	117	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	181/201 (90%)	178 (98%)	3 (2%)	60 78
2	B	131/191 (69%)	131 (100%)	0	100 100
All	All	312/392 (80%)	309 (99%)	3 (1%)	76 86

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	110	ARG
1	A	243	MET
1	A	287	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	235	ASN
2	B	687	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	201/232 (86%)	-0.20	0 100 100	147, 206, 248, 285	0
2	B	167/238 (70%)	-0.31	2 (1%) 79 70	145, 179, 240, 340	0
All	All	368/470 (78%)	-0.25	2 (0%) 91 86	145, 195, 248, 340	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	657	ASN	3.1
2	B	658	GLU	2.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	ZN	A	301	1/1	0.94	0.08	210,210,210,210	0

6.5 Other polymers [i](#)

There are no such residues in this entry.