

Jul 3, 2024 - 06:18 am BST

PDB ID 6S0K : EMDB ID EMD-10073 : Title : Ribosome nascent chain in complex with SecA Authors Jomaa, A.; Wang, S.; Shan, S.; Ban, N. : Deposited on 2019-06-17 : 3.10 Å(reported) Resolution : Based on initial model 5GAG :

This is a Full wwPDB EM Validation Report for a publicly released PDB entry. We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at

https://www.wwpdb.org/validation/2017/EMValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

:	0.0.1. dev 92
:	4.02b-467
:	20191225.v01 (using entries in the PDB archive December 25th 2019)
:	1.9.13
:	Engh & Huber (2001)
:	Parkinson et al. (1996)
:	2.37.1
	: : : : :

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $ELECTRON\ MICROSCOPY$

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f EM} {f structures} \ (\#{f Entries})$		
Ramachandran outliers	154571	4023		
Sidechain outliers	154315	3826		
RNA backbone	4643	859		

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for $\geq=3, 2, 1$ and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq=5\%$ The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion < 40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	2	3	67% 33	3%
2	А	2883	83%	17% ·
3	В	120	84%	16%
4	С	273	99%	
5	D	209	100%	
6	Е	201	100%	
7	F	179	98%	••
8	G	177	99%	



Mol	Chain	Length	Quality of chain	
9	Н	149	25% 99%	— .
10	Ι	165	65% • 24%	_
11	J	142	94%	6%
12	K	142	100%	-
13	L	123	100%	
14	М	144	• 100%	-
15	Ν	136	100%	-
16	0	127	98%	••
17	Р	117	100%	_
18	Q	115	99%	•
19	R	118	99%	
20	S	103	99%	- .
21	Т	110	100%	_
22	U	100	• 92%	%
23	V	104	99%	
24	W	94	100%	-
25	Х	85	89% 11%)
26	Y	78	99%	
27	Z	63	98%	•
28	a	59	• 97%	•••
29	b	57	98%	•
30	с	55	/5% 91% • 79	%
31	d	46	- 100%	
32	е	65	94% 59	%•
33	f	38	100%	-



Mol	Chain	Length	Quality of chain
			61%
34	h	838	97% •
			53%
35	k	57	100%



2 Entry composition (i)

There are 37 unique types of molecules in this entry. The entry contains 98708 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a RNA chain called tRNA-CCA.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	2	3	Total 62	C 28	N 11	O 20	Р 3	0	0

• Molecule 2 is a RNA chain called 23S ribosomal RNA.

Mol	Chain	Residues			AltConf	Trace			
2	А	2883	Total 61902	C 27613	N 11397	O 20009	Р 2883	0	0

• Molecule 3 is a RNA chain called 5S ribosomal RNA.

Mol	Chain	Residues		At	AltConf	Trace			
3	В	120	Total 2569	C 1144	N 468	0 837	Р 120	0	0

• Molecule 4 is a protein called 50S ribosomal protein L2.

Mol	Chain	Residues		At	AltConf	Trace			
4	С	272	Total 2083	C 1288	N 424	0 364	${ m S} 7$	0	1

• Molecule 5 is a protein called 50S ribosomal protein L3.

Mol	Chain	Residues		At	oms	AltConf	Trace		
5	D	209	Total 1565	C 979	N 288	0 294	${S \atop 4}$	0	0

• Molecule 6 is a protein called 50S ribosomal protein L4.

Mol	Chain	Residues		At	oms	AltConf	Trace		
6	Е	201	Total 1552	C 974	N 283	O 290	${ m S}{ m 5}$	0	0



• Molecule 7 is a protein called 50S ribosomal protein L5.

Mol	Chain	Residues		At	oms	AltConf	Trace		
7	F	178	Total 1411	C 899	N 250	O 256	S 6	0	1

• Molecule 8 is a protein called 50S ribosomal protein L6.

Mol	Chain	Residues		At	oms			AltConf	Trace
8	G	176	Total 1323	C 832	N 243	0 246	${ m S} { m 2}$	0	0

• Molecule 9 is a protein called 50S ribosomal protein L9.

Mol	Chain	Residues		At	oms			AltConf	Trace
9	Н	149	Total 1110	C 699	N 197	0 213	S 1	0	0

• Molecule 10 is a protein called 50S ribosomal protein L10.

Mol	Chain	Residues		At	oms			AltConf	Trace
10	Ι	125	Total 946	C 598	N 169	0 175	$\frac{S}{4}$	0	0

• Molecule 11 is a protein called 50S ribosomal protein L11.

Mol	Chain	Residues		At	oms			AltConf	Trace
11	J	134	Total 979	C 619	N 169	0 185	S 6	0	0

• Molecule 12 is a protein called 50S ribosomal protein L13.

Mol	Chain	Residues		At	oms		AltConf	Trace	
12	K	142	Total 1129	C 714	N 212	0 199	${S \atop 4}$	0	0

• Molecule 13 is a protein called 50S ribosomal protein L14.

Mol	Chain	Residues		At	oms			AltConf	Trace
13	L	123	Total 946	C 593	N 181	0 166	${f S}{f 6}$	0	0

• Molecule 14 is a protein called 50S ribosomal protein L15.



Mol	Chain	Residues		At	oms			AltConf	Trace
14	М	144	Total 1053	C 654	N 207	O 190	${ m S} { m 2}$	0	0

• Molecule 15 is a protein called 50S ribosomal protein L16.

Mol	Chain	Residues		At	oms			AltConf	Trace
15	Ν	136	Total 1074	C 686	N 205	0 177	S 6	0	0

• Molecule 16 is a protein called 50S ribosomal protein L17.

Mol	Chain	Residues		At	oms			AltConf	Trace
16	Ο	125	Total 993	C 613	N 202	0 173	${ m S}{ m 5}$	0	0

• Molecule 17 is a protein called 50S ribosomal protein L18.

Mol	Chain	Residues		At	oms			AltConf	Trace
17	Р	117	Total 900	C 557	N 179	0 163	S 1	0	0

• Molecule 18 is a protein called 50S ribosomal protein L19.

Mol	Chain	Residues		At	oms	AltConf	Trace		
18	Q	114	Total 917	C 574	N 179	O 163	S 1	0	0

• Molecule 19 is a protein called 50S ribosomal protein L20.

Mol	Chain	Residues		Ato	ms		AltConf	Trace
19	R	117	Total	С	Ν	0	0	0
			947	604	192	151	, i i i i i i i i i i i i i i i i i i i	Ŭ

• Molecule 20 is a protein called 50S ribosomal protein L21.

Mol	Chain	Residues		At	oms			AltConf	Trace
20	S	103	Total 816	C 516	N 153	0 145	${S \over 2}$	0	0

• Molecule 21 is a protein called 50S ribosomal protein L22.



Mol	Chain	Residues		At	oms			AltConf	Trace
21	Т	110	Total 857	C 532	N 166	O 156	${ m S} { m 3}$	0	0

• Molecule 22 is a protein called 50S ribosomal protein L23.

Mol	Chain	Residues		At	oms	AltConf	Trace		
22	U	92	Total 730	C 461	N 138	O 130	S 1	0	0

• Molecule 23 is a protein called 50S ribosomal protein L24.

Mol	Chain	Residues		Ato	ms	AltConf	Trace	
23	V	103	Total 780	C 492	N 147	0 141	0	1

• Molecule 24 is a protein called 50S ribosomal protein L25.

Mol	Chain	Residues		At	\mathbf{oms}			AltConf	Trace
24	W	94	Total 753	C 479	N 137	0 134	${ m S} { m 3}$	0	0

• Molecule 25 is a protein called 50S ribosomal protein L27.

Mol	Chain	Residues		At	oms			AltConf	Trace
25	Х	76	Total 580	C 359	N 117	0 103	S 1	0	0

• Molecule 26 is a protein called 50S ribosomal protein L28.

Mol	Chain	Residues		At	oms			AltConf	Trace
26	Y	77	Total 625	C 388	N 129	O 106	${ m S} { m 2}$	0	0

• Molecule 27 is a protein called 50S ribosomal protein L29.

Mol	Chain	Residues		Ato	\mathbf{ms}	AltConf	Trace		
27	Z	62	Total 501	C 308	N 98	0 94	S 1	0	0

• Molecule 28 is a protein called 50S ribosomal protein L30.



	$T \rightarrow 1$	a		-			
28 a 58	Total 440	C 281	N 87	0 70	S 2	0	0

• Molecule 29 is a protein called 50S ribosomal protein L32.

Mol	Chain	Residues		Ato	\mathbf{ms}	AltConf	Trace		
29	b	56	Total	С	N	0	S	0	0
			444	269	94	80	T		

• Molecule 30 is a protein called 50S ribosomal protein L33.

Mol	Chain	Residues		Aton	ıs	AltConf	Trace	
30	с	51	Total 414	C 266	N 76	O 72	0	0

• Molecule 31 is a protein called 50S ribosomal protein L34.

Mol	Chain	Residues		Ato	\mathbf{ms}	AltConf	Trace		
31	d	46	Total 377	C 228	N 90	O 57	${ m S} { m 2}$	0	0

• Molecule 32 is a protein called 50S ribosomal protein L35.

Mol	Chain	Residues	Atoms				AltConf	Trace	
32	е	64	Total 504	C 323	N 105	0 74	$\begin{array}{c} \mathrm{S} \\ \mathrm{2} \end{array}$	0	0

• Molecule 33 is a protein called 50S ribosomal protein L36.

Mol	Chain	Residues	Atoms			AltConf	Trace		
33	f	38	Total	С	Ν	Ο	S	0	0
00	1	30	302	185	65	48	4	0	0

• Molecule 34 is a protein called Protein translocase subunit SecA.

Mol	Chain	Residues	Atoms					AltConf	Trace
34	h	838	Total 6672	C 4181	N 1181	O 1276	S 34	0	0

There are 5 discrepancies between the modelled and reference sequences:



Chain	Residue	Modelled	Actual	Comment	Reference
h	-3	GLN	-	expression tag	UNP A0A037YQ84
h	-2	GLY	-	expression tag	UNP A0A037YQ84
h	-1	HIS	-	expression tag	UNP A0A037YQ84
h	0	MET	-	expression tag	UNP A0A037YQ84
h	12	CYS	SER	conflict	UNP A0A037YQ84

• Molecule 35 is a protein called Cytoskeleton protein RodZ.

Mol	Chain	Residues	Atoms			AltConf	Trace		
35	k	57	Total 374	C 247	N 63	O 63	S 1	0	0

• Molecule 36 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	AltConf
36	А	65	Total Mg 65 65	0
36	В	2	Total Mg 2 2	0
36	b	1	Total Mg 1 1	0

• Molecule 37 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	AltConf
37	f	1	Total Zn 1 1	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: tRNA-CCA













 \bullet Molecule 12: 50S ribosomal protein L13

Chain K:	
There are no outlier residues recorded for this chain.	
• Molecule 13: 50S ribosomal protein L14	
Chain L: 100%	
There are no outlier residues recorded for this chain.	
• Molecule 14: 50S ribosomal protein L15	
Chain M:	
\bullet Molecule 15: 50S ribosomal protein L16	
Chain N: 100%	
There are no outlier residues recorded for this chain.	
\bullet Molecule 16: 50S ribosomal protein L17	
Chain O: 98%	
M1 T57 A122 A122 A124 A126 GUU	
• Molecule 17: 50S ribosomal protein L18	
Chain P: 100%	
• Molecule 18: 50S ribosomal protein L19	
Chain Q: 99%	
MET S2 MI16 MI16	



• Molecule 19: 50S ribosomal protein L20
Chain R: 99%
A118
\bullet Molecule 20: 50S ribosomal protein L21
Chain S: 99%
\bullet Molecule 21: 50S ribosomal protein L22
Chain T:
\bullet Molecule 22: 50S ribosomal protein L23
Chain U: 92% 8%
MET 12 193 BHE GLY GLY GLY GLY
\bullet Molecule 23: 50S ribosomal protein L24
Chain V: 99%
MET K104
• Molecule 24: 50S ribosomal protein L25
Chain W: 100%
There are no outlier residues recorded for this chain.
\bullet Molecule 25: 50S ribosomal protein L27
Chain X: 89% 11%
MET ALA LYS LYS GLY SER FIG E85



\bullet Molecule 26: 50S ribosomal protein L28	
Chain Y: 99% .	
\bullet Molecule 27: 50S ribosomal protein L29	
Chain Z: 98% ·	
MET 46.3 A6.3	
\bullet Molecule 28: 50S ribosomal protein L30	
Chain a: 97% ···	
MET 14 165 165	
\bullet Molecule 29: 50S ribosomal protein L32	
Chain b: 98% ·	
\bullet Molecule 30: 50S ribosomal protein L33	
75% Chain c: 91% • 7%	
MET LYS LYS R6 E7 R6 R6 R5 R5 R5 R11 V12 R11 R12 R12 R12 R12 R12 R12 R12 R12 R	LYS
\bullet Molecule 31: 50S ribosomal protein L34	
Chain d: 100%	
\bullet Molecule 32: 50S ribosomal protein L35	
Chain e: 94% 5% •	





• Molecule 33: 50S ribosomal protein L36







• Molecule 35: Cytoskeleton protein RodZ





4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	37334	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE	Depositor
	CORRECTION	
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{\AA}^2)$	40	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	FEI FALCON III $(4k \ge 4k)$	Depositor
Maximum map value	6.126	Depositor
Minimum map value	-2.568	Depositor
Average map value	0.015	Depositor
Map value standard deviation	0.411	Depositor
Recommended contour level	0.82	Depositor
Map size (Å)	444.8, 444.8, 444.8	wwPDB
Map dimensions	320, 320, 320	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.39, 1.39, 1.39	Depositor



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		Bond	lengths	Bond angles		
	Ullalli	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	2	0.42	0/68	1.13	0/103	
2	А	0.57	0/69329	1.00	144/108152~(0.1%)	
3	В	0.49	0/2872	1.02	8/4478~(0.2%)	
4	С	0.35	0/2122	0.53	0/2854	
5	D	0.35	0/1586	0.51	0/2134	
6	Ε	0.33	0/1571	0.47	0/2113	
7	F	0.29	0/1435	0.50	1/1928~(0.1%)	
8	G	0.30	0/1343	0.49	0/1816	
9	Н	0.30	0/1121	0.53	0/1515	
10	Ι	0.32	0/958	0.65	0/1290	
11	J	0.31	0/993	0.60	0/1341	
12	Κ	0.35	0/1152	0.47	0/1551	
13	L	0.34	0/955	0.54	0/1279	
14	М	0.32	0/1062	0.52	0/1413	
15	Ν	0.34	0/1093	0.51	0/1460	
16	0	0.33	0/1006	0.51	0/1345	
17	Р	0.31	0/910	0.48	0/1219	
18	Q	0.34	0/929	0.47	0/1242	
19	R	0.34	0/960	0.41	0/1278	
20	\mathbf{S}	0.35	0/829	0.55	0/1107	
21	Т	0.32	0/864	0.51	0/1156	
22	U	0.34	0/736	0.51	0/984	
23	V	0.33	0/788	0.51	0/1053	
24	W	0.31	0/766	0.49	0/1025	
25	Х	0.34	0/587	0.46	0/776	
26	Y	0.34	0/635	0.51	0/848	
27	Ζ	0.30	0/502	0.45	0/667	
28	a	0.31	0/453	0.45	0/605	
29	b	0.32	0/450	0.47	0/599	
30	с	0.29	0/421	0.51	$0/\overline{561}$	
31	d	0.32	0/380	0.51	0/498	
32	е	0.33	0/513	0.62	0/676	



Mal	Chain	Bond lengths		Bond angles		
		RMSZ	# Z > 5	RMSZ	# Z > 5	
33	f	0.33	0/303	0.47	0/397	
34	h	0.31	0/6782	0.60	3/9146~(0.0%)	
35	k	0.27	0/384	0.58	0/531	
All	All	0.50	0/106858	0.89	156/159140~(0.1%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
10	Ι	0	2
32	е	0	1
34	h	0	3
All	All	0	6

There are no bond length outliers.

All (156) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	А	1313	U	C2-N1-C1'	10.24	129.99	117.70
2	А	1313	U	N1-C2-O2	9.89	129.72	122.80
2	А	1533	С	C2-N1-C1'	9.56	129.32	118.80
2	А	1313	U	N3-C2-O2	-9.54	115.52	122.20
2	А	1314	С	C2-N1-C1'	8.66	128.33	118.80
2	А	1348	С	N1-C2-O2	8.54	124.03	118.90
2	А	837	С	N3-C2-O2	-7.70	116.51	121.90
2	А	1956	U	N1-C2-O2	7.67	128.17	122.80
2	А	837	С	N1-C2-O2	7.66	123.50	118.90
2	А	1774	С	N3-C2-O2	-7.66	116.54	121.90
2	А	1314	С	C6-N1-C2	-7.61	117.25	120.30
2	А	1956	U	N3-C2-O2	-7.54	116.92	122.20
2	А	897	С	N1-C2-O2	7.49	123.39	118.90
2	А	234	U	N3-C2-O2	-7.12	117.22	122.20
3	В	31	С	C2-N1-C1'	7.11	126.62	118.80
2	А	1348	С	N3-C2-O2	-7.09	116.93	121.90
2	А	2666	С	N1-C2-O2	7.09	123.16	118.90
2	А	2430	А	C2-N3-C4	6.98	114.09	110.60
2	A	1314	С	C5-C6-N1	6.89	124.44	121.00
2	A	1313	U	C6-N1-C1'	-6.86	111.59	121.20
2	А	1774	С	N1-C2-O2	6.86	123.01	118.90
2	А	2617	U	N3-C2-O2	-6.86	117.40	122.20



Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
3	В	31	С	C6-N1-C2	-6.85	117.56	120.30
2	А	607	U	N3-C2-O2	-6.71	117.50	122.20
2	А	2063	С	C6-N1-C2	-6.67	117.63	120.30
2	А	2063	С	C2-N1-C1'	6.64	126.10	118.80
2	А	1533	С	C6-N1-C2	-6.61	117.66	120.30
2	А	1060	U	N3-C4-O4	6.57	124.00	119.40
2	А	1533	С	C6-N1-C1'	-6.54	112.95	120.80
2	А	1101	U	N1-C2-O2	6.54	127.38	122.80
2	А	1533	С	N1-C2-O2	6.53	122.82	118.90
2	А	2063	С	N1-C2-O2	6.50	122.80	118.90
2	А	353	С	N1-C2-O2	6.50	122.80	118.90
2	А	613	А	P-O3'-C3'	6.48	127.48	119.70
2	А	897	С	C6-N1-C2	-6.48	117.71	120.30
2	А	1060	U	C5-C4-O4	-6.46	122.02	125.90
2	А	901	С	N1-C2-O2	6.42	122.75	118.90
2	А	1072	С	C6-N1-C2	-6.34	117.76	120.30
2	А	897	С	C2-N1-C1'	6.31	125.74	118.80
2	А	1294	U	N3-C2-O2	-6.27	117.81	122.20
2	А	1101	U	N3-C2-O2	-6.27	117.81	122.20
2	А	2395	С	C5-C6-N1	6.24	124.12	121.00
3	В	30	С	C6-N1-C2	-6.21	117.81	120.30
2	А	1489	С	N3-C2-O2	-6.20	117.56	121.90
2	А	1533	С	C5-C6-N1	6.17	124.08	121.00
2	А	2474	U	N1-C2-O2	6.14	127.09	122.80
2	А	985	С	C2-N1-C1'	6.13	125.54	118.80
2	А	114	U	C2-N1-C1'	6.12	125.04	117.70
2	А	2063	С	N3-C2-O2	-6.07	117.65	121.90
34	h	288	LYS	C-N-CA	6.06	136.85	121.70
2	А	2617	U	N1-C2-O2	6.06	127.04	122.80
2	А	867	С	N1-C2-O2	6.02	122.51	118.90
2	А	897	С	N3-C2-O2	-5.99	117.71	121.90
2	А	2666	С	N3-C2-O2	-5.98	117.71	121.90
2	А	2195	U	N1-C2-O2	5.98	126.98	122.80
2	А	2149	U	N1-C2-O2	5.97	126.98	122.80
2	А	1489	С	N1-C2-O2	5.97	122.48	118.90
2	А	1320	С	N3-C2-O2	-5.93	117.75	121.90
2	А	1348	С	C2-N1-C1'	5.92	125.32	118.80
2	А	1343	G	C4-N9-C1'	5.88	134.15	126.50
2	А	2158	А	P-O3'-C3'	5.84	126.71	119.70
2	А	370	G	O4'-C1'-N9	-5.80	103.56	108.20
2	А	919	U	N1-C2-O2	5.78	126.84	122.80

А

2

U

N3-C2-O2

2474

Continued on next page...

122.20

118.16



-5.78

Conti	nued from	n previo	ous page.	
Mol	Chain	Res	Type	Atoms
-			a	Q = 3 = . Q =

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	А	837	С	C6-N1-C2	-5.77	117.99	120.30
2	А	790	U	C2-N1-C1'	5.76	124.61	117.70
2	А	12	U	N3-C2-O2	-5.76	118.17	122.20
2	А	2149	U	C2-N1-C1'	5.75	124.60	117.70
2	А	2667	С	C6-N1-C2	-5.74	118.00	120.30
2	А	1267	U	N1-C2-O2	5.70	126.79	122.80
2	А	1294	U	N1-C2-O2	5.68	126.77	122.80
2	А	2602	А	P-O3'-C3'	5.67	126.51	119.70
3	В	26	С	N1-C2-O2	5.67	122.30	118.90
2	А	2656	U	N1-C2-O2	5.66	126.76	122.80
7	F	152	LEU	CA-CB-CG	5.66	128.31	115.30
2	А	2149	U	N3-C2-O2	-5.65	118.24	122.20
2	А	607	U	N1-C2-O2	5.64	126.75	122.80
2	А	2474	U	C2-N1-C1'	5.64	124.47	117.70
3	В	31	С	N3-C2-O2	-5.64	117.95	121.90
2	А	135	U	N1-C2-O2	5.63	126.74	122.80
2	А	234	U	N1-C2-O2	5.62	126.74	122.80
2	А	2248	С	C2-N1-C1'	5.62	124.98	118.80
2	А	1398	С	C2-N1-C1'	5.60	124.96	118.80
2	А	353	С	N3-C2-O2	-5.60	117.98	121.90
2	А	2195	U	N3-C2-O2	-5.59	118.28	122.20
3	В	25	U	N1-C2-O2	5.59	126.72	122.80
2	А	135	U	C2-N1-C1'	5.59	124.41	117.70
2	А	784	G	P-O3'-C3'	5.59	126.41	119.70
2	А	897	С	C5-C6-N1	5.59	123.79	121.00
3	В	31	С	N1-C2-O2	5.58	122.25	118.90
2	А	783	А	C2-N3-C4	5.57	113.39	110.60
34	h	733	GLU	C-N-CA	5.56	135.60	121.70
2	А	1494	А	P-O3'-C3'	5.55	126.36	119.70
2	А	2127	G	P-O3'-C3'	5.53	126.33	119.70
2	А	1352	U	N3-C2-O2	-5.51	118.34	122.20
2	А	1267	U	C2-N1-C1'	5.50	124.30	117.70
2	А	243	U	N1-C2-O2	5.49	126.65	122.80
2	А	343	С	C6-N1-C2	-5.49	118.10	120.30
2	А	1314	С	C6-N1-C1'	-5.48	114.22	120.80
2	А	1314	С	N1-C2-O2	5.47	122.18	118.90
2	А	2431	U	N3-C2-O2	-5.47	118.37	122.20
2	А	1102	С	N1-C2-O2	5.46	122.18	118.90
2	А	1115	G	O4'-C1'-N9	5.46	112.57	108.20
2	А	1343	G	C8-N9-C1'	-5.44	119.93	127.00
2	A	2395	С	C6-N1-C2	-5.44	118.13	120.30
2	А	323	С	N1-C2-O2	5.42	122.15	118.90



α \cdot \cdot \cdot	C	•	
Continued	trom	previous	<i>paae</i>
00100010000	J. 01.0	proceed as	<i>p</i> @ <i>g</i> o

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	А	635	С	C6-N1-C2	-5.42	118.13	120.30
2	А	1585	С	N1-C2-O2	5.39	122.13	118.90
2	А	323	С	C2-N1-C1'	5.38	124.72	118.80
2	А	1267	U	N3-C2-O2	-5.38	118.43	122.20
2	А	984	A	C2-N3-C4	5.37	113.28	110.60
2	А	847	U	N3-C2-O2	-5.36	118.45	122.20
2	А	2195	U	C2-N1-C1'	5.33	124.10	117.70
2	А	183	С	N1-C2-O2	5.31	122.08	118.90
2	А	1956	U	C2-N1-C1'	5.31	124.07	117.70
2	А	1320	С	C6-N1-C2	-5.30	118.18	120.30
2	А	1509	A	O4'-C1'-N9	5.29	112.43	108.20
2	А	2720	U	N3-C2-O2	-5.29	118.50	122.20
34	h	800	LYS	C-N-CA	5.29	134.93	121.70
2	А	2421	G	N3-C4-N9	5.27	129.16	126.00
2	А	901	С	N3-C2-O2	-5.27	118.21	121.90
2	А	2656	U	C2-N1-C1'	5.26	124.02	117.70
2	А	919	U	N3-C2-O2	-5.26	118.52	122.20
2	А	1167	С	C2-N1-C1'	5.25	124.58	118.80
2	А	1379	U	N3-C2-O2	-5.25	118.53	122.20
2	А	1669	A	C4-N9-C1'	5.25	135.75	126.30
2	А	867	С	N3-C2-O2	-5.20	118.26	121.90
2	А	135	U	N3-C2-O2	-5.19	118.57	122.20
2	А	2656	U	N3-C2-O2	-5.19	118.57	122.20
2	А	1779	U	C2-N1-C1'	5.18	123.92	117.70
2	А	1313	U	C5-C6-N1	5.18	125.29	122.70
2	А	2739	U	N3-C2-O2	-5.17	118.58	122.20
2	А	2430	A	N3-C4-N9	5.17	131.54	127.40
2	А	353	C	C2-N1-C1'	5.15	124.46	118.80
2	А	2043	C	C6-N1-C2	-5.14	118.25	120.30
2	А	985	C	N1-C2-O2	5.13	121.98	118.90
2	А	2585	U	N3-C2-O2	-5.13	118.61	122.20
2	А	243	U	N3-C2-O2	-5.12	118.61	122.20
2	А	2636	C	C2-N1-C1'	5.10	124.41	118.80
2	А	1076	C	C6-N1-C2	-5.09	118.26	120.30
2	A	1394	U	C6-N1-C1'	5.09	128.33	121.20
2	A	915	C	C2-N1-C1'	5.08	124.39	118.80
3	В	70	C	C6-N1-C2	-5.08	118.27	120.30
2	А	1352	U	N1-C2-O2	5.08	126.36	122.80
2	A	484	C	C5-C6-N1	5.07	123.54	121.00
2	A	1658	C	C6-N1-C2	-5.05	118.28	120.30
2	A	2430	A	N3-C4-C5	-5.05	123.27	126.80
2	A	783	A	C4-N9-C1'	5.05	135.38	126.30



Mol	Chain	\mathbf{Res}	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	А	206	U	N1-C2-O2	5.04	126.33	122.80
2	А	1656	С	C5-C6-N1	5.04	123.52	121.00
2	А	343	С	C2-N1-C1'	5.03	124.33	118.80
2	А	985	С	C6-N1-C2	-5.03	118.29	120.30
2	А	277	G	N3-C4-C5	-5.02	126.09	128.60
2	А	2667	С	N1-C2-O2	5.02	121.91	118.90
2	А	158	U	C2-N1-C1'	5.02	123.72	117.70
2	А	2646	С	C5-C6-N1	5.01	123.51	121.00

There are no chirality outliers.

All (6) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
10	Ι	105	LYS	Peptide
10	Ι	107	GLU	Peptide
32	е	31	HIS	Peptide
34	h	289	GLU	Peptide
34	h	618	ILE	Peptide
34	h	801	GLN	Peptide

5.2 Too-close contacts (i)

Due to software issues we are unable to calculate clashes - this section is therefore empty.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
4	С	270/273~(99%)	262 (97%)	8 (3%)	0	100	100
5	D	207/209~(99%)	199 (96%)	8 (4%)	0	100	100
6	Ε	199/201~(99%)	197 (99%)	2(1%)	0	100	100



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
7	F	176/179~(98%)	169~(96%)	7~(4%)	0	100	100
8	G	174/177~(98%)	167~(96%)	7~(4%)	0	100	100
9	Η	147/149~(99%)	141 (96%)	6~(4%)	0	100	100
10	Ι	123/165~(74%)	114 (93%)	7~(6%)	2(2%)	9	37
11	J	132/142~(93%)	122 (92%)	10 (8%)	0	100	100
12	Κ	140/142~(99%)	136 (97%)	4 (3%)	0	100	100
13	L	121/123 (98%)	119 (98%)	2(2%)	0	100	100
14	М	142/144~(99%)	138 (97%)	4 (3%)	0	100	100
15	Ν	134/136~(98%)	128 (96%)	6 (4%)	0	100	100
16	О	123/127~(97%)	119 (97%)	4 (3%)	0	100	100
17	Р	115/117~(98%)	114 (99%)	1 (1%)	0	100	100
18	Q	112/115~(97%)	110 (98%)	2(2%)	0	100	100
19	R	115/118 (98%)	115 (100%)	0	0	100	100
20	S	101/103~(98%)	100 (99%)	1 (1%)	0	100	100
21	Т	108/110 (98%)	107 (99%)	1 (1%)	0	100	100
22	U	90/100 (90%)	86 (96%)	4 (4%)	0	100	100
23	V	101/104 (97%)	94 (93%)	7 (7%)	0	100	100
24	W	92/94~(98%)	91 (99%)	1 (1%)	0	100	100
25	Х	74/85~(87%)	70~(95%)	4 (5%)	0	100	100
26	Y	75/78~(96%)	75 (100%)	0	0	100	100
27	Ζ	60/63~(95%)	59~(98%)	1 (2%)	0	100	100
28	a	56/59~(95%)	55~(98%)	1 (2%)	0	100	100
29	b	54/57~(95%)	54 (100%)	0	0	100	100
30	с	49/55~(89%)	48 (98%)	1 (2%)	0	100	100
31	d	44/46~(96%)	44 (100%)	0	0	100	100
32	е	62/65~(95%)	55 (89%)	5 (8%)	2(3%)	4	22
33	f	36/38~(95%)	36 (100%)	0	0	100	100
34	h	836/838 (100%)	753 (90%)	75 (9%)	8 (1%)	15	49
35	k	53/57~(93%)	46 (87%)	7 (13%)	0	100	100
All	All	4321/4469~(97%)	4123 (95%)	186 (4%)	12 (0%)	44	73

All (12) Ramachandran outliers are listed below:



Mol	Chain	Res	Type
32	е	32	ILE
34	h	734	THR
32	е	33	LEU
34	h	38	LEU
34	h	258	GLN
34	h	231	ASP
34	h	257	PHE
34	h	801	GLN
34	h	404	ILE
34	h	287	VAL
10	Ι	108	VAL
10	Ι	67	THR

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percer	ntiles
4	С	216/218~(99%)	215 (100%)	1 (0%)	88	94
5	D	164/164~(100%)	164 (100%)	0	100	100
6	Ε	165/165~(100%)	164 (99%)	1 (1%)	86	94
7	F	148/150~(99%)	147~(99%)	1 (1%)	84	93
8	G	137/138~(99%)	136~(99%)	1 (1%)	84	93
9	Н	114/114~(100%)	113 (99%)	1 (1%)	78	91
10	Ι	95/123~(77%)	93~(98%)	2(2%)	53	79
11	J	104/110~(94%)	103 (99%)	1 (1%)	76	90
12	Κ	116/116~(100%)	116 (100%)	0	100	100
13	L	104/104~(100%)	104 (100%)	0	100	100
14	М	103/103~(100%)	103 (100%)	0	100	100
15	Ν	109/109~(100%)	109 (100%)	0	100	100
16	Ο	102/103~(99%)	101 (99%)	1 (1%)	76	90
17	Р	87/87~(100%)	87 (100%)	0	100	100
18	Q	99/100~(99%)	99 (100%)	0	100	100



Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
19	R	89/90~(99%)	89 (100%)	0	100	100
20	S	84/84~(100%)	83~(99%)	1 (1%)	71	88
21	Т	93/93~(100%)	93 (100%)	0	100	100
22	U	79/84~(94%)	79~(100%)	0	100	100
23	V	83/85~(98%)	83 (100%)	0	100	100
24	W	78/78~(100%)	78 (100%)	0	100	100
25	Х	57/63~(90%)	57 (100%)	0	100	100
26	Y	67/68~(98%)	67 (100%)	0	100	100
27	Ζ	54/55~(98%)	54 (100%)	0	100	100
28	a	48/49~(98%)	47 (98%)	1 (2%)	53	79
29	b	47/48~(98%)	47 (100%)	0	100	100
30	с	45/49~(92%)	44 (98%)	1 (2%)	52	78
31	d	38/38~(100%)	38 (100%)	0	100	100
32	е	51/52~(98%)	51 (100%)	0	100	100
33	f	34/34~(100%)	34 (100%)	0	100	100
34	h	714/714 (100%)	703~(98%)	11 (2%)	65	85
35	k	21/30~(70%)	21 (100%)	0	100	100
All	All	3545/3618 (98%)	3522 (99%)	23 (1%)	86	94

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	С	174	LEU
6	Е	170	ARG
7	F	74	VAL
8	G	49	THR
9	Н	124	THR
10	Ι	64	VAL
10	Ι	72	LEU
11	J	80	LEU
16	0	57	THR
20	S	63	VAL
28	a	4	THR
30	с	47	VAL
34	h	37	LYS
34	h	148	GLU



Conti	Continuea from previous page				
Mol	Chain	Res	Type		
34	h	236	TYR		
34	h	272	VAL		
34	h	289	GLU		
34	h	305	MET		
34	h	396	THR		
34	h	401	PHE		
34	h	475	LYS		
34	h	477	ASN		
34	h	730	LEU		

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (37) such sidechains are listed below:

Mol	Chain	Res	Type
4	С	260	ASN
5	D	49	GLN
5	D	173	GLN
6	Е	136	GLN
7	F	23	ASN
7	F	63	GLN
8	G	38	ASN
9	Н	33	GLN
9	Н	135	HIS
10	Ι	103	ASN
11	J	12	GLN
11	J	30	GLN
14	М	54	GLN
15	N	3	GLN
16	0	18	GLN
16	0	62	ASN
18	Q	41	GLN
18	Q	75	GLN
19	R	44	GLN
19	R	72	ASN
19	R	81	ASN
20	S	82	HIS
22	U	48	GLN
22	U	92	ASN
23	V	74	ASN
26	Y	6	GLN
27	Ζ	27	ASN
27	Ζ	39	GLN
34	h	29	ASN



Continued from previous page...

Mol	Chain	Res	Type
34	h	87	GLN
34	h	354	HIS
34	h	415	ASN
34	h	520	GLN
34	h	533	GLN
34	h	629	ASN
34	h	662	GLN
34	h	731	HIS

5.3.3 RNA (i)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	2	2/3~(66%)	1 (50%)	0
2	А	2878/2883~(99%)	446 (15%)	8~(0%)
3	В	119/120~(99%)	14 (11%)	0
All	All	2999/3006~(99%)	461 (15%)	8~(0%)

All (461) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	2	76	А
2	А	10	А
2	А	12	U
2	А	15	G
2	А	34	U
2	А	35	G
2	А	46	G
2	А	58	G
2	А	63	А
2	А	71	А
2	А	74	А
2	А	75	G
2	А	84	А
2	А	91	А
2	А	101	А
2	А	102	U
2	А	118	А
2	А	119	А
2	А	120	U
2	А	125	А
2	А	136	G



Mol	Chain	Res	Type
2	А	137	U
2	А	138	U
2	А	139	U
2	А	141	G
2	А	163	С
2	А	165	А
2	А	181	А
2	А	196	А
2	А	199	А
2	А	215	G
2	А	216	А
2	А	221	А
2	А	222	A
2	A	226	A
2	А	248	G
2	A	266	G
2	А	272	А
2	А	276	U
2	А	277	G
2	А	279	А
2	А	302	C
2	А	311	А
2	А	329	G
2	А	330	А
2	А	353	С
2	А	354	А
2	А	361	G
2	А	362	А
2	А	372	G
2	A	386	G
2	А	396	G
2	A	399	U
2	A	404	A
2	A	405	U
2	A	406	G
2	A	411	G
2	A	412	A
2	A	420	C
2	A	424	G
2	A	435	C
2	A	455	С
2	А	467	G



Mol	Chain	Res	Type
2	А	477	А
2	А	479	А
2	А	481	G
2	А	491	G
2	А	505	А
2	А	509	С
2	А	529	А
2	А	530	G
2	А	531	С
2	А	532	А
2	А	543	G
2	А	549	G
2	А	551	G
2	A	563	A
2	A	573	U
2	A	575	A
2	А	592	А
2	А	603	А
2	А	613	А
2	А	614	А
2	А	615	U
2	А	621	А
2	А	627	А
2	А	637	А
2	А	645	С
2	А	646	U
2	А	647	G
2	А	651	G
2	А	653	U
2	A	654	A
2	A	655	A
2	A	686	U
2	A	714	U
2	A	717	С
2	A	726	G
2	A	730	A
2	A	747	U
2	A	762	U
2	A	764	A
2	A	765	C
2	A	775	G
2	А	776	G



Mol	Chain	Res	Type
2	А	777	G
2	А	782	А
2	А	784	G
2	А	785	G
2	А	790	U
2	А	791	С
2	А	792	А
2	А	800	А
2	А	801	G
2	А	805	G
2	А	811	U
2	А	812	С
2	А	827	U
2	А	828	U
2	А	831	G
2	А	845	A
2	А	846	U
2	А	847	U
2	А	859	G
2	А	869	G
2	А	878	А
2	А	896	А
2	А	897	С
2	А	907	G
2	А	910	А
2	А	934	U
2	А	946	С
2	А	953	G
2	А	961	С
2	А	973	А
2	A	974	G
2	A	983	A
2	А	989	G
2	A	996	A
2	A	1005	С
2	A	1009	A
2	А	1012	U
2	A	1013	С
2	А	1023	U
2	A	1025	G
2	A	1026	G
2	А	1033	U



Mol	Chain	Res	Type
2	А	1040	А
2	А	1046	А
2	А	1060	U
2	А	1069	А
2	А	1070	А
2	А	1071	G
2	А	1073	А
2	А	1074	G
2	А	1083	U
2	А	1084	А
2	А	1087	G
2	А	1088	А
2	A	1097	U
2	А	1101	U
2	A	1112	G
2	A	1122	G
2	А	1128	G
2	А	1132	U
2	А	1133	А
2	А	1135	С
2	А	1142	А
2	А	1155	А
2	А	1173	U
2	А	1178	С
2	А	1212	G
2	А	1218	G
2	А	1227	G
2	А	1236	G
2	A	1238	G
2	А	1247	А
2	A	1250	G
2	A	1253	A
2	A	1256	G
2	А	1271	G
2	A	$127\overline{2}$	A
2	A	1275	A
2	А	1300	G
2	A	1301	A
2	А	1321	А
2	A	$1\overline{329}$	U
2	A	1345	С
2	А	1352	U



Mol	Chain	Res	Type
2	А	1360	G
2	А	1365	А
2	А	1378	А
2	А	1379	U
2	А	1383	А
2	А	1416	G
2	А	1417	С
2	А	1421	G
2	А	1427	А
2	А	1428	С
2	А	1434	А
2	А	1435	G
2	А	1437	С
2	A	1451	С
2	А	1453	А
2	А	1460	U
2	А	1482	G
2	А	1491	G
2	А	1493	С
2	А	1494	А
2	А	1495	А
2	А	1497	U
2	А	1508	А
2	А	1510	G
2	А	1515	А
2	А	1524	G
2	А	1533	C
2	А	1534	U
2	А	1536	С
2	A	1537	G
2	А	1557	С
2	А	1558	С
2	A	1566	A
2	A	1569	A
2	А	1578	U
2	A	1583	A
2	А	1584	U
2	A	1585	С
2	A	1607	С
2	A	1608	A
2	A	1610	A
2	A	1634	A



Mol	Chain	Res	Type
2	А	1647	U
2	А	1648	U
2	А	1649	G
2	А	1669	А
2	А	1674	G
2	А	1698	А
2	А	1715	G
2	А	1729	U
2	А	1730	С
2	А	1732	С
2	А	1738	G
2	А	1756	G
2	А	1757	А
2	A	1758	U
2	А	1764	С
2	A	1773	А
2	А	1781	U
2	А	1782	U
2	А	1784	А
2	А	1800	С
2	А	1801	А
2	А	1802	А
2	А	1808	А
2	А	1811	G
2	А	1816	С
2	А	1829	А
2	А	1870	С
2	А	1871	А
2	А	1872	А
2	А	1873	G
2	A	1906	G
2	А	1929	G
2	A	1936	A
2	A	1938	A
2	А	1939	U
2	A	1955	U
2	A	1967	С
2	A	1970	A
2	A	1971	U
2	A	1972	G
2	А	1982	U
2	А	1991	U



Mol	Chain	Res	Type
2	А	1992	G
2	А	1993	U
2	А	1997	С
2	А	2020	А
2	А	2023	С
2	А	2030	А
2	А	2031	А
2	А	2033	А
2	А	2043	С
2	А	2055	С
2	А	2056	G
2	А	2060	А
2	А	2061	G
2	A	2062	A
2	А	2069	G
2	A	2093	G
2	А	2097	А
2	А	2104	С
2	А	2106	U
2	А	2108	А
2	А	2111	U
2	А	2112	G
2	А	2113	U
2	A	2114	А
2	A	2115	G
2	А	2117	А
2	A	2118	U
2	A	2119	A
2	А	2120	G
2	A	2123	G
2	A	2125	G
2	A	2126	A
2	A	2128	G
2	A	2130	U
2	A	2131	U
2	A	2132	U
2	A	2133	G
2	A	2134	A
2	A	2137	U
2	A	2145	С
2	A	2146	С
2	A	2148	G



Mol	Chain	Res	Type
2	А	2158	А
2	А	2159	G
2	А	2161	С
2	А	2162	G
2	А	2163	A
2	А	2164	С
2	А	2165	С
2	А	2167	U
2	А	2168	G
2	А	2169	А
2	А	2170	А
2	А	2171	А
2	А	2172	U
2	А	2173	А
2	А	2177	С
2	А	2179	С
2	А	2184	А
2	А	2186	G
2	А	2187	U
2	А	2188	U
2	А	2191	А
2	А	2192	U
2	А	2198	А
2	А	2199	А
2	А	2203	U
2	А	2204	G
2	А	2210	U
2	А	2211	А
2	А	2223	G
2	А	2225	А
2	A	2238	G
2	А	2239	G
2	A	2266	A
2	А	2278	A
2	A	2279	G
2	A	2283	С
2	А	2287	А
2	A	2288	A
2	А	2305	U
2	A	2308	G
2	А	2309	A
2	А	2310	С



Mol	Chain	Res	Type
2	А	2322	А
2	А	2325	G
2	А	2333	А
2	А	2334	U
2	А	2347	С
2	А	2350	С
2	А	2361	G
2	А	2383	G
2	А	2385	С
2	А	2391	G
2	А	2402	U
2	А	2406	А
2	А	2422	С
2	A	2424	С
2	А	2425	А
2	А	2428	G
2	А	2429	G
2	А	2430	А
2	А	2431	U
2	А	2432	А
2	А	2434	А
2	А	2435	А
2	А	2441	U
2	А	2447	G
2	А	2448	А
2	А	2470	G
2	А	2476	А
2	А	2478	А
2	А	2491	U
2	А	2494	G
2	A	2498	С
2	A	2502	G
2	A	2505	G
2	A	2518	A
2	A	2520	С
2	A	2529	G
2	А	2547	А
2	A	2562	U
2	А	2566	А
2	A	2567	G
2	A	2572	A
2	А	2585	U



Mol	Chain	Res	Type
2	А	2586	U
2	А	2602	А
2	А	2603	G
2	А	2609	U
2	А	2613	U
2	А	2615	U
2	А	2621	G
2	А	2629	U
2	А	2646	С
2	А	2654	А
2	А	2663	G
2	А	2682	А
2	А	2684	U
2	А	2689	U
2	А	2690	U
2	А	2714	G
2	А	2716	С
2	А	2726	А
2	А	2733	А
2	А	2739	U
2	А	2744	G
2	А	2748	А
2	А	2757	А
2	А	2765	А
2	А	2778	А
2	А	2779	U
2	А	2791	G
2	А	2792	А
2	А	2798	U
2	А	2799	А
2	А	2800	А
2	А	2818	U
2	А	2820	А
2	А	2825	G
2	A	2833	U
2	А	2835	А
2	А	2836	U
2	А	2849	U
2	А	2861	U
2	А	2867	G
2	А	2872	А
2	А	2873	A



Mol	Chain	Res	Type
2	А	2879	А
2	А	2880	С
2	А	2883	А
2	А	2884	U
2	А	2886	А
2	А	2891	U
3	В	9	G
3	В	13	G
3	В	24	G
3	В	35	С
3	В	41	G
3	В	56	G
3	В	66	А
3	В	67	G
3	В	88	С
3	В	89	U
3	В	90	С
3	В	105	G
3	В	109	А
3	В	120	U

All (8) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
2	А	613	А
2	А	784	G
2	А	1494	А
2	А	2112	G
2	А	2127	G
2	А	2158	А
2	А	2602	А
2	А	2756	U

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.



5.6 Ligand geometry (i)

Of 69 ligands modelled in this entry, 69 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
2	А	4
35	k	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	k	76:GLN	С	109:ALA	Ν	27.78
1	А	882:G	O3'	894:U	Р	15.86
1	А	1912:A	O3'	1917:U	Р	14.04
1	А	545:U	O3'	548:G	Р	13.58
1	А	1173:U	O3'	1177:G	Р	13.39



6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-10073. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections (i)

6.1.1 Primary map



The images above show the map projected in three orthogonal directions.

6.2 Central slices (i)

6.2.1 Primary map



X Index: 160



Z Index: 160



The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices (i)

6.3.1 Primary map



X Index: 170

Y Index: 157

Z Index: 165

The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) (i)

6.4.1 Primary map



The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.



6.5 Orthogonal surface views (i)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.82. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.6 Mask visualisation (i)

This section was not generated. No masks/segmentation were deposited.



7 Map analysis (i)

This section contains the results of statistical analysis of the map.

7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



7.2 Volume estimate (i)



The volume at the recommended contour level is 2156 nm^3 ; this corresponds to an approximate mass of 1948 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



7.3 Rotationally averaged power spectrum (i)



*Reported resolution corresponds to spatial frequency of 0.323 \AA^{-1}



8 Fourier-Shell correlation (i)

This section was not generated. No FSC curve or half-maps provided.



9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-10073 and PDB model 6S0K. Per-residue inclusion information can be found in section 3 on page 11.

9.1 Map-model overlay (i)



The images above show the 3D surface view of the map at the recommended contour level 0.82 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.



9.2 Q-score mapped to coordinate model (i)



The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.82).



9.4 Atom inclusion (i)



At the recommended contour level, 89% of all backbone atoms, 90% of all non-hydrogen atoms, are inside the map.



9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (0.82) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.8960	0.4800
2	0.9190	0.5050
А	0.9780	0.5100
В	0.9900	0.4970
С	0.9230	0.5540
D	0.9250	0.5420
Е	0.9060	0.5090
F	0.8530	0.4200
G	0.9070	0.4680
Н	0.5930	0.3700
Ι	0.1720	0.1650
J	0.2120	0.1100
K	0.9370	0.5470
L	0.8840	0.5390
М	0.9340	0.5410
Ν	0.9210	0.5420
0	0.9210	0.5330
Р	0.9400	0.4950
Q	0.9010	0.5330
R	0.9380	0.5330
S	0.9250	0.5250
Т	0.8930	0.5240
U	0.8590	0.5130
V	0.9150	0.4960
W	0.9250	0.5170
X	0.9330	0.5570
Y	0.9070	0.5330
Z	0.8510	0.4730
a	0.9110	0.5310
b	0.9020	0.5380
с	0.2360	0.4130
d	0.9180	0.5580
e	0.9330	0.5520
f	0.9320	0.5440
h	0.3510	0.1690
k	0.4930	0.3020

0.0 <0.0

1.0

