

wwPDB X-ray Structure Validation Summary Report (i)

Mar 24, 2022 – 03:28 pm GMT

PDB ID : 6S0R

Title : The crystal structure of kanamycin B dioxygenase (KanJ) from Streptomyces

kanamyceticus complex with nickel, sulfate and chloride

Authors: Mrugala, B.; Porebski, P.J.; Niedzialkowska, E.; Cymborowski, M.T.; Minor,

W.; Borowski, T.

Deposited on : 2019-06-18

Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul: 1.8.4, CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.27

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0267

CCP4 : 7.1.010 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.27

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY\ DIFFRACTION$

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \text{Whole archive} \\ (\#\text{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar \ resolution} \\ (\#{\rm Entries, \ resolution \ range(\AA)}) \end{array}$
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	288	94%	
1	В	288	93%	5% •
1	С	288	94%	
1	D	288	91%	6% •
1	Е	288	91%	5% 5%



Continued from previous page...

Mol	Chain	Length	Quality of chain	
1	F	288	94%	



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 13998 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Kanamycin B dioxygenase.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	A	283	Total	С	N	О	S	0	0	0
1	Λ	200	2202	1401	387	404	10	0	0	0
1	В	284	Total	С	N	О	S	0	3	0
1	Ъ	204	2230	1418	391	411	10	0	3	U
1	С	279	Total	С	N	О	S	0	3	0
1		219	2185	1388	383	404	10	0		0
1	D	280	Total	С	N	О	S	0	2	0
1	D	280	2193	1392	386	405	10	0		0
1	Е	275	Total	С	N	О	S	0	1	0
1	l L	210	2142	1364	377	392	9	0	1	0
1	F	278	Total	С	N	О	S	0	2	0
1	I'	210	2172	1380	381	401	10	U	<u> </u>	U

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	expression tag	UNP Q6L732
A	-1	ASN	-	expression tag	UNP Q6L732
A	0	ALA	-	expression tag	UNP Q6L732
В	-2	SER	-	expression tag	UNP Q6L732
В	-1	ASN	-	expression tag	UNP Q6L732
В	0	ALA	-	expression tag	UNP Q6L732
С	-2	SER	-	expression tag	UNP Q6L732
С	-1	ASN	-	expression tag	UNP Q6L732
С	0	ALA	-	expression tag	UNP Q6L732
D	-2	SER	-	expression tag	UNP Q6L732
D	-1	ASN	-	expression tag	UNP Q6L732
D	0	ALA	-	expression tag	UNP Q6L732
Е	-2	SER	-	expression tag	UNP Q6L732
Е	-1	ASN	-	expression tag	UNP Q6L732
Е	0	ALA	=	expression tag	UNP Q6L732
F	-2	SER	=	expression tag	UNP Q6L732
F	-1	ASN	=	expression tag	UNP Q6L732




Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	0	ALA	-	expression tag	UNP Q6L732

• Molecule 2 is NICKEL (II) ION (three-letter code: NI) (formula: Ni) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Ni 1 1	0	0
2	В	1	Total Ni 1 1	0	0
2	C	1	Total Ni 1 1	0	0
2	D	1	Total Ni 1 1	0	0
2	Е	1	Total Ni 1 1	0	0
2	F	1	Total Ni 1 1	0	0

• Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0

Continued from previous page...

Mol	Chain	$egin{array}{c} n \ previous \ pa \ \hline \mathbf{Residues} \end{array}$	Aton	ns		ZeroOcc	AltConf
3	A	1		O 4	S 1	0	0
3	A	1	5	O 4	S 1	0	0
3	В	1	5	O 4	S 1	0	0
3	В	1	5	O 4	S 1	0	0
3	С	1	5	O 4	S 1	0	0
3	С	1	5	O 4	S 1	0	0
3	С	1	5	O 4	S 1	0	0
3	D	1	5	O 4	S 1	0	0
3	D	1	5	O 4	S 1	0	0
3	D	1	5	O 4	S 1	0	0
3	D	1	10	O 8	S 2	0	1
3	E	1	5	O 4	S 1	0	0
3	E	1	5	O 4	S 1	0	0
3	Е	1	5	O 4	S 1	0	0
3	F	1	5	O 4	S 1	0	0
3	F	1		O 4	S 1	0	0
3	F	1		O 4	S 1	0	0
3	F	1		O 4	S 1	0	0
3	F	1		O 4	S 1	0	0

• Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	В	1	Total Cl 1 1	0	0

• Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	122	Total O 122 122	0	0
5	В	226	Total O 226 226	0	0
5	С	100	Total O 100 100	0	0
5	D	140	Total O 140 140	0	0
5	E	63	Total O 63 63	0	0
5	F	106	Total O 106 106	0	0



3 Residue-property plots (i)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Kanamycin B dioxygenase





 \bullet Molecule 1: Kanamycin B dioxygenase





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	50.37Å 184.21Å 110.15Å	Depositor
a, b, c, α , β , γ	90.00° 94.23° 90.00°	Depositor
Resolution (Å)	47.22 - 2.50	Depositor
Resolution (A)	47.17 - 2.50	EDS
% Data completeness	87.5 (47.22-2.50)	Depositor
(in resolution range)	87.6 (47.17-2.50)	EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.64 (at 2.51Å)	Xtriage
Refinement program	REFMAC 5.8.0238	Depositor
P. P.	0.170 , 0.208	Depositor
R, R_{free}	0.177 , 0.215	DCC
R_{free} test set	3043 reflections $(5.04%)$	wwPDB-VP
Wilson B-factor (Å ²)	37.7	Xtriage
Anisotropy	0.078	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	(Not available), (Not available)	EDS
L-test for twinning ²	$ < L > = 0.48, < L^2> = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	13998	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.89% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, NI, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.65	0/2271	0.79	0/3124	
1	В	0.70	0/2308	0.82	0/3174	
1	С	0.65	0/2257	0.79	0/3104	
1	D	0.65	0/2265	0.81	0/3114	
1	Е	0.65	0/2211	0.79	0/3042	
1	F	0.65	0/2244	0.81	0/3086	
All	All	0.66	0/13556	0.80	0/18644	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2202	0	2149	5	0
1	В	2230	0	2181	12	0
1	С	2185	0	2140	10	0
1	D	2193	0	2151	6	0
1	Е	2142	0	2098	6	0
1	F	2172	0	2121	5	0
2	A	1	0	0	0	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	В	1	0	0	0	0
2	С	1	0	0	0	0
2	D	1	0	0	0	0
2	Е	1	0	0	0	0
2	F	1	0	0	0	0
3	A	20	0	0	0	0
3	В	10	0	0	0	0
3	С	15	0	0	0	0
3	D	25	0	0	0	0
3	Ε	15	0	0	0	0
3	F	25	0	0	0	0
4	В	1	0	0	0	0
5	A	122	0	0	1	0
5	В	226	0	0	0	0
5	С	100	0	0	0	0
5	D	140	0	0	2	0
5	Е	63	0	0	2	0
5	F	106	0	0	0	0
All	All	13998	0	12840	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 36 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$egin{aligned} ext{Interatomic} \ ext{distance} \ (ext{Å}) \end{aligned}$	$egin{aligned} ext{Clash} \ ext{overlap } (ext{Å}) \end{aligned}$
1:F:269:VAL:HG21	1:F:273:VAL:HG22	1.76	0.66
1:D:111:ASP:HB2	5:D:431:HOH:O	1.95	0.66
1:C:192:GLU:HB3	1:F:65:LEU:HD13	1.81	0.62
1:B:48[A]:ARG:HD3	1:C:48:ARG:HA	1.81	0.62
1:D:14:LEU:HD21	1:D:21:LEU:CD2	2.32	0.59

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



The Analysed column shows	the number	of residues	for which	the ba	ackbone	conformation	was
analysed, and the total number	r of residue	es.					

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	$281/288 \ (98\%)$	277 (99%)	4 (1%)	0	100	100
1	В	285/288~(99%)	281 (99%)	4 (1%)	0	100	100
1	C	280/288~(97%)	276 (99%)	4 (1%)	0	100	100
1	D	280/288~(97%)	277 (99%)	3 (1%)	0	100	100
1	E	$274/288\ (95\%)$	270 (98%)	4 (2%)	0	100	100
1	F	$278/288 \ (96\%)$	273 (98%)	5 (2%)	0	100	100
All	All	1678/1728 (97%)	1654 (99%)	24 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	243/248 (98%)	241 (99%)	2 (1%)	81 93
1	В	248/248 (100%)	245 (99%)	3 (1%)	71 88
1	С	243/248 (98%)	241 (99%)	2 (1%)	81 93
1	D	244/248 (98%)	237 (97%)	7 (3%)	42 69
1	Е	237/248 (96%)	232 (98%)	5 (2%)	53 78
1	F	240/248 (97%)	238 (99%)	2 (1%)	81 93
All	All	1455/1488 (98%)	1434 (99%)	21 (1%)	65 86

5 of 21 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	Ε	15	THR
1	Ε	133	LEU
1	F	272	ASP
1	Е	240	GLU
1	Е	86	ARG



Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	68	GLN
1	A	248	GLN
1	В	67	GLN
1	С	67	GLN
1	F	52	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 29 ligands modelled in this entry, 7 are monoatomic - leaving 22 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trme	Chain	Dog	Res Link		ond leng	gths	В	ond ang	gles
MIOI	Type	Chain	nes	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	F	305	-	4,4,4	0.34	0	6,6,6	0.05	0
3	SO4	A	304	-	4,4,4	0.35	0	6,6,6	0.09	0
3	SO4	D	303	-	4,4,4	0.37	0	6,6,6	0.08	0
3	SO4	D	305[A]	-	4,4,4	0.40	0	6,6,6	0.09	0
3	SO4	С	303	-	4,4,4	0.36	0	6,6,6	0.09	0
3	SO4	F	306	-	4,4,4	0.38	0	6,6,6	0.04	0
3	SO4	A	303	-	4,4,4	0.35	0	6,6,6	0.12	0



Mol	Trino	Chain	Res	Link	В	Bond lengths		В	ond ang	gles
MIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	Е	304	-	4,4,4	0.37	0	6,6,6	0.09	0
3	SO4	С	302	-	4,4,4	0.26	0	6,6,6	0.15	0
3	SO4	F	304	-	4,4,4	0.38	0	6,6,6	0.09	0
3	SO4	В	303	-	4,4,4	0.95	0	6,6,6	0.26	0
3	SO4	С	304	-	4,4,4	0.37	0	6,6,6	0.04	0
3	SO4	A	305	-	4,4,4	0.36	0	6,6,6	0.06	0
3	SO4	В	302	-	4,4,4	0.32	0	6,6,6	0.12	0
3	SO4	D	302	-	4,4,4	0.26	0	6,6,6	0.12	0
3	SO4	Е	302	-	4,4,4	0.30	0	6,6,6	0.08	0
3	SO4	D	304	-	4,4,4	0.34	0	6,6,6	0.10	0
3	SO4	Е	303	-	4,4,4	0.37	0	6,6,6	0.08	0
3	SO4	F	303	-	4,4,4	0.36	0	6,6,6	0.12	0
3	SO4	D	305[B]	-	4,4,4	0.38	0	6,6,6	0.10	0
3	SO4	F	302	-	4,4,4	0.28	0	6,6,6	0.09	0
3	SO4	A	302	-	4,4,4	0.37	0	6,6,6	0.06	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.













































































































5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\#\mathrm{RSRZ}{>}2$		$\mathbf{OWAB}(\mathbf{\mathring{A}}^2)$	Q < 0.9	
1	A	283/288 (98%)	0.04	5 (1%)	68	71	23, 39, 78, 125	0
1	В	284/288 (98%)	-0.11	3 (1%)	80	82	14, 30, 61, 111	0
1	С	279/288 (96%)	0.14	6 (2%)	62	65	24, 51, 102, 128	0
1	D	280/288 (97%)	-0.00	4 (1%)	75	77	22, 38, 73, 100	0
1	E	275/288 (95%)	0.52	19 (6%)	16	17	37, 59, 95, 123	0
1	F	278/288 (96%)	0.01	4 (1%)	75	77	24, 45, 82, 109	0
All	All	1679/1728 (97%)	0.10	41 (2%)	59	62	14, 44, 88, 128	0

The worst 5 of 41 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	Е	274	ASP	4.7
1	Е	273	VAL	4.5
1	Е	272	ASP	4.2
1	A	279	ASP	3.9
1	Е	271	GLY	3.5

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.




6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
3	SO4	A	304	5/5	0.81	0.34	84,92,103,107	0
3	SO4	D	303	5/5	0.81	0.19	95,103,105,109	0
3	SO4	Е	304	5/5	0.82	0.21	103,107,110,114	0
3	SO4	F	304	5/5	0.84	0.39	107,117,126,129	0
3	SO4	С	304	5/5	0.85	0.28	126,133,135,136	0
3	SO4	F	305	5/5	0.87	0.14	93,94,99,103	0
3	SO4	F	306	5/5	0.88	0.21	110,114,119,120	0
3	SO4	A	303	5/5	0.89	0.18	73,77,82,90	0
3	SO4	В	303	5/5	0.89	0.18	61,68,74,77	0
3	SO4	F	303	5/5	0.91	0.12	87,92,98,104	0
3	SO4	Е	303	5/5	0.92	0.26	91,91,97,103	0
3	SO4	A	305	5/5	0.92	0.40	111,117,121,122	0
3	SO4	С	303	5/5	0.93	0.15	79,89,93,94	0
3	SO4	D	305[A]	5/5	0.93	0.15	38,40,41,42	5
3	SO4	D	305[B]	5/5	0.93	0.15	40,43,45,45	5
3	SO4	D	304	5/5	0.95	0.14	76,82,84,85	0
3	SO4	Е	302	5/5	0.96	0.11	69,70,80,82	0
2	NI	A	301	1/1	0.98	0.03	27,27,27,27	0
2	NI	D	301	1/1	0.98	0.04	31,31,31,31	0
3	SO4	С	302	5/5	0.98	0.12	45,52,57,58	0
3	SO4	F	302	5/5	0.98	0.18	55,56,65,65	0
2	NI	В	301	1/1	0.99	0.04	22,22,22,22	0
2	NI	Е	301	1/1	0.99	0.08	42,42,42,42	0
3	SO4	D	302	5/5	0.99	0.14	41,49,51,52	0
3	SO4	A	302	5/5	0.99	0.16	53,54,56,57	0
2	NI	С	301	1/1	0.99	0.05	27,27,27,27	0
4	CL	В	304	1/1	0.99	0.09	41,41,41,41	0
2	NI	F	301	1/1	1.00	0.07	31,31,31,31	0
3	SO4	В	302	5/5	1.00	0.12	37,40,41,42	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.















Electron density around SO4 E 304:
























Electron density around SO4 F 306:






























Electron density around SO4 A 305: $2\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)




Electron density around SO4 C 303:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)








Electron density around SO4 D 305 (A): $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${ m mF}_o{ m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)





Electron density around SO4 D 305 (B): 2mFo-DFc (at 0.7 rmsd) in gray mFo-DFc (at 3 rmsd) in purple (negative) and green (positive)




Electron density around SO4 D 304:





























Electron density around SO4 F 302:

















Electron density around SO4 D 302:




















Electron density around CL B 304: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_{o}\text{-}\mathrm{DF}_{c}$ (at 3 rmsd) in purple (negative) and green (positive)









6.5 Other polymers (i)

There are no such residues in this entry.

