Full wwPDB X-ray Structure Validation Report (i)

Jul 13, 2023 – 10:13 AM EDT

PDB ID	:	8SBC
Title	:	Co-structure of Phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic subunit alpha isoform and brain penetrant inhibitors
Authors	:	Knapp, M.S.; Elling, R.A.; Tang, J.
Deposited on	:	2023-04-03
Resolution	:	2.30 Å(reported)

The following versions of software and data (see references (i)) were used in the production of this report:

|		
MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.34
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.34

1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.

Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1074	5% / 78% / 12% / 10%
2	B	293	5% / 78% / 12% / 10%

2 Entry composition ⓘ

There are 5 unique types of molecules in this entry. The entry contains 9775 atoms, of which 26 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic subunit alpha isoform.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	970	Total 7580	C 4853	N 1285	O 1377	S 65	0	3	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	232	LYS	MET	engineered mutation	UNP P42336
A	233	LYS	LEU	engineered mutation	UNP P42336
A	780	LYS	TRP	engineered mutation	UNP P42336
A	800	MET	ILE	engineered mutation	UNP P42336
A	850	TRP	VAL	engineered mutation	UNP P42336
A	930	VAL	PHE	conflict	UNP P42336
A	1069	HIS	-	expression tag	UNP P42336
A	1070	HIS	-	expression tag	UNP P42336
A	1071	HIS	-	expression tag	UNP P42336
A	1072	HIS	-	expression tag	UNP P42336
A	1073	HIS	-	expression tag	UNP P42336
A	1074	HIS	-	expression tag	UNP P42336

- Molecule 2 is a protein called Phosphatidylinositol 3-kinase regulatory subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	264	Total 2101	C 1320	N 369	O 407	S 5	0	0	0

There is a discrepancy between the modelled and reference sequences:

| Chain	Residue	Modelled	Actual	Comment	Reference
B	306	TYR	THR	engineered mutation	UNP P27986

- Molecule 3 is (2M)-7-[(3R)-3-methylmorpholin-4-yl]-5-[(3S)-3-methylmorpholin-4-yl]-2-(

pyridin-2-yl)-1H-imidazo[4,5-b]pyridine (three-letter code: ZTV) (formula: $C_{21}H_{26}N_6O_2$) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	N	O		
3	A	1	55	21	26	6	2	0	0

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O_4S).

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
4	A	1	5	4	1	0	0

• Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	27	Total O 27 27	0	0
5	B	7	Total O 7 7	0	0

3 Residue-property plots ⓘ

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic subunit alpha isoform

• Molecule 2: Phosphatidylinositol 3-kinase regulatory subunit alpha

4 Data and refinement statistics ⓘ

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α, β, γ	100.08Å 107.85Å 133.72Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	83.95 – 2.30 83.95 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.4 (83.95-2.30) 99.4 (83.95-2.30)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$<I/\sigma(I)>$ [1]	1.07 (at 2.29Å)	Xtriage
Refinement program	PHENIX (1.20.1_4487: ???)	Depositor
R, R_{free}	0.241 , 0.271 0.235 , 0.267	Depositor DCC
R_{free} test set	3181 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å^2)	60.8	Xtriage
Anisotropy	0.536	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.29 , 52.9	EDS
L-test for twinning[2]	$<|L|> = 0.49, <L^2> = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.95	EDS
Total number of atoms	9775	wwPDB-VP
Average B, all atoms (Å^2)	89.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.64% of the height of the origin peak. No significant pseudotranslation is detected.*

---

[1]Intensities estimated from amplitudes.
[2]Theoretical values of $<|L|>$, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: ZTV, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#$|Z|$>5	RMSZ	#$|Z|$>5
1	A	0.24	0/7757	0.43	0/10546
2	B	0.23	0/2137	0.45	0/2885
All	All	0.24	0/9894	0.44	0/13431

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7580	0	7206	78	0
2	B	2101	0	1940	24	0
3	A	29	26	0	0	0
4	A	5	0	0	0	0
5	A	27	0	0	0	0
5	B	7	0	0	0	0
All	All	9749	26	9146	99	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (99) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:158:SER:CB	1:A:159:PRO:HD3	2.22	0.69
1:A:619:LEU:HD22	1:A:623:LEU:HD22	1.76	0.68
1:A:158:SER:CB	1:A:159:PRO:CD	2.73	0.67
1:A:346:VAL:HG21	1:A:351:ILE:HG13	1.75	0.66
1:A:357[B]:ARG:HG2	1:A:372:ASN:OD1	2.00	0.62
1:A:216:VAL:HG12	1:A:218:GLU:H	1.64	0.62
2:B:512:PHE:HB2	2:B:521:ILE:HD11	1.82	0.62
2:B:372:LEU:HD21	2:B:422:VAL:CG1	2.30	0.61
1:A:216:VAL:HG13	1:A:217:PRO:HD2	1.84	0.60
1:A:124:PRO:HG2	1:A:127:GLU:HG3	1.84	0.58
1:A:193:VAL:HG22	1:A:208:THR:HG22	1.85	0.58
1:A:744:PHE:CE1	1:A:748:LEU:HD22	2.39	0.57
2:B:557:ARG:O	2:B:561:LYS:HG3	2.05	0.57
1:A:326:SER:HB3	1:A:329:VAL:HG23	1.86	0.56
1:A:937:PHE:HD1	1:A:938:LEU:HG	1.70	0.56
1:A:856:THR:HG22	1:A:922:MET:HG2	1.88	0.56
2:B:332:GLU:HB2	2:B:429:SER:HB2	1.88	0.56
1:A:19:ARG:HG2	1:A:37:LEU:HD23	1.88	0.56
1:A:328:TRP:HA	1:A:394:PRO:HB3	1.90	0.54
1:A:355:TYR:OH	1:A:419:HIS:ND1	2.30	0.54
1:A:79:ARG:HD2	2:B:489:GLU:OE2	2.07	0.53
1:A:355:TYR:HE2	1:A:357[B]:ARG:HG3	1.74	0.53
1:A:355:TYR:CE2	1:A:357[B]:ARG:HG3	2.44	0.52
2:B:509:ILE:HG23	2:B:521:ILE:HD12	1.92	0.52
1:A:136:VAL:HG13	1:A:686:LEU:HD11	1.93	0.51
1:A:9:GLU:OE2	1:A:38:ARG:NH2	2.36	0.51
2:B:456:PHE:CE1	2:B:574:ARG:HB2	2.45	0.51
1:A:793:LEU:HG	1:A:794:PHE:HD1	1.75	0.51
2:B:354:THR:HA	2:B:426:TYR:O	2.11	0.51
1:A:772:MET:HB2	1:A:778:PRO:HG2	1.92	0.50
2:B:422:VAL:HG22	2:B:423:LYS:N	2.27	0.50
2:B:567:LYS:O	2:B:571:ILE:HG12	2.12	0.50
1:A:39:GLU:HB2	1:A:88:ARG:HH11	1.77	0.50
1:A:489:MET:HA	1:A:492:ILE:HG12	1.94	0.50
2:B:581:LEU:O	2:B:585:THR:HG23	2.13	0.49
2:B:499:GLN:O	2:B:503:ARG:HG3	2.11	0.49
1:A:131:VAL:HG11	1:A:136:VAL:HG21	1.95	0.49
1:A:538:ASP:HB2	1:A:999:ILE:HD12	1.95	0.49
1:A:25:LEU:HD13	2:B:497:GLN:HG3	1.95	0.49
1:A:917:HIS:CE1	1:A:919:SER:HB2	2.48	0.48
1:A:153:LEU:HA	1:A:156:LEU:HD23	1.95	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:829:LEU:HB3	1:A:831:LEU:HD13	1.96	0.47
1:A:351:ILE:C	1:A:412:ARG:HH22	2.18	0.47
1:A:151:VAL:HG21	1:A:302:PHE:HB2	1.97	0.47
1:A:939:ASP:HB3	1:A:1021:TYR:CD1	2.49	0.47
1:A:145:ASN:OD1	1:A:651:ARG:NH2	2.47	0.47
1:A:427:ILE:HD11	1:A:443:LEU:HD22	1.97	0.46
1:A:360:ILE:HG22	1:A:367:LEU:HD12	1.97	0.46
1:A:42:LEU:HD21	1:A:92:LEU:HD11	1.98	0.46
2:B:571:ILE:HG22	2:B:575:LYS:HE2	1.97	0.46
1:A:273:ILE:O	1:A:277:ILE:HG13	2.15	0.46
1:A:488:ASP:O	1:A:492:ILE:HG23	2.16	0.46
1:A:373:THR:HG22	1:A:387:LEU:HD11	1.97	0.46
1:A:937:PHE:CD1	1:A:938:LEU:HG	2.51	0.45
2:B:454:THR:O	2:B:458:GLU:HG2	2.16	0.45
2:B:372:LEU:HD21	2:B:422:VAL:HG11	1.97	0.45
1:A:348:ILE:HG22	1:A:378:CYS:SG	2.57	0.45
1:A:420[B]:CYS:SG	2:B:568:PRO:HB3	2.57	0.45
1:A:430:PHE:O	1:A:644:TYR:OH	2.26	0.44
1:A:861:GLN:HA	1:A:876:THR:HG21	1.99	0.44
1:A:145:ASN:HA	1:A:148:LYS:HZ2	1.83	0.44
1:A:981:GLN:OE1	1:A:1051:TRP:HA	2.16	0.44
1:A:433:THR:O	1:A:485:LYS:NZ	2.49	0.44
1:A:937:PHE:O	1:A:938:LEU:HB2	2.18	0.44
1:A:624:THR:HG23	1:A:627:LYS:H	1.83	0.44
1:A:854:SER:HB2	1:A:922:MET:HE3	2.00	0.44
1:A:855:HIS:HB2	1:A:860:ILE:HD11	1.99	0.43
2:B:405:ILE:O	2:B:409:ARG:HG3	2.18	0.43
1:A:587:VAL:HA	1:A:590:TRP:HB2	2.01	0.43
1:A:860:ILE:HD12	1:A:880:TRP:CE2	2.54	0.43
1:A:734:PHE:O	1:A:738:GLN:HG2	2.18	0.43
1:A:921:ILE:HD13	1:A:954:PHE:HE1	1.84	0.42
1:A:357[B]:ARG:O	1:A:404:LEU:HA	2.19	0.42
1:A:157:ASN:CB	1:A:161:SER:HB3	2.50	0.42
1:A:830:ASP:O	1:A:899:ARG:HD3	2.20	0.42
1:A:357[A]:ARG:O	1:A:404:LEU:HA	2.20	0.42
1:A:821:GLU:O	1:A:825:GLN:HG2	2.19	0.42
2:B:453:ASN:O	2:B:457:GLN:HG2	2.19	0.42
2:B:567:LYS:N	2:B:568:PRO:HD2	2.34	0.42
2:B:422:VAL:HG22	2:B:423:LYS:H	1.85	0.41
1:A:714:ASN:O	1:A:718:ILE:HG13	2.19	0.41
1:A:979:ARG:O	1:A:983:MET:HG3	2.20	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:807:LEU:HD12	1:A:846:GLY:HA3	2.02	0.41
1:A:873:ASN:HB3	1:A:876:THR:OG1	2.21	0.41
1:A:924:LYS:HD3	1:A:928:GLN:HB3	2.02	0.41
1:A:532:LYS:O	1:A:536:THR:HG23	2.20	0.41
1:A:173:SER:OG	1:A:624:THR:HG21	2.20	0.41
1:A:334:LEU:HD22	1:A:396:LEU:HD11	2.02	0.41
1:A:997:LEU:O	1:A:1001:LEU:HG	2.21	0.41
2:B:341:GLU:OE1	2:B:341:GLU:N	2.49	0.41
2:B:494:PHE:HB3	2:B:535:ILE:HG12	2.01	0.41
2:B:520:GLU:OE2	2:B:523:ARG:NH1	2.54	0.41
1:A:392:TYR:CD2	1:A:394:PRO:HD2	2.55	0.40
1:A:565:LEU:HB3	1:A:566:PRO:HD3	2.03	0.40
1:A:267:LEU:HG	1:A:273:ILE:HG13	2.01	0.40
1:A:68:TYR:HB3	1:A:101:VAL:HG13	2.03	0.40
1:A:723:LYS:HE3	1:A:734:PHE:CD1	2.57	0.40
1:A:739:MET:HA	1:A:744:PHE:CD1	2.57	0.40
1:A:809:GLN:HG2	1:A:1010:MET:HE2	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	961/1074 (90%)	928 (97%)	31 (3%)	2 (0%)	47	58
2	B	260/293 (89%)	254 (98%)	6 (2%)	0	100	100
All	All	1221/1367 (89%)	1182 (97%)	37 (3%)	2 (0%)	47	58

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	158	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	762	GLY

5.3.2   Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	792/980 (81%)	791 (100%)	1 (0%)	93	97
2	B	207/272 (76%)	207 (100%)	0	100	100
All	All	999/1252 (80%)	998 (100%)	1 (0%)	93	97

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	971	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3   RNA ⓘ

There are no RNA molecules in this entry.

5.4   Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5   Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6   Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	#$|Z| > 2$	Counts	RMSZ	#$|Z| > 2$
3	ZTV	A	1101	-	31,33,33	1.07	3 (9%)	33,47,47	1.32	5 (15%)
4	SO4	A	1102	-	4,4,4	0.14	0	6,6,6	0.05	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ZTV	A	1101	-	-	7/12/34/34	0/5/5/5

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1101	ZTV	C4-C3	-3.50	1.36	1.42
3	A	1101	ZTV	C1-N	2.42	1.36	1.32
3	A	1101	ZTV	C2-N1	2.32	1.39	1.34

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(º)	Ideal(º)
3	A	1101	ZTV	C4-C3-C2	3.61	121.31	117.79
3	A	1101	ZTV	C2-C3-N2	-2.99	106.43	109.47
3	A	1101	ZTV	C13-O1-C12	-2.46	106.56	109.97
3	A	1101	ZTV	C1-N-C2	-2.38	115.85	117.65
3	A	1101	ZTV	C14-N4-C11	2.27	118.75	114.66

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1101	ZTV	C20-C16-C5-N1
3	A	1101	ZTV	N5-C16-C5-N1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	A	1101	ZTV	N-C1-N4-C14
3	A	1101	ZTV	C-C1-N4-C11
3	A	1101	ZTV	N-C1-N4-C11
3	A	1101	ZTV	C-C4-N3-C6
3	A	1101	ZTV	C20-C16-C5-N2

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand ZTV A 1101

Bond lengths

Bond angles

Torsions

Rings

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	970/1074 (90%)	0.45	51 (5%)	26	33	45, 86, 141, 178	0
2	B	264/293 (90%)	0.49	14 (5%)	26	33	51, 86, 138, 164	0
All	All	1234/1367 (90%)	0.46	65 (5%)	26	33	45, 86, 140, 178	0

All (65) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	968	ALA	5.2
1	A	551	LEU	4.6
2	B	431	TYR	4.5
2	B	521	ILE	4.2
2	B	512	PHE	4.1
1	A	213	HIS	3.8
1	A	261	PHE	3.8
1	A	483	VAL	3.7
1	A	252	LEU	3.7
1	A	255	CYS	3.6
1	A	886	LYS	3.4
1	A	835	PRO	3.4
1	A	477	PHE	3.4
1	A	945	PHE	3.4
1	A	411	GLY	3.3
2	B	440	ASP	3.2
1	A	834	LEU	3.2
1	A	192	VAL	3.2
1	A	881	LEU	3.1
1	A	430	PHE	3.1
1	A	54	ARG	3.0
1	A	525	GLU	3.0
1	A	287	LEU	3.0
1	A	484	VAL	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	436	VAL	2.9
1	A	183	ASN	2.9
1	A	190	ILE	2.9
1	A	480	PHE	2.8
2	B	591	GLN	2.8
1	A	51	LYS	2.8
1	A	194	ILE	2.8
1	A	873	ASN	2.7
2	B	583	TRP	2.7
2	B	328	LEU	2.7
2	B	566	ILE	2.7
1	A	310	ARG	2.7
1	A	188	GLY	2.7
1	A	951	ARG	2.6
2	B	348	ARG	2.6
1	A	259	GLU	2.6
2	B	418	PRO	2.5
1	A	265	TYR	2.5
1	A	262	LEU	2.5
1	A	264	LYS	2.5
1	A	829	LEU	2.4
1	A	286	MET	2.4
1	A	889	ILE	2.4
1	A	182	TYR	2.4
1	A	562	PRO	2.3
1	A	552	TRP	2.3
1	A	273	ILE	2.2
1	A	351	ILE	2.2
1	A	526	ASN	2.2
1	A	764	LEU	2.2
1	A	793	LEU	2.2
1	A	291	GLU	2.2
1	A	216	VAL	2.2
1	A	222	ALA	2.2
1	A	527	ASP	2.1
1	A	161	SER	2.1
2	B	372	LEU	2.1
1	A	171	VAL	2.1
1	A	492	ILE	2.1
2	B	441	ASN	2.1
2	B	404	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no monosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
4	SO4	A	1102	5/5	0.91	0.55	80,94,100,166	0
3	ZTV	A	1101	29/29	0.92	0.20	83,102,124,140	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around ZTV A 1101:

$2mF_o\text{-}DF_c$ (at 0.7 rmsd) in gray
$mF_o\text{-}DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)

6.5 Other polymers ⓘ

There are no such residues in this entry.