



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 11, 2024 – 02:15 PM EST

PDB ID : 1SC7
Title : Human DNA Topoisomerase I (70 Kda) In Complex With The Indenoisoquinoline MJ-II-38 and Covalent Complex With A 22 Base Pair DNA Duplex
Authors : Staker, B.L.; Feese, M.D.; Cushman, M.; Pommier, Y.; Zembower, D.; Stewart, L.; Burgin, A.B.
Deposited on : 2004-02-11
Resolution : 3.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	180529	2866 (3.00-3.00)
Ramachandran outliers	177936	2778 (3.00-3.00)
Sidechain outliers	177891	2781 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	B	10	
2	C	12	
3	D	22	
4	A	592	

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 5629 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called 5'-D(*AP*AP*AP*AP*AP*GP*AP*CP*TP*T)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
1	B	10	203	99	42	53	9	0	0	0

- Molecule 2 is a DNA chain called 5'-D(*(TGP)P*GP*AP*AP*AP*AP*AP*TP*TP*TP*TP*P*T)-3'.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	P	S			
2	C	12	246	120	45	69	11	1	0	0	0

- Molecule 3 is a DNA chain called 5'-D(*AP*AP*AP*AP*AP*TP*TP*TP*TP*TP*CP*CP*AP*AP*GP*TP*CP*TP*TP*TP*TP*T)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
3	D	22	443	217	71	134	21	0	0	0


- Molecule 4 is a protein called DNA topoisomerase I.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	P	S			
4	A	567	4699	2990	822	860	1	26	0	0	0

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	723	PTR	TYR	modified residue	UNP P11387

- Molecule 5 is 4-(5,11-DIOXO-5H-INDENO[1,2-C]ISOQUINOLIN-6(11H)-YL)BUTANOATE (three-letter code: M38) (formula: C₂₀H₁₅NO₄).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
5	C	1	25	20	1	4	0	0

- Molecule 6 is TETRAETHYLENE GLYCOL (three-letter code: PG4) (formula: $C_8H_{18}O_5$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
6	A	1	13	8	5	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: 5'-D(*AP*AP*AP*AP*AP*GP*AP*CP*TP*T)-3'




- Molecule 2: 5'-D(*(TGP)P*GP*AP*AP*AP*AP*AP*TP*TP*TP*TP*T)-3'



- Molecule 3: 5'-D(*AP*AP*AP*AP*AP*TP*TP*TP*TP*TP*CP*CP*AP*AP*GP*TP*CP*TP*TP*TP*TP*T)-3'



- Molecule 4: DNA topoisomerase I



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	260.94Å 74.66Å 57.49Å 90.00° 96.94° 90.00°	Depositor
Resolution (Å)	19.85 – 3.00	Depositor
% Data completeness (in resolution range)	87.9 (19.85-3.00)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
Refinement program	CNX 2002	Depositor
R, R_{free}	0.233 , 0.285	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	5629	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: M38, PG4, TGP, PTR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	B	0.57	0/229	0.83	0/351
2	C	0.43	0/254	0.74	0/390
3	D	0.52	0/494	0.83	0/760
4	A	0.63	0/4783	0.74	2/6420 (0.0%)
All	All	0.61	0/5760	0.75	2/7921 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	D	0	1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
4	A	617	LEU	CA-CB-CG	5.83	128.70	115.30
4	A	335	LEU	CA-CB-CG	5.04	126.89	115.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	D	112	DC	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	203	0	112	10	0
2	C	246	0	138	7	0
3	D	443	0	255	18	0
4	A	4699	0	4726	21	0
5	C	25	0	14	1	0
6	A	13	0	18	1	0
All	All	5629	0	5263	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:17:DA:H2''	2:C:18:DT:H5''	1.44	0.96
4:A:599:GLN:HE22	4:A:765:PHE:H	1.02	0.92
1:B:1:DA:H2'	1:B:2:DA:C8	2.03	0.92
4:A:419:ASN:HD22	4:A:419:ASN:H	1.35	0.74
3:D:121:DT:H2'	3:D:122:DT:H72	1.73	0.71
1:B:7:DA:H2''	1:B:8:DC:O5'	1.92	0.69
3:D:101:DA:H2'	3:D:102:DA:C8	2.26	0.69
1:B:8:DC:H2'	1:B:9:DT:C7	2.24	0.68
2:C:18:DT:H2'	2:C:19:DT:H72	1.75	0.67
2:C:17:DA:C2'	2:C:18:DT:H5''	2.23	0.67
4:A:599:GLN:NE2	4:A:765:PHE:H	1.85	0.66
4:A:740:ILE:H	4:A:740:ILE:HD13	1.61	0.64
4:A:419:ASN:H	4:A:419:ASN:ND2	1.99	0.60
1:B:6:DG:OP2	4:A:424:ILE:HD11	2.03	0.59
4:A:599:GLN:HE22	4:A:765:PHE:N	1.87	0.58
3:D:121:DT:H2''	3:D:122:DT:C6	2.39	0.57
2:C:13:DA:OP2	4:A:636:PRO:HD3	2.05	0.56
2:C:18:DT:H2''	2:C:19:DT:C6	2.40	0.56
3:D:121:DT:H2'	3:D:122:DT:C7	2.37	0.55
1:B:8:DC:H2'	1:B:9:DT:H72	1.88	0.54
3:D:106:DT:H5'	3:D:106:DT:H6	1.72	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:18:DT:H2'	2:C:19:DT:C7	2.39	0.53
4:A:296:ASN:HD21	4:A:299:LYS:HD2	1.74	0.52
2:C:11:TGP:H2'	2:C:12:DG:N7	2.25	0.51
3:D:105:DA:H2''	3:D:106:DT:C5'	2.41	0.51
3:D:115:DG:H2'	3:D:116:DT:C7	2.41	0.51
4:A:505:CYS:SG	4:A:573:LEU:HD12	2.52	0.50
4:A:626:VAL:HG11	4:A:724:LEU:HD21	1.94	0.50
4:A:383:ILE:HG23	4:A:403:GLU:HG3	1.93	0.49
1:B:5:DA:H2''	1:B:6:DG:O5'	2.14	0.48
4:A:511:HIS:HE2	6:A:911:PG4:H81	1.80	0.47
1:B:5:DA:C8	1:B:5:DA:H5'	2.50	0.47
3:D:105:DA:H2''	3:D:106:DT:H5'	1.97	0.47
1:B:5:DA:H5'	1:B:5:DA:H8	1.79	0.47
3:D:118:DT:C6	3:D:119:DT:H72	2.50	0.47
3:D:116:DT:OP1	4:A:488:ARG:HB3	2.15	0.46
4:A:485:LEU:HD21	4:A:541:VAL:HG11	1.96	0.46
3:D:102:DA:H2''	3:D:103:DA:OP2	2.15	0.45
3:D:109:DT:H2''	3:D:110:DT:O5'	2.15	0.45
3:D:115:DG:H2'	3:D:116:DT:H72	1.99	0.45
3:D:112:DC:H2''	3:D:113:DA:H5''	1.97	0.45
4:A:273:ILE:H	4:A:273:ILE:HD12	1.82	0.44
3:D:120:DT:H5'	3:D:120:DT:H6	1.84	0.43
3:D:120:DT:H2''	3:D:121:DT:OP2	2.18	0.43
1:B:4:DA:H1'	1:B:5:DA:H5''	2.00	0.43
5:C:990:M38:H222	5:C:990:M38:H17	2.01	0.43
4:A:254:GLU:O	4:A:258:THR:HG23	2.19	0.42
1:B:8:DC:H5''	4:A:428:MET:SD	2.60	0.42
4:A:541:VAL:HA	4:A:542:PRO:HD3	1.94	0.41
3:D:111:DC:H5'	3:D:111:DC:C6	2.56	0.41
4:A:341:CYS:SG	4:A:429:LEU:HD21	2.60	0.41
4:A:216:LYS:HB3	4:A:435:ILE:HD11	2.03	0.40
3:D:106:DT:H5'	3:D:106:DT:C6	2.53	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
4	A	564/592 (95%)	521 (92%)	33 (6%)	10 (2%)	7 32

All (10) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	A	375	ARG
4	A	636	PRO
4	A	676	LYS
4	A	759	ALA
4	A	213	GLU
4	A	518	LEU
4	A	629	LEU
4	A	641	GLU
4	A	634	ARG
4	A	397	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
4	A	505/535 (94%)	476 (94%)	29 (6%)	17 49

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	A	207	GLU
4	A	210	ARG
4	A	236	GLU
4	A	284	LYS
4	A	288	ASN
4	A	304	GLN
4	A	325	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
4	A	328	LYS
4	A	335	LEU
4	A	397	PRO
4	A	418	GLU
4	A	419	ASN
4	A	430	ASN
4	A	439	LYS
4	A	461	TYR
4	A	498	THR
4	A	557	ASN
4	A	573	LEU
4	A	623	ASN
4	A	629	LEU
4	A	636	PRO
4	A	648	GLN
4	A	696	GLU
4	A	702	GLU
4	A	708	ARG
4	A	718	THR
4	A	721	LEU
4	A	740	ILE
4	A	760	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
4	A	277	ASN
4	A	367	HIS
4	A	419	ASN
4	A	442	GLN
4	A	459	ASN
4	A	460	GLN
4	A	491	ASN
4	A	599	GLN
4	A	611	ASN
4	A	623	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	TGP	C	11	2,3	18,21,25	1.21	2 (11%)	19,31,38	0.75	0
4	PTR	A	723	1,4	15,16,17	1.19	1 (6%)	17,22,24	0.83	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TGP	C	11	2,3	-	0/2/18/22	0/3/3/3
4	PTR	A	723	1,4	-	0/10/11/13	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	11	TGP	C5-C6	-3.32	1.40	1.47
2	C	11	TGP	C8-N7	-2.39	1.31	1.34
4	A	723	PTR	CE1-CD1	2.20	1.42	1.38

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	11	TGP	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	M38	C	990	-	26,28,28	3.45	13 (50%)	31,41,41	1.62	7 (22%)
6	PG4	A	911	-	12,12,12	0.48	0	11,11,11	0.25	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	M38	C	990	-	-	3/6/18/18	0/4/4/4
6	PG4	A	911	-	-	3/10/10/10	-

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	C	990	M38	C8-N10	9.78	1.52	1.39
5	C	990	M38	C7-C6	9.05	1.54	1.41
5	C	990	M38	C13-C8	6.15	1.52	1.41
5	C	990	M38	C14-C12	3.23	1.44	1.38
5	C	990	M38	C16-C17	3.13	1.43	1.36
5	C	990	M38	C2-C1	2.90	1.42	1.36
5	C	990	M38	C3-C4	2.87	1.42	1.36
5	C	990	M38	C12-C11	-2.85	1.48	1.53
5	C	990	M38	C9-N10	2.82	1.42	1.38
5	C	990	M38	C15-C14	2.44	1.43	1.38
5	C	990	M38	C7-C8	-2.15	1.37	1.41

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	C	990	M38	C17-C13	2.07	1.46	1.42
5	C	990	M38	C16-C15	2.06	1.42	1.38

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	C	990	M38	C5-C9-N10	4.76	119.27	116.13
5	C	990	M38	C14-C12-C11	-3.33	129.69	134.59
5	C	990	M38	C13-C12-C11	3.13	109.32	106.92
5	C	990	M38	C7-C8-C13	-2.79	106.48	110.62
5	C	990	M38	C12-C11-C7	-2.57	103.98	105.61
5	C	990	M38	C8-N10-C9	-2.13	119.55	121.72
5	C	990	M38	C22-N10-C8	2.06	121.73	118.99

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	C	990	M38	N10-C22-C23-C25
6	A	911	PG4	O1-C1-C2-O2
6	A	911	PG4	C1-C2-O2-C3
6	A	911	PG4	C4-C3-O2-C2
5	C	990	M38	O20-C24-C25-C23
5	C	990	M38	O19-C24-C25-C23


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	C	990	M38	1	0
6	A	911	PG4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and

any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.