



Full wwPDB X-ray Structure Validation Report ⓘ

May 13, 2020 – 11:18 am BST

PDB ID : 6SJZ
Title : HsNMT1 in complex with both MyrCoA and Acetylated-GNCFKPR substrates
Authors : Dian, C.; Riviere, F.B.; Asensio, T.; Giglione, C.; Meinnel, T.
Deposited on : 2019-08-14
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11

2 Entry composition i

There are 6 unique types of molecules in this entry. The entry contains 7209 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glycylpeptide N-tetradecanoyltransferase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	391	3194	2070	536	571	17	0	6	0
1	B	389	3153	2047	532	558	16	0	7	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	95	GLY	-	expression tag	UNP P30419
A	96	GLY	-	expression tag	UNP P30419
A	97	SER	-	expression tag	UNP P30419
A	98	GLU	-	expression tag	UNP P30419
B	95	GLY	-	expression tag	UNP P30419
B	96	GLY	-	expression tag	UNP P30419
B	97	SER	-	expression tag	UNP P30419
B	98	GLU	-	expression tag	UNP P30419

- Molecule 2 is a protein called Apoptosis-inducing factor 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	E	9	60	37	10	12	1	0	0	0
2	F	8	54	34	9	10	1	0	0	0

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
E	0	ACE	-	acetylation	UNP Q96NN9
E	2	ASN	GLY	engineered mutation	UNP Q96NN9
E	8	ARG	LYS	conflict	UNP Q96NN9
F	0	ACE	-	acetylation	UNP Q96NN9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	2	ASN	GLY	engineered mutation	UNP Q96NN9
F	8	ARG	LYS	conflict	UNP Q96NN9

- Molecule 3 is TETRADECANOYL-COA (three-letter code: MYA) (formula: $C_{35}H_{62}N_7O_{17}P_3S$) (labeled as "Ligand of Interest" by author).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	N	O	P			S
3	A	1	Total	C	N	O	P	S	0	0
			63	35	7	17	3	1		
3	B	1	Total	C	N	O	P	S	0	0
			63	35	7	17	3	1		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total Cl 1 1	0	0
4	A	1	Total Cl 1 1	0	0

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	B	1	Total C O 6 3 3	0	0
5	B	1	Total C O 6 3 3	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	327	Total O 327 327	0	1
6	B	263	Total O 263 263	0	0
6	E	3	Total O 3 3	0	0
6	F	3	Total O 3 3	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Glycylpeptide N-tetradecanoyltransferase 1




- Molecule 1: Glycylpeptide N-tetradecanoyltransferase 1



- Molecule 2: Apoptosis-inducing factor 3



- Molecule 2: Apoptosis-inducing factor 3



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	79.91Å 178.90Å 58.38Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.89 – 2.00 48.89 – 1.99	Depositor EDS
% Data completeness (in resolution range)	99.3 (48.89-2.00) 99.5 (48.89-1.99)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.33 (at 2.00Å)	Xtrriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.172 , 0.202 0.174 , 0.203	Depositor DCC
R_{free} test set	2845 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å ²)	21.7	Xtrriage
Anisotropy	0.506	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 50.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7209	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 18.32% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MYA, GOL, ACE, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.28	1/3284 (0.0%)	0.46	0/4468
1	B	0.27	0/3242	0.46	0/4417
2	E	0.31	0/59	0.85	0/77
2	F	0.35	0/53	0.60	0/70
All	All	0.28	1/6638 (0.0%)	0.46	0/9032

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	123	GLN	C-N	5.02	1.43	1.34

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3194	0	3129	27	0
1	B	3153	0	3086	23	0
2	E	60	0	53	6	0
2	F	54	0	51	10	0
3	A	63	0	58	0	0
3	B	63	0	58	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	12	0	16	1	0
5	B	12	0	16	1	0
6	A	327	0	0	3	0
6	B	263	0	0	2	0
6	E	3	0	0	0	0
6	F	3	0	0	0	0
All	All	7209	0	6467	52	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (52) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:403:LEU:HD23	2:E:0:ACE:H3	1.48	0.95
1:A:496:GLN:HE22	5:A:503:GOL:H11	1.48	0.78
1:A:107:LYS:NZ	6:A:601:HOH:O	2.23	0.72
1:A:403:LEU:HD23	2:F:0:ACE:H3	1.71	0.70
1:B:403:LEU:CD2	2:E:0:ACE:H3	2.22	0.69
1:A:177[B]:ASN:ND2	1:A:191:ASP:OD1	2.33	0.62
1:B:420:TYR:CE2	2:E:0:ACE:H1	2.37	0.59
1:A:177[A]:ASN:ND2	6:A:607:HOH:O	2.36	0.58
1:B:410:HIS:CD2	1:B:413:HIS:H	2.22	0.56
1:A:474:LEU:HD13	2:F:3:CYS:HB3	1.87	0.56
1:A:170:LYS:HA	1:A:173[B]:TYR:CE2	2.41	0.55
1:B:458:ASN:HA	1:B:461:PHE:CE2	2.42	0.54
1:A:145:ILE:HD12	1:A:272:HIS:HB3	1.90	0.53
1:B:496:GLN:HE22	5:B:503:GOL:H11	1.74	0.53
1:A:355[A]:ARG:NH1	1:B:143:ASP:OD2	2.43	0.52
1:A:351:GLN:O	1:A:355[A]:ARG:HG2	2.09	0.52
1:B:319:THR:HG23	1:B:322:ARG:H	1.73	0.52
1:B:410:HIS:HD2	1:B:412:THR:H	1.57	0.52
1:A:403:LEU:CD2	2:F:0:ACE:H3	2.41	0.51
1:A:420:TYR:CE2	2:F:0:ACE:H1	2.47	0.50
1:A:458:ASN:HA	1:A:461:PHE:CE2	2.46	0.50
1:B:130:GLU:OE2	6:B:601:HOH:O	2.20	0.49
6:B:667:HOH:O	2:E:2:ASN:HB3	2.11	0.49
1:A:405:SER:HG	2:F:4:PHE:HE2	1.55	0.49
1:B:145:ILE:HD12	1:B:272:HIS:HB3	1.94	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:311:PHE:CE1	2:F:6:LYS:HB3	2.48	0.48
2:F:1:GLY:O	2:F:3:CYS:N	2.39	0.48
1:A:282:THR:C	2:F:2:ASN:HD21	2.17	0.47
1:B:410:HIS:CD2	1:B:412:THR:H	2.31	0.47
1:A:162:ASP:HA	6:A:801:HOH:O	2.15	0.46
1:A:355[A]:ARG:HG2	1:A:355[A]:ARG:H	1.60	0.46
1:A:410:HIS:HB3	1:A:413:HIS:CD2	2.52	0.45
1:A:303:PRO:O	1:A:307:ILE:HG12	2.16	0.45
1:A:188:PHE:HB2	2:F:4:PHE:CZ	2.52	0.45
1:B:259:VAL:O	1:B:262[B]:VAL:HG22	2.17	0.45
1:A:413:HIS:N	1:A:413:HIS:CD2	2.84	0.45
1:B:195:GLU:HB3	1:B:381:ILE:HD11	1.97	0.45
1:B:405:SER:OG	2:E:4:PHE:HE2	1.99	0.45
1:B:182:GLU:OE2	1:B:189:ARG:NH1	2.45	0.44
1:B:303:PRO:O	1:B:307:ILE:HG12	2.17	0.44
1:B:251:HIS:CE1	1:B:253:LYS:HG2	2.53	0.43
1:A:195:GLU:HB3	1:A:381:ILE:HD11	1.99	0.43
1:B:401:TYR:CE1	2:E:0:ACE:H2	2.52	0.43
1:A:170:LYS:HA	1:A:173[B]:TYR:CD2	2.54	0.43
1:A:417:LYS:HG2	1:A:447:PHE:HA	2.02	0.42
1:A:130:GLU:O	1:A:289:LYS:HE3	2.20	0.41
1:B:410:HIS:HD2	1:B:413:HIS:H	1.68	0.41
1:B:289:LYS:HA	1:B:290:PRO:HD3	1.95	0.41
2:F:4:PHE:HD1	2:F:4:PHE:C	2.24	0.41
1:B:458:ASN:O	1:B:462:LEU:HG	2.21	0.41
1:B:288:PRO:HA	1:B:289:LYS:HA	1.95	0.40
1:A:314:LEU:HD21	1:A:318:MET:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	393/402 (98%)	385 (98%)	7 (2%)	1 (0%)	41	37
1	B	392/402 (98%)	382 (97%)	10 (3%)	0	100	100
2	E	7/9 (78%)	6 (86%)	1 (14%)	0	100	100
2	F	6/9 (67%)	5 (83%)	1 (17%)	0	100	100
All	All	798/822 (97%)	778 (98%)	19 (2%)	1 (0%)	51	49

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	314	LEU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	346/362 (96%)	340 (98%)	6 (2%)	60	65
1	B	338/362 (93%)	332 (98%)	6 (2%)	59	63
2	E	6/7 (86%)	5 (83%)	1 (17%)	2	1
2	F	6/7 (86%)	2 (33%)	4 (67%)	0	0
All	All	696/738 (94%)	679 (98%)	17 (2%)	50	51

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	108	THR
1	A	318	MET
1	A	334	LYS
1	A	355[A]	ARG
1	A	355[B]	ARG
1	A	455	LEU
1	B	108	THR
1	B	113	SER
1	B	115	ARG
1	B	240	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	322	ARG
1	B	455	LEU
2	E	3	CYS
2	F	2	ASN
2	F	3	CYS
2	F	4	PHE
2	F	6	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	413	HIS
1	A	496	GLN
1	B	147	GLN
1	B	410	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 2 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	MYA	B	501	-	54,65,65	1.31	5 (9%)	67,91,91	1.60	9 (13%)
5	GOL	B	503	-	5,5,5	0.37	0	5,5,5	0.20	0
5	GOL	A	503	-	5,5,5	0.39	0	5,5,5	0.20	0
5	GOL	A	504	-	5,5,5	0.38	0	5,5,5	0.20	0
5	GOL	B	502	-	5,5,5	0.37	0	5,5,5	0.20	0
3	MYA	A	501	-	54,65,65	1.30	6 (11%)	67,91,91	1.57	8 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MYA	B	501	-	-	7/59/80/80	0/3/3/3
5	GOL	B	503	-	-	2/4/4/4	-
5	GOL	A	503	-	-	2/4/4/4	-
5	GOL	A	504	-	-	0/4/4/4	-
5	GOL	B	502	-	-	0/4/4/4	-
3	MYA	A	501	-	-	6/59/80/80	0/3/3/3

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	501	MYA	C2A-N3A	5.06	1.40	1.32
3	A	501	MYA	C2A-N3A	4.67	1.39	1.32
3	B	501	MYA	O4X-C1X	3.55	1.46	1.41
3	B	501	MYA	C2A-N1A	3.21	1.39	1.33
3	A	501	MYA	C2A-N1A	3.20	1.39	1.33
3	A	501	MYA	O4X-C1X	2.79	1.45	1.41
3	A	501	MYA	C5A-C4A	-2.69	1.33	1.40
3	B	501	MYA	C5A-C4A	-2.54	1.34	1.40
3	A	501	MYA	C6A-C5A	-2.51	1.34	1.43
3	B	501	MYA	C6A-C5A	-2.50	1.34	1.43
3	A	501	MYA	P3X-O8A	-2.31	1.46	1.54

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	501	MYA	N3A-C2A-N1A	-7.28	117.30	128.68
3	B	501	MYA	O2M-C2M-C3M	6.70	120.95	109.02
3	B	501	MYA	N3A-C2A-N1A	-6.56	118.42	128.68

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	501	MYA	O2M-C2M-C3M	6.26	120.16	109.02
3	B	501	MYA	C2X-C3X-C4X	-3.21	97.54	103.22
3	B	501	MYA	O4X-C1X-C2X	-2.79	102.85	106.93
3	A	501	MYA	O4X-C1X-C2X	-2.61	103.11	106.93
3	A	501	MYA	C2X-C3X-C4X	-2.47	98.85	103.22
3	A	501	MYA	O4X-C4X-C5X	-2.44	101.34	109.37
3	B	501	MYA	C5A-C6A-N6A	-2.41	116.69	120.35
3	B	501	MYA	O6A-P2A-O4A	2.33	118.19	109.07
3	B	501	MYA	P2A-O3A-P1A	-2.33	124.83	132.83
3	A	501	MYA	C4A-C5A-N7A	-2.28	107.02	109.40
3	A	501	MYA	P2A-O3A-P1A	-2.27	125.03	132.83
3	B	501	MYA	O5A-P2A-O4A	2.17	122.97	112.24
3	A	501	MYA	C5A-C6A-N6A	-2.17	117.06	120.35
3	B	501	MYA	O5A-P2A-O6A	2.09	117.44	107.75

There are no chirality outliers.

All (17) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	B	501	MYA	C12-O6A-P2A-O4A
3	A	501	MYA	C5X-O5X-P1A-O2A
5	B	503	GOL	O1-C1-C2-C3
5	A	503	GOL	O1-C1-C2-C3
5	A	503	GOL	O1-C1-C2-O2
5	B	503	GOL	O1-C1-C2-O2
3	A	501	MYA	C5X-O5X-P1A-O3A
3	B	501	MYA	CAM-CBM-CCM-CDM
3	A	501	MYA	S1-C2M-C3M-C4M
3	B	501	MYA	P1A-O3A-P2A-O5A
3	A	501	MYA	C3X-C4X-C5X-O5X
3	B	501	MYA	C8M-C9M-CAM-CBM
3	A	501	MYA	C5M-C6M-C7M-C8M
3	A	501	MYA	O4X-C4X-C5X-O5X
3	B	501	MYA	C11-C12-O6A-P2A
3	B	501	MYA	C5X-O5X-P1A-O2A
3	B	501	MYA	S1-C2M-C3M-C4M

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	503	GOL	1	0
5	A	503	GOL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	391/402 (97%)	0.27	29 (7%) 14 13	9, 21, 57, 80	0
1	B	389/402 (96%)	0.26	21 (5%) 25 24	12, 26, 55, 70	0
2	E	8/9 (88%)	2.94	7 (87%) 0 0	41, 46, 50, 64	0
2	F	7/9 (77%)	6.38	7 (100%) 0 0	44, 54, 59, 59	0
All	All	795/822 (96%)	0.35	64 (8%) 12 11	9, 23, 56, 80	0

All (64) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	F	3	CYS	9.3
2	F	7	PRO	8.0
2	F	4	PHE	7.5
2	F	6	LYS	6.2
2	F	5	SER	6.2
1	A	412	THR	5.9
1	A	105	PRO	4.9
1	B	113	SER	4.8
1	A	409	ASN	4.7
1	A	410	HIS	4.5
1	A	313	HIS	4.4
1	B	108	THR	4.4
2	F	2	ASN	4.4
2	E	7	PRO	4.2
1	A	184	ASP	4.1
1	A	106	ALA	3.9
2	E	8	ARG	3.9
1	B	114	LYS	3.8
1	A	411	PRO	3.7
2	E	1	GLY	3.7
1	A	408	MET	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	114	LYS	3.4
1	B	116	SER	3.4
1	A	108	THR	3.4
1	B	313	HIS	3.4
1	A	314	LEU	3.4
1	A	307	ILE	3.3
1	B	320	MET	3.3
1	A	107	LYS	3.2
1	A	185	ASP	3.1
1	A	315	SER	3.1
2	F	1	GLY	3.1
1	A	312	SER	3.0
2	E	4	PHE	3.0
1	B	324	MET	2.9
2	E	5	SER	2.8
1	B	112	ALA	2.8
1	A	254	LEU	2.6
1	A	309	VAL	2.6
1	B	309	VAL	2.6
1	A	109	MET	2.5
1	A	326	LEU	2.5
1	A	187	MET	2.4
1	B	115	ARG	2.4
1	B	307	ILE	2.4
1	A	318	MET	2.4
1	B	154	GLN	2.4
1	A	413	HIS	2.3
1	B	319	THR	2.3
1	B	444	MET	2.3
1	A	188	PHE	2.3
1	A	115	ARG	2.3
1	A	186	ASN	2.2
1	B	323	THR	2.2
1	B	256	SER	2.2
1	B	128	LEU	2.2
1	B	105	PRO	2.1
2	E	2	ASN	2.1
2	E	6	LYS	2.1
1	B	322	ARG	2.0
1	A	183	ASP	2.0
1	B	471	ASP	2.0
1	B	312	SER	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	111	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	GOL	B	503	6/6	0.89	0.18	25,30,33,34	0
3	MYA	B	501	63/63	0.90	0.15	12,33,47,51	0
5	GOL	A	503	6/6	0.91	0.14	27,30,31,32	0
3	MYA	A	501	63/63	0.94	0.11	9,23,30,31	0
4	CL	A	502	1/1	0.95	0.08	39,39,39,39	0
5	GOL	B	502	6/6	0.96	0.09	20,24,29,30	0
5	GOL	A	504	6/6	0.96	0.10	13,17,20,25	0
4	CL	B	504	1/1	0.98	0.13	39,39,39,39	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around MYA B 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.