



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 10, 2023 – 02:53 AM EDT

PDB ID : 7SMV  
Title : Crystallization of feline coronavirus Mpro with GC376 reveals mechanism of inhibition  
Authors : Khan, M.B.; Lu, J.; Young, H.S.; Lemieux, M.J.  
Deposited on : 2021-10-26  
Resolution : 1.93 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 1.93 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4310 (1.96-1.92)
Clashscore	141614	1023 (1.94-1.94)
Ramachandran outliers	138981	1007 (1.94-1.94)
Sidechain outliers	138945	1007 (1.94-1.94)
RSRZ outliers	127900	4250 (1.96-1.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	302	
1	B	302	

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 9504 atoms, of which 4564 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 3C-like proteinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	299	Total	C	H	N	O	S	0	0	0
			4548	1454	2249	390	439	16			
1	B	299	Total	C	H	N	O	S	0	0	0
			4554	1454	2255	390	439	16			

- Molecule 2 is N 2 -[(benzyloxy)carbonyl]-N-[(2S)-1-hydroxy-3-[(3S)-2-oxopyrrolidin-3-yl]propan-2-yl]-L-leucinamide (three-letter code: UED) (formula: C<sub>21</sub>H<sub>31</sub>N<sub>3</sub>O<sub>5</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	N	O		
2	A	1	Total	C	H	N	O	0	0
			59	21	30	3	5		
2	B	1	Total	C	H	N	O	0	0
			59	21	30	3	5		


- Molecule 3 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
3	A	112	Total 112	O 112	0	0
3	B	172	Total 172	O 172	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 3C-like proteinase



- Molecule 1: 3C-like proteinase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	67.80Å 77.42Å 111.75Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.31 – 1.93 45.31 – 1.93	Depositor EDS
% Data completeness (in resolution range)	99.9 (45.31-1.93) 99.9 (45.31-1.93)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.13 (at 1.94Å)	Xtrriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, $R_{free}$	0.191 , 0.245 0.193 , 0.246	Depositor DCC
$R_{free}$ test set	2247 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	39.2	Xtrriage
Anisotropy	0.359	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 45.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	9504	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	50.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.78% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: UED

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.93	8/2346 (0.3%)	1.45	20/3179 (0.6%)
1	B	0.74	0/2346	0.86	3/3179 (0.1%)
All	All	0.84	8/4692 (0.2%)	1.19	23/6358 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	5

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	234	LYS	CB-CG	14.33	1.91	1.52
1	A	234	LYS	CE-NZ	12.73	1.80	1.49
1	A	61	ARG	NE-CZ	-12.24	1.17	1.33
1	A	234	LYS	CD-CE	10.17	1.76	1.51
1	A	61	ARG	N-CA	8.72	1.63	1.46
1	A	227	GLU	CB-CG	7.13	1.65	1.52
1	A	234	LYS	CG-CD	6.55	1.74	1.52
1	A	61	ARG	CD-NE	5.78	1.56	1.46

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	227	GLU	OE1-CD-OE2	-32.05	84.83	123.30
1	A	61	ARG	CB-CG-CD	25.91	178.96	111.60
1	A	61	ARG	CD-NE-CZ	-23.67	90.46	123.60

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	227	GLU	CG-CD-OE1	22.16	162.62	118.30
1	A	61	ARG	CA-CB-CG	-19.26	71.03	113.40
1	B	277	ILE	CA-CB-CG1	-17.11	78.48	111.00
1	A	234	LYS	CB-CG-CD	-16.42	68.91	111.60
1	A	227	GLU	CG-CD-OE2	-14.92	88.46	118.30
1	A	61	ARG	NE-CZ-NH2	14.14	127.37	120.30
1	A	61	ARG	NH1-CZ-NH2	-14.12	103.87	119.40
1	A	234	LYS	N-CA-CB	-13.53	86.25	110.60
1	A	234	LYS	CD-CE-NZ	-11.66	84.88	111.70
1	A	234	LYS	CB-CA-C	9.12	128.64	110.40
1	A	61	ARG	CB-CA-C	8.92	128.25	110.40
1	B	277	ILE	CG1-CB-CG2	7.89	128.75	111.40
1	A	50	VAL	CA-CB-CG2	6.84	121.17	110.90
1	A	61	ARG	N-CA-C	-6.21	94.22	111.00
1	A	234	LYS	CA-CB-CG	-5.51	101.28	113.40
1	A	240	GLU	CA-CB-CG	-5.45	101.41	113.40
1	B	186	ASP	CB-CG-OD2	-5.39	113.45	118.30
1	A	226	LEU	CA-CB-CG	5.10	127.03	115.30
1	A	50	VAL	N-CA-CB	5.10	122.71	111.50
1	A	234	LYS	N-CA-C	5.02	124.54	111.00

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	226	LEU	Peptide
1	A	227	GLU	Sidechain
1	A	233	ALA	Peptide,Mainchain
1	A	61	ARG	Sidechain

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2299	2249	2256	61	0
1	B	2299	2255	2256	29	0
2	A	29	30	0	0	0

Continued on next page...



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	29	30	0	0	0
3	A	112	0	0	5	0
3	B	172	0	0	4	1
All	All	4940	4564	4512	89	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (89) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:234:LYS:CD	1:A:234:LYS:CG	1.74	1.60
1:A:234:LYS:CD	1:A:234:LYS:CE	1.76	1.60
1:A:234:LYS:CG	1:A:234:LYS:CB	1.91	1.48
1:A:234:LYS:CE	1:A:234:LYS:NZ	1.80	1.41
1:A:234:LYS:CD	1:A:234:LYS:CB	2.07	1.33
1:A:60:VAL:O	1:A:61:ARG:HD3	1.37	1.19
1:A:234:LYS:HD2	1:A:234:LYS:HB3	1.21	1.18
1:A:234:LYS:CB	1:A:234:LYS:HD2	1.75	1.13
1:A:225:THR:HG22	1:A:227:GLU:H	1.17	1.10
1:B:217:TRP:HZ2	1:B:277:ILE:HD11	1.17	1.08
1:B:276:THR:C	1:B:277:ILE:HG13	1.70	1.06
1:B:272:PHE:CE1	1:B:277:ILE:HD12	1.91	1.06
1:A:234:LYS:CD	1:A:234:LYS:HB3	1.80	1.05
1:B:120:CYS:SG	3:B:668:HOH:O	2.15	1.02
1:B:217:TRP:CZ2	1:B:277:ILE:HD11	1.94	1.01
1:A:60:VAL:O	1:A:61:ARG:CD	2.07	1.01
1:A:60:VAL:C	1:A:61:ARG:HD3	1.90	0.92
1:A:225:THR:HG22	1:A:227:GLU:N	1.90	0.87
1:A:234:LYS:CD	1:A:234:LYS:NZ	2.41	0.84
1:B:272:PHE:CD1	1:B:277:ILE:HD12	2.12	0.84
1:A:120:CYS:SG	3:A:603:HOH:O	2.36	0.83
1:B:272:PHE:CE1	1:B:277:ILE:CD1	2.64	0.80
1:B:276:THR:O	1:B:277:ILE:HG13	1.81	0.79
1:A:225:THR:CG2	1:A:227:GLU:H	1.96	0.78
1:A:69:LYS:HE3	3:A:536:HOH:O	1.83	0.77
1:B:264:CYS:O	1:B:268:LEU:HD13	1.85	0.76
1:A:234:LYS:HG3	1:A:240:GLU:OE1	1.86	0.76
1:B:248:ASN:O	3:B:501:HOH:O	2.10	0.69
1:A:234:LYS:CG	1:A:234:LYS:CA	2.72	0.67
1:A:201:ASP:OD1	3:A:501:HOH:O	2.14	0.65

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:6:MET:HE3	1:A:295:GLN:HG3	1.80	0.62
1:A:234:LYS:CG	1:A:240:GLU:OE1	2.48	0.61
1:A:120:CYS:SG	3:A:605:HOH:O	2.57	0.61
3:A:530:HOH:O	1:B:1:SER:HB2	2.00	0.61
1:A:211:LEU:HD13	1:A:256:TYR:HD2	1.67	0.59
1:B:49:ARG:HE	1:B:50:VAL:H	1.50	0.58
1:A:57:LEU:HA	1:A:60:VAL:HG23	1.85	0.57
1:A:59:SER:C	1:A:61:ARG:NH1	2.58	0.56
1:A:230:ASN:O	1:A:234:LYS:HB2	2.05	0.56
1:A:211:LEU:HD13	1:A:256:TYR:CD2	2.41	0.55
1:B:276:THR:C	1:B:277:ILE:CG1	2.60	0.55
1:A:60:VAL:HG12	1:A:61:ARG:N	2.21	0.55
1:A:61:ARG:HB3	1:A:63:HIS:CE1	2.43	0.54
1:A:69:LYS:HG3	1:A:69:LYS:O	2.08	0.52
1:A:112:ASN:O	1:A:148:GLY:HA2	2.09	0.52
1:A:130:ARG:HH22	1:A:136:LYS:HE2	1.75	0.52
1:A:57:LEU:HA	1:A:60:VAL:CG2	2.40	0.51
1:B:272:PHE:CD1	1:B:277:ILE:CD1	2.89	0.50
1:A:130:ARG:NH2	1:A:136:LYS:HE2	2.26	0.50
1:A:216:ARG:O	1:A:219:VAL:HG12	2.10	0.50
1:A:19:ARG:O	1:A:67:ILE:HA	2.12	0.49
1:A:189:SER:O	1:A:190:MET:HG2	2.12	0.49
1:A:220:THR:OG1	1:A:222:THR:OG1	2.29	0.49
1:A:189:SER:O	1:A:190:MET:CG	2.61	0.48
1:B:112:ASN:O	1:B:148:GLY:HA2	2.13	0.48
1:A:61:ARG:HD3	1:A:61:ARG:HH11	0.95	0.48
1:B:253:LYS:HD2	1:B:297:TYR:CZ	2.48	0.48
1:B:40:ARG:HG2	1:B:84:VAL:O	2.15	0.47
1:B:6:MET:CE	1:B:295:GLN:CD	2.83	0.47
1:B:190:MET:HA	1:B:190:MET:CE	2.46	0.46
1:A:190:MET:HG3	1:A:190:MET:O	2.16	0.46
1:B:6:MET:HE1	1:B:295:GLN:CG	2.45	0.46
1:A:138:SER:O	1:A:138:SER:OG	2.34	0.46
1:A:225:THR:C	1:A:227:GLU:N	2.68	0.45
1:A:272:PHE:CE2	1:A:277:ILE:HD12	2.52	0.45
1:B:6:MET:HE1	1:B:295:GLN:HG3	1.99	0.45
1:A:189:SER:O	1:A:190:MET:CB	2.64	0.44
1:B:31:TRP:CE2	1:B:74:LEU:HD11	2.53	0.44
1:A:208:TYR:O	1:A:212:ILE:HG13	2.17	0.44
1:B:49:ARG:NE	1:B:50:VAL:H	2.15	0.44
1:B:201:ASP:OD1	3:B:502:HOH:O	2.21	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:56:GLU:O	1:A:60:VAL:HG23	2.18	0.43
1:A:27:LEU:HD12	1:A:27:LEU:C	2.38	0.43
1:B:6:MET:HE1	1:B:295:GLN:CD	2.39	0.43
1:A:57:LEU:CA	1:A:60:VAL:HG23	2.48	0.43
1:A:231:SER:HA	1:A:234:LYS:HE3	2.01	0.43
1:A:240:GLU:H	1:A:240:GLU:HG3	1.53	0.43
1:A:225:THR:HG22	1:A:228:SER:H	1.84	0.42
1:B:296:MET:HB3	1:B:296:MET:HE2	1.88	0.42
1:A:263:GLU:HG2	1:A:267:ARG:HH21	1.83	0.42
1:A:59:SER:O	1:A:61:ARG:CZ	2.68	0.42
1:A:231:SER:O	1:A:234:LYS:HD2	2.19	0.42
1:A:60:VAL:O	1:A:61:ARG:HG2	2.14	0.41
1:B:107:PRO:HA	1:B:129:MET:HG3	2.03	0.41
1:B:136:LYS:HD2	3:B:525:HOH:O	2.21	0.41
1:A:56:GLU:O	1:A:60:VAL:CG2	2.69	0.40
1:A:49:ARG:O	1:A:50:VAL:C	2.60	0.40
1:A:264:CYS:O	1:A:268:LEU:HG	2.21	0.40
1:A:125:TYR:CD1	1:B:6:MET:HG2	2.57	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:608:HOH:O	3:B:645:HOH:O[4_445]	2.16	0.04

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	297/302 (98%)	287 (97%)	8 (3%)	2 (1%)	22	11
1	B	297/302 (98%)	293 (99%)	4 (1%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	594/604 (98%)	580 (98%)	12 (2%)	2 (0%)	41 32

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	190	MET
1	A	234	LYS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	254/257 (99%)	243 (96%)	11 (4%)	29 14
1	B	254/257 (99%)	251 (99%)	3 (1%)	71 64
All	All	508/514 (99%)	494 (97%)	14 (3%)	43 29

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	27	LEU
1	A	69	LYS
1	A	80	LYS
1	A	123	SER
1	A	130	ARG
1	A	190	MET
1	A	220	THR
1	A	222	THR
1	A	234	LYS
1	A	269	ASN
1	A	285	ASP
1	B	45	SER
1	B	60	VAL
1	B	123	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	269	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	UED	A	401	1	30,30,30	4.66	14 (46%)	37,39,39	1.56	8 (21%)
2	UED	B	401	1	30,30,30	4.33	10 (33%)	37,39,39	1.93	10 (27%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	UED	A	401	1	-	1/27/37/37	0/2/2/2
2	UED	B	401	1	-	2/27/37/37	0/2/2/2

All (24) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	UED	C29-N28	19.21	1.54	1.33
2	B	401	UED	C29-N28	17.47	1.52	1.33
2	A	401	UED	C3-C2	7.08	1.53	1.38
2	A	401	UED	C6-C1	6.59	1.53	1.38
2	B	401	UED	C3-C2	6.50	1.52	1.38
2	B	401	UED	C6-C1	6.49	1.52	1.38
2	B	401	UED	C17-N19	6.10	1.47	1.34
2	A	401	UED	C5-C4	5.88	1.53	1.38
2	A	401	UED	C17-N19	5.75	1.46	1.34
2	B	401	UED	C5-C4	5.48	1.52	1.38
2	A	401	UED	C9-N11	4.79	1.46	1.34
2	A	401	UED	C27-N28	4.76	1.56	1.46
2	B	401	UED	O8-C9	4.73	1.44	1.35
2	B	401	UED	C27-N28	4.44	1.55	1.46
2	B	401	UED	C9-N11	4.43	1.45	1.34
2	A	401	UED	O8-C9	3.60	1.42	1.35
2	B	401	UED	C26-C25	-3.37	1.45	1.54
2	A	401	UED	C24-C25	3.19	1.61	1.53
2	A	401	UED	C26-C25	-3.18	1.45	1.54
2	A	401	UED	O18-C17	-2.83	1.17	1.23
2	A	401	UED	O30-C29	-2.78	1.17	1.23
2	A	401	UED	C5-C6	-2.70	1.33	1.38
2	B	401	UED	C5-C6	-2.68	1.33	1.38
2	A	401	UED	C2-C1	-2.48	1.33	1.38

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	401	UED	C24-C20-C21	-4.56	105.31	111.65
2	B	401	UED	O10-C9-N11	-4.50	117.48	124.85
2	A	401	UED	C27-N28-C29	-4.31	105.38	113.84
2	B	401	UED	O22-C21-C20	-3.79	102.16	111.95
2	B	401	UED	O8-C9-N11	3.28	117.17	110.50
2	A	401	UED	C26-C25-C29	3.02	106.80	102.88
2	A	401	UED	O10-C9-N11	-2.97	119.98	124.85
2	B	401	UED	C26-C25-C29	2.94	106.70	102.88
2	A	401	UED	C27-C26-C25	2.86	110.40	105.75
2	B	401	UED	C27-C26-C25	2.83	110.33	105.75
2	B	401	UED	C13-C12-N11	-2.82	104.07	110.58
2	B	401	UED	C20-N19-C17	-2.78	118.77	123.20
2	B	401	UED	C27-N28-C29	-2.50	108.94	113.84
2	A	401	UED	O8-C9-N11	2.43	115.44	110.50

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	UED	O22-C21-C20	-2.16	106.36	111.95
2	B	401	UED	O18-C17-N19	-2.16	118.94	122.93
2	A	401	UED	C24-C25-C26	-2.13	109.49	117.31
2	A	401	UED	C14-C13-C12	-2.10	109.67	115.43

There are no chirality outliers.



All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	401	UED	C12-C13-C14-C16
2	B	401	UED	N11-C12-C13-C14
2	A	401	UED	N11-C12-C13-C14

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	299/302 (99%)	0.76	26 (8%) 10 15	30, 47, 71, 87	0
1	B	299/302 (99%)	0.52	11 (3%) 41 49	29, 40, 67, 90	0
All	All	598/604 (99%)	0.64	37 (6%) 20 27	29, 44, 69, 90	0

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	277	ILE	6.9
1	A	60	VAL	5.3
1	A	299	VAL	5.1
1	A	189	SER	4.9
1	B	280	TYR	4.6
1	A	216	ARG	4.5
1	A	217	TRP	4.1
1	A	61	ARG	3.9
1	A	190	MET	3.8
1	B	2	GLY	3.6
1	A	278	LEU	3.6
1	B	1	SER	3.6
1	A	226	LEU	3.5
1	B	192	LEU	3.5
1	A	223	SER	3.5
1	A	234	LYS	3.4
1	A	252	ALA	3.4
1	A	1	SER	3.3
1	A	280	TYR	2.9
1	A	214	GLY	2.9
1	A	219	VAL	2.8
1	B	273	GLY	2.6
1	A	50	VAL	2.5
1	A	258	VAL	2.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	240	GLU	2.4
1	A	221	ASN	2.3
1	B	190	MET	2.3
1	A	192	LEU	2.2
1	B	216	ARG	2.2
1	A	117	TYR	2.2
1	A	222	THR	2.2
1	B	3	LEU	2.1
1	A	279	SER	2.1
1	B	220	THR	2.1
1	B	211	LEU	2.1
1	A	57	LEU	2.1
1	A	272	PHE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	UED	A	401	29/29	0.92	0.14	39,48,58,61	0
2	UED	B	401	29/29	0.93	0.12	32,41,52,54	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around UED A 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.