

# Full wwPDB X-ray Structure Validation Report (i)

#### Apr 22, 2024 – 04:14 PM EDT

:	8SM1
:	CRYSTAL STRUCTURE OF HUMAN ANTIBODY 769A9 IN COMPLEX
	WITH EPSTEIN-BARR VIRUS MAJOR GLYCOPROTEIN GP350
:	Chen, WH.; Bu, W.; Cohen, J.I.; Kanekiyo, M.; Joyce, M.G.
:	2023-04-25
:	3.29 Å(reported)
	::

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36.2
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36.2

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY \, DIFFRACTION$ 

The reported resolution of this entry is 3.29 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Motrie	Whole archive	Similar resolution
WIEUTIC	$(\# { m Entries})$	$(\# { m Entries},  { m resolution}  { m range}({ m \AA}))$
$R_{free}$	130704	1149 (3.34-3.26)
Clashscore	141614	1205 (3.34-3.26)
Ramachandran outliers	138981	1183 (3.34-3.26)
Sidechain outliers	138945	1182 (3.34-3.26)
RSRZ outliers	127900	1115 (3.34-3.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain						
1	G	431	76%	20%	••				
2	Н	228	6%           67%         22%	•	9%				
3	L	215	80%	18%	•				

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:



Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	NAG	G	502	-	-	-	Х
4	NAG	G	503	-	-	-	Х
4	NAG	G	505	-	-	-	Х
4	NAG	G	506	-	-	-	Х



#### 8SM1

# 2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 6513 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Envelope glycoprotein gp350.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	G	420	Total 3201	C 2024	N 519	0 641	S 17	0	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
G	426	HIS	-	expression tag	UNP P03200
G	427	HIS	-	expression tag	UNP P03200
G	428	HIS	-	expression tag	UNP P03200
G	429	HIS	-	expression tag	UNP P03200
G	430	HIS	-	expression tag	UNP P03200
G	431	HIS	-	expression tag	UNP P03200

• Molecule 2 is a protein called 769A9 Fab heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	Н	208	Total 1551	C 982	N 256	O 308	${ m S}{ m 5}$	0	0	0

• Molecule 3 is a protein called 769A9 Fab light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	L	214	Total 1642	C 1030	N 277	0 331	${S \atop 4}$	0	0	0

• Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	G	1	Total C N O $14$ 8 1 5	0	0
4	G	1	TotalCNO	0	0
		-	14 8 1 5 Total C N O		
4	G	1	14 8 1 5	0	0
4	G	1	$\begin{array}{cccc} \text{Total} & \text{C} & \text{N} & \text{O} \\ 14 & 8 & 1 & 5 \end{array}$	0	0
4	G	1	Total C N O	0	0
	~		$\begin{array}{c ccccccccccccccccccccccccccccccccccc$		-
4	G	1	14 8 1 5	0	0
4	G	1	$\begin{array}{cccc} \text{Total} & \text{C} & \text{N} & \text{O} \\ 14 & 8 & 1 & 5 \end{array}$	0	0
4	G	1	Total C N O	0	0
		-	14  8  1  5		

• Molecule 5 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	G	1	Total 6	${ m C} { m 3}$	O 3	0	0

• Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	G	1	Total O 1 1	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Envelope glycoprotein gp350





## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants	164.08Å 54.68Å 137.22Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $111.71^{\circ}$ $90.00^{\circ}$	Depositor
Bosolution(Å)	45.78 - 3.29	Depositor
Resolution (A)	45.78 - 3.28	EDS
% Data completeness	82.9 (45.78-3.29)	Depositor
(in resolution range)	82.9 (45.78-3.28)	EDS
$R_{merge}$	0.13	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	$2.18 (at 3.25 \text{\AA})$	Xtriage
Refinement program	PHENIX 1.11.1_2575	Depositor
D D .	0.258 , $0.298$	Depositor
$n, n_{free}$	0.278 , $0.323$	DCC
$R_{free}$ test set	732 reflections $(5.00\%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	74.9	Xtriage
Anisotropy	0.322	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.34 , $67.4$	EDS
L-test for $twinning^2$	$ < L >=0.48, < L^2>=0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.84	EDS
Total number of atoms	6513	wwPDB-VP
Average B, all atoms $(Å^2)$	72.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.25% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond	lengths	Bond	angles
	Unam	RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	G	0.35	0/3278	0.48	0/4486
2	Н	0.32	0/1591	0.52	0/2177
3	L	0.60	0/1678	0.58	0/2278
All	All	0.42	0/6547	0.52	0/8941

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	G	3201	0	3070	49	0
2	Н	1551	0	1505	40	0
3	L	1642	0	1602	52	0
4	G	112	0	104	1	0
5	G	6	0	8	0	0
6	G	1	0	0	0	0
All	All	6513	0	6289	137	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.



		Interatomic	Clash
Atom-1 Atom-2		distance (Å)	overlap (Å)
3:L:47:LEU:HD13	3:L:62:PHE:CD2	1.70	1.25
3:L:55:ALA:O	3:L:58:ILE:HG13	1.46	1.15
3:L:47:LEU:CD1	3:L:62:PHE:CE2	2.32	1.13
2:H:11:LEU:HD23	2:H:116:THR:HG22	1.22	1.11
3:L:47:LEU:HD11	3:L:62:PHE:HE2	1.08	1.06
3:L:47:LEU:CD1	3:L:62:PHE:CD2	2.41	1.03
3:L:47:LEU:HD11	3:L:62:PHE:CE2	1.89	1.03
3:L:47:LEU:O	3:L:58:ILE:HD12	1.57	1.03
3:L:47:LEU:HD13	3:L:62:PHE:HD2	1.16	0.99
2:H:38:ARG:HH12	2:H:86:ASP:HA	1.36	0.90
2:H:11:LEU:CD2	2:H:116:THR:HG22	2.04	0.88
3:L:54:ARG:HD2	3:L:62:PHE:O	1.78	0.84
2:H:66:ARG:HD2	2:H:82(B):ASN:CG	2.00	0.80
3:L:55:ALA:HB3	3:L:58:ILE:HD11	1.65	0.79
2:H:11:LEU:HD23	2:H:116:THR:CG2	2.11	0.77
3:L:55:ALA:O	3:L:58:ILE:CG1	2.31	0.77
3:L:47:LEU:HD13	3:L:62:PHE:CE2	2.07	0.75
2:H:66:ARG:HD2	2:H:82(B):ASN:CB	2.18	0.74
3:L:15:PRO:HD3	3:L:107:LYS:HB3	1.68	0.73
3:L:47:LEU:O	3:L:58:ILE:CD1	2.35	0.73
3:L:61:ARG:HH22	3:L:79:GLU:CD	1.93	0.72
3:L:120:PRO:HD3	3:L:132:VAL:HG22	1.71	0.71
3:L:35:TRP:CE3	3:L:73:LEU:HD23	2.26	0.69
2:H:66:ARG:HD2	2:H:82(B):ASN:OD1	1.91	0.68
3:L:35:TRP:CE3	3:L:73:LEU:CD2	2.77	0.68
2:H:18:VAL:HG11	2:H:82:VAL:HG12	1.76	0.68
1:G:199:LYS:HD2	2:H:98:TYR:HB3	1.77	0.66
2:H:100(C):THR:HB	3:L:92:GLY:HA2	1.78	0.66
2:H:66:ARG:NH1	2:H:86:ASP:OD2	2.29	0.66
3:L:61:ARG:NH2	3:L:79:GLU:CD	2.50	0.66
1:G:46:VAL:HG13	1:G:65:PHE:HB2	1.79	0.65
2:H:19:LYS:O	2:H:19:LYS:HD3	1.98	0.64
3:L:61:ARG:NH1	3:L:79:GLU:OE2	2.31	0.63
3:L:144:ALA:HB2	3:L:198:HIS:HD2	1.64	0.63
3:L:47:LEU:HA	3:L:58:ILE:HD13	1.81	0.62
1:G:246:THR:HG23	1:G:263:SER:HB3	1.80	0.62
3:L:183:LYS:NZ	3:L:187:GLU:OE2	2.33	0.61
1:G:341:ALA:O	1:G:387:ARG:NH1	2.33	0.61
3:L:61:ARG:HH22	3:L:79:GLU:CG	2.14	0.61
3:L:125:LEU:HB3	3:L:183:LYS:HE2	1.82	0.61

All (137) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



		Interatomic	Clash
Atom-1	Atom-2	distance $(\text{\AA})$	overlap (Å)
1:G:275:LEU:HB2	1:G:305:PHE:HZ	1.65	0.60
1:G:213:MET:HA	1:G:230:ILE:HA	1.82	0.60
3:L:61:ARG:NH2	3:L:79:GLU:OE2	2.34	0.60
2:H:38:ARG:HG3	2:H:46:GLU:HG3	1.82	0.59
2:H:184:VAL:HG21	2:H:189:LEU:HD21	1.83	0.59
2:H:63:PHE:O	2:H:66:ARG:HG3	2.03	0.59
1:G:297:TYR:HA	4:G:501:NAG:H61	1.85	0.58
3:L:47:LEU:CD1	3:L:62:PHE:HE2	1.78	0.58
1:G:87:ASN:O	1:G:90:ASN:OD1	2.21	0.57
2:H:146:PHE:HB2	2:H:175:LEU:HD23	1.86	0.57
1:G:309:ILE:HD12	1:G:310:PRO:HD2	1.86	0.57
3:L:61:ARG:HH22	3:L:79:GLU:HG2	1.71	0.56
2:H:16:ALA:HB3	2:H:82(B):ASN:H	1.71	0.55
3:L:47:LEU:CD1	3:L:62:PHE:HD2	1.95	0.55
1:G:196:PHE:HB3	1:G:218:ILE:HD12	1.89	0.55
3:L:35:TRP:CZ3	3:L:73:LEU:HD23	2.42	0.55
1:G:175:ARG:HB2	1:G:310:PRO:HG3	1.87	0.54
3:L:47:LEU:HA	3:L:58:ILE:CD1	2.36	0.54
1:G:16:ILE:HD11	1:G:235:TYR:HB3	1.89	0.54
3:L:187:GLU:O	3:L:211:ARG:NH1	2.41	0.54
1:G:136:PHE:HB3	1:G:365:LYS:HD2	1.89	0.54
1:G:366:TRP:HD1	1:G:371:GLY:HA2	1.72	0.54
1:G:225:ASP:H	1:G:246:THR:HA	1.72	0.54
2:H:201:LYS:HB2	2:H:202:PRO:HD3	1.91	0.53
2:H:23:LYS:HA	2:H:77:THR:HG22	1.91	0.53
2:H:119:PRO:HB3	2:H:145:TYR:HB3	1.90	0.53
2:H:155:ASN:HB3	2:H:158:ALA:HB3	1.92	0.52
2:H:18:VAL:O	2:H:18:VAL:HG13	2.09	0.52
1:G:131:TYR:HB3	1:G:139:MET:HE3	1.92	0.52
1:G:132:PHE:HB3	1:G:140:TRP:HB2	1.92	0.52
1:G:176:ALA:O	1:G:269:ARG:NH2	2.43	0.52
2:H:7:SER:O	2:H:20:VAL:HG13	2.11	0.51
2:H:39:GLN:HB2	2:H:45:LEU:HD23	1.93	0.51
1:G:279:SER:HB3	1:G:303:ILE:HG23	1.93	0.50
1:G:87:ASN:N	1:G:87:ASN:ND2	2.59	0.50
1:G:338:SER:HB2	1:G:412:ALA:HA	1.93	0.50
3:L:39:LYS:HG2	3:L:84:ALA:HB2	1.93	0.49
1:G:200:THR:HB	1:G:211:CYS:HB2	1.94	0.49
2:H:51:ILE:HG13	2:H:57:THR:HG22	1.95	0.48
3:L:49:TYR:O	3:L:91:TYR:OH	2.29	0.48
3:L:142:ARG:O	3:L:142:ARG:HG3	2.11	0.48



		Interatomic	Clash
Atom-1	Atom-2	distance $(\text{\AA})$	overlap (Å)
2:H:1:GLN:OE1	2:H:1:GLN:N	2.47	0.48
3:L:61:ARG:NH2	3:L:79:GLU:HG2	2.29	0.48
1:G:344:PRO:HD2	1:G:410:THR:HG21	1.95	0.48
3:L:61:ARG:CZ	3:L:79:GLU:OE2	2.62	0.47
1:G:244:ILE:HG23	1:G:265:ARG:HB3	1.94	0.47
3:L:35:TRP:CD2	3:L:73:LEU:CD2	2.98	0.47
1:G:179:LEU:HB3	1:G:268:PRO:HG2	1.97	0.47
1:G:403:LEU:HB3	1:G:419:VAL:HG23	1.96	0.47
2:H:181:VAL:HG21	3:L:135:LEU:HD22	1.96	0.47
1:G:77:GLN:HE21	1:G:90:ASN:HA	1.79	0.47
1:G:48:VAL:HB	1:G:63:LEU:HB2	1.96	0.47
3:L:182:SER:HB3	3:L:185:ASP:HB2	1.96	0.47
2:H:10:GLU:HB2	2:H:109:VAL:HG23	1.97	0.46
2:H:47:TRP:CZ2	2:H:49:GLY:HA2	2.51	0.46
2:H:11:LEU:HD11	2:H:146:PHE:HE2	1.81	0.46
1:G:210:GLU:OE1	2:H:100(A):ARG:N	2.49	0.45
1:G:275:LEU:HB2	1:G:305:PHE:CZ	2.49	0.45
2:H:38:ARG:NH1	2:H:86:ASP:HA	2.18	0.45
2:H:126:PRO:HB3	2:H:138:LEU:HB3	1.97	0.45
2:H:3:GLN:O	2:H:24:ALA:HA	2.17	0.45
1:G:129:ASP:OD1	1:G:143:HIS:ND1	2.46	0.45
3:L:12:SER:HA	3:L:105:GLU:HB2	1.97	0.45
2:H:89:VAL:HG22	2:H:108:LEU:HG	1.97	0.44
3:L:35:TRP:CD2	3:L:73:LEU:HD22	2.52	0.44
1:G:269:ARG:HB3	1:G:270:PRO:HD3	1.99	0.44
2:H:12:LYS:HD3	2:H:12:LYS:HA	1.66	0.44
1:G:22:ASP:OD2	1:G:108:SER:OG	2.35	0.44
1:G:95:GLU:OE1	1:G:273:ARG:HD2	2.18	0.43
2:H:100(F):LEU:HB2	2:H:103:TRP:CZ3	2.53	0.43
2:H:159:LEU:HD13	2:H:182:VAL:HG11	2.00	0.43
1:G:340:ASP:OD1	1:G:341:ALA:N	2.46	0.43
2:H:66:ARG:HD2	2:H:82(B):ASN:HB3	1.97	0.43
1:G:125:LEU:HD21	1:G:128:VAL:HG23	1.99	0.43
3:L:122:ASP:OD1	3:L:122:ASP:N	2.51	0.43
1:G:23:PRO:HD3	1:G:234:GLY:HA2	2.00	0.43
1:G:26:PHE:HD1	1:G:235:TYR:CZ	2.37	0.42
3:L:198:HIS:ND1	3:L:199:GLN:O	2.52	0.42
3:L:79:GLU:HB2	3:L:80:PRO:HD2	2.01	0.42
3:L:61:ARG:NH2	3:L:79:GLU:CG	2.83	0.42
1:G:352:TRP:HH2	1:G:393:VAL:HG11	1.85	0.42
3:L:80:PRO:HA	3:L:106:ILE:HG13	2.02	0.42



Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:G:333:VAL:HG21	1:G:405:ILE:HG21	2.01	0.41
2:H:124:LEU:HB2	2:H:139:GLY:C	2.41	0.41
3:L:32:TYR:HB3	3:L:91:TYR:O	2.20	0.41
1:G:91:LEU:HA	1:G:107:ARG:HD3	2.02	0.41
1:G:107:ARG:HD2	1:G:274:PHE:CE1	2.55	0.41
3:L:11:LEU:HB3	3:L:13:LEU:HG	2.02	0.41
3:L:43:ALA:HB1	3:L:44:PRO:CD	2.51	0.41
3:L:61:ARG:H	3:L:61:ARG:HG2	1.63	0.41
1:G:408:THR:HB	1:G:414:THR:HG23	2.03	0.41
1:G:350:ALA:HB1	1:G:403:LEU:HD11	2.02	0.40
1:G:13:GLN:HB3	1:G:143:HIS:HB3	2.03	0.40
1:G:198:VAL:HG21	1:G:264:LEU:HD21	2.03	0.40
1:G:213:MET:HB3	1:G:230:ILE:HG22	2.04	0.40
1:G:26:PHE:N	1:G:106:MET:O	2.55	0.40
1:G:93:LEU:HD23	1:G:273:ARG:HD3	2.03	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	G	418/431 (97%)	374 (90%)	44 (10%)	0	100 100
2	Н	204/228~(90%)	191 (94%)	13~(6%)	0	100 100
3	L	212/215~(99%)	198 (93%)	14 (7%)	0	100 100
All	All	834/874~(95%)	763~(92%)	71 (8%)	0	100 100

There are no Ramachandran outliers to report.



#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the side chain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percei	ntiles
1	G	364/373~(98%)	347~(95%)	17 (5%)	26	57
2	Н	174/192~(91%)	165~(95%)	9(5%)	23	54
3	L	184/185~(100%)	177 (96%)	7 (4%)	33	62
All	All	722/750~(96%)	689~(95%)	33 (5%)	27	58

All (33) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	6	LEU
1	G	59	HIS
1	G	87	ASN
1	G	89	THR
1	G	90	ASN
1	G	102	LEU
1	G	110	LYS
1	G	156	THR
1	G	159	TYR
1	G	161	LYS
1	G	164	ASN
1	G	246	THR
1	G	281	LEU
1	G	303	ILE
1	G	364	CYS
1	G	367	THR
1	G	383	PHE
2	Н	19	LYS
2	Н	82(C)	LEU
2	Н	84	SER
2	Н	100(A)	ARG
2	Н	103	TRP
2	Н	110	THR
2	Н	144	ASP
2	Н	175	LEU
2	Н	201	LYS



Continued from previous page...

Mol	Chain	Res	Type
3	L	46	LEU
3	L	47	LEU
3	L	58	ILE
3	L	67	SER
3	L	70	ASP
3	L	73	LEU
3	L	142	ARG

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such side chains are listed below:

Mol	Chain	Res	Type
1	G	77	GLN
1	G	87	ASN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry (i)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mal Tur		Chain	Dec	Tink	Bo	Bond lengths			Bond angles		
INIOI	туре	Unam	nes	LIIIK	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2	
4	NAG	G	507	1	14,14,15	0.23	0	17,19,21	0.53	0	
4	NAG	G	505	1	14,14,15	0.24	0	17,19,21	0.42	0	
4	NAG	G	502	1	14,14,15	0.22	0	$17,\!19,\!21$	0.42	0	
4	NAG	G	503	1	14,14,15	0.23	0	17,19,21	0.40	0	
5	GOL	G	509	-	$5,\!5,\!5$	0.22	0	$5,\!5,\!5$	0.47	0	
4	NAG	G	501	1	14,14,15	0.18	0	17,19,21	0.41	0	
4	NAG	G	504	1	14,14,15	0.23	0	17,19,21	0.40	0	
4	NAG	G	506	1	14,14,15	0.27	0	17,19,21	0.51	0	
4	NAG	G	508	1	14,14,15	0.23	0	17,19,21	0.43	0	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	G	507	1	-	2/6/23/26	0/1/1/1
4	NAG	G	505	1	-	2/6/23/26	0/1/1/1
4	NAG	G	502	1	-	4/6/23/26	0/1/1/1
4	NAG	G	503	1	-	2/6/23/26	0/1/1/1
5	GOL	G	509	-	-	2/4/4/4	-
4	NAG	G	501	1	-	2/6/23/26	0/1/1/1
4	NAG	G	504	1	-	2/6/23/26	0/1/1/1
4	NAG	G	506	1	-	2/6/23/26	0/1/1/1
4	NAG	G	508	1	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (18) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	G	501	NAG	C4-C5-C6-O6
4	G	505	NAG	O5-C5-C6-O6
4	G	501	NAG	O5-C5-C6-O6
4	G	505	NAG	C4-C5-C6-O6
4	G	502	NAG	O5-C5-C6-O6
4	G	502	NAG	C4-C5-C6-O6
4	G	502	NAG	C8-C7-N2-C2



Mol	Chain	Res	Type	Atoms
4	G	502	NAG	O7-C7-N2-C2
4	G	504	NAG	C4-C5-C6-O6
4	G	506	NAG	O5-C5-C6-O6
4	G	507	NAG	O5-C5-C6-O6
4	G	503	NAG	C4-C5-C6-O6
4	G	503	NAG	O5-C5-C6-O6
4	G	504	NAG	O5-C5-C6-O6
4	G	506	NAG	C3-C2-N2-C7
4	G	507	NAG	C3-C2-N2-C7
5	G	509	GOL	O2-C2-C3-O3
5	G	509	GOL	C1-C2-C3-O3

Continued from previous page...

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	G	501	NAG	1	0

## 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ>2		$OWAB(Å^2)$	Q<0.9
1	G	420/431~(97%)	0.38	17 (4%) 38	36	41, 67, 105, 129	0
2	Н	208/228~(91%)	0.54	14 (6%) 17	17	47, 79, 109, 120	0
3	L	214/215~(99%)	0.54	10 (4%) 31	29	30, 75, 96, 107	0
All	All	842/874~(96%)	0.46	41 (4%) 29	27	30, 72, 105, 129	0

All (41) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	159	TYR	5.1
2	Н	137	ALA	4.2
1	G	158	PRO	4.1
1	G	290	PRO	3.5
2	Н	189	LEU	3.3
2	Н	82	VAL	3.1
2	Н	190	GLY	3.1
1	G	155	GLU	3.1
1	G	289	GLY	2.9
1	G	116	THR	2.9
1	G	383	PHE	2.8
1	G	380	SER	2.7
3	L	79	GLU	2.7
1	G	222	LEU	2.7
2	Н	185	PRO	2.7
1	G	163	ASP	2.6
1	G	387	ARG	2.5
2	Н	188	SER	2.4
3	L	19	ALA	2.4
2	Н	138	LEU	2.4
3	L	75	ILE	2.4
2	Н	82(C)	LEU	2.3
3	L	205	VAL	2.3



Mol	Chain	Res	Type	RSRZ
3	L	202	SER	2.3
3	L	129	THR	2.3
1	G	248	THR	2.3
2	Н	203	SER	2.3
2	Н	82(B)	ASN	2.3
1	G	223	PRO	2.3
3	L	80	PRO	2.2
3	L	137	ASN	2.2
3	L	64	GLY	2.2
1	G	191	ALA	2.1
2	Н	43	GLU	2.1
3	L	28	VAL	2.1
1	G	220	GLN	2.1
1	G	341	ALA	2.1
2	Н	194	TYR	2.1
2	Н	135	THR	2.1
2	Н	159	LEU	2.1
1	G	225	ASP	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

#### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(A^2)$	Q<0.9
4	NAG	G	505	14/15	0.74	0.55	$25,\!26,\!27,\!27$	0
4	NAG	G	503	14/15	0.75	0.57	16,17,18,18	0
4	NAG	G	506	14/15	0.78	0.57	26,27,29,30	0
4	NAG	G	502	14/15	0.80	0.44	7,7,8,9	0
4	NAG	G	508	14/15	0.81	0.26	24,24,26,26	0



Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathrm{\AA}^2)$	Q<0.9
4	NAG	G	504	14/15	0.82	0.62	22,23,24,24	0
4	NAG	G	507	14/15	0.83	0.37	19,20,21,24	0
4	NAG	G	501	14/15	0.89	0.23	5, 6, 9, 9	0
5	GOL	G	509	6/6	0.93	0.23	37,37,37,37	0

Continued from previous page...

## 6.5 Other polymers (i)

There are no such residues in this entry.

