

wwPDB EM Validation Summary Report (i)

Oct 14, 2024 – 03:48 AM EDT

PDB ID : 7SSD

EMDB ID : EMD-25407

Title: Mid translocation intermediate with EF-G bound with GDP (Structure IV)

Authors: Carbone, C.E.; Korostelev, A.A.

Deposited on : 2021-11-10

Resolution : 3.30 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*A user guide is available at

https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev113

Mogul : 2022.3.0, CSD as543be (2022)

MolProbity : 4.02b-467 buster-report : 1.1.7 (2018)

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

 $MapQ \quad : \quad 1.9.13$

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


 $\begin{tabular}{lll} Validation Pipeline (wwPDB-VP) & : & 2.39 \end{tabular}$

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $ELECTRON\ MICROSCOPY$

The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	EM structures
Wietric	$(\# ext{Entries})$	$(\# ext{Entries})$
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415
RNA backbone	6643	2191

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion <40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	3	1539	87%	13%
2	1	2903	87%	13%
3	2	120	89%	11%
4	8	697	93%	
5	6	77	78%	22%
6	b	271	97%	
7	С	209	99%	•
8	d	201	98%	•



Continued from previous page...

Mol	Chain	Length	Quality of chain
9	e	177	96%
10	f	176	99%
11	g	149	98%
12	a	234	56% • 43%
13	i	142	98%
14	j	142	100%
15	k	122	98%
16	1	143	98%
17	m	136	
18	n	120	
19	0	116	
20		114	
	р		100%
21	q	117	100%
22	r	103	97%
23	S	110	100%
24	t	93	99%
25	u	102	100%
26	V	94	100%
27	W	75	100%
28	X	77	100%
29	у	63	100%
30	Z	58	98%
31	A	66	70% 30%
32	В	56	100%
33	С	50	100%



Continued from previous page...

Mol	Chain	Length	Quality of chain	
34	D	46	96%	•
35	Е	64	94%	5% •
36	F	38	100%	
37	4	39	36% · · 59%	
38	5	77	78%	22%
39	G	225	89%	8% •
40	Н	206	99%	
41	I	205	99%	
42	J	157	86%	13% •
43	K	100	96%	•••
44	L	151	97%	•
45	M	129	100%	· ·
46	N	127	97%	
47	O	98	<u></u>	•
	P		100%	
48		116	97%	•
49	Q	123	93%	7% •
50	R	114	98%	••
51	S	100	98%	•
52	T	88	100%	
53	U	82	98%	·
54	V	80	95%	5%
55	W	65	95%	5%
56	X	79	96%	
57	Y	85	100%	
58	Z	65	94%	6%



2 Entry composition (i)

There are 59 unique types of molecules in this entry. The entry contains 153049 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a RNA chain called 16S rRNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	3	1539	Total 33012	C 14725	N 6052	O 10697	P 1538	0	0

• Molecule 2 is a RNA chain called 23S rRNA.

Mol	Chain	Residues			Atoms			AltConf	Trace
2	1	2903	Total 62317	C 27801	N 11468	O 20146	P 2902	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
1	747	С	U	$\operatorname{conflict}$	GB 802133627

• Molecule 3 is a RNA chain called 5S RNA.

Mo	ol (Chain	Residues		\mathbf{A}^{1}	toms			AltConf	Trace
3		2	120	Total 2568	C 1145	N 471	O 833	P 119	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
2	120	A	-	insertion	GB 1266961702

• Molecule 4 is a protein called Elongation factor G.

Mol	Chain	Residues		At	oms			AltConf	Trace
4	8	666	Total 5157	C 3256	N 886	O 992	S 23	0	0

• Molecule 5 is a RNA chain called tRNA fMet.



Mol	Chain	Residues		Atoms				AltConf	Trace
5	6	77	Total 1640	C 732	N 297	O 535	P 76	0	0

• Molecule 6 is a protein called 50S ribosomal protein L2.

Mol	Chain	Residues		At	oms			AltConf	Trace
6	b	271	Total 2083	C 1288	N 423	O 365	S 7	0	0

• Molecule 7 is a protein called 50S ribosomal protein L3.

Mol	Chain	Residues		At	oms			AltConf	Trace
7	c	209	Total	С	N	0	S	0	0
			1565	979	288	294	4		

• Molecule 8 is a protein called 50S ribosomal protein L4.

Mol	Chain	Residues		At	oms	AltConf	Trace		
0	J	201	Total	С	N	О	S	0	0
0	a	201	1552	974	283	290	5	U	U

• Molecule 9 is a protein called 50S ribosomal protein L5.

Mol	Chain	Residues		At	oms	AltConf	Trace		
9	e	177	Total 1411	C 899	N 249	O 257	S 6	0	0

• Molecule 10 is a protein called 50S ribosomal protein L6.

Mol	Chain	Residues		At	oms	AltConf	Trace		
10	t	176	Total	С	N	О	S	0	0
10	1	170	1323	832	243	246	2	0	U

• Molecule 11 is a protein called 50S ribosomal protein L9.

Mol	Chain	Residues		At	oms	AltConf	Trace		
11	g	149	Total	C 699	N 197	O 214	S 1	0	0

• Molecule 12 is a protein called 50S ribosomal protein L1.



Mol	Chain	Residues		At	oms	AltConf	Trace		
19		134	Total	С	N	О	S	0	0
12	a	104	1026	645	186	193	2	0	U

• Molecule 13 is a protein called 50S ribosomal protein L11.

Mol	Chain	Residues		At	oms	AltConf	Trace		
12	i	141	Total	С	N	О	S	0	0
10		141	1032	651	179	196	6		U

• Molecule 14 is a protein called 50S ribosomal protein L13.

Mol	Chain	Residues		At	oms	AltConf	Trace		
1./	i	142	Total	С	N	О	S	0	0
14	J	142	1129	714	212	199	4		0

• Molecule 15 is a protein called 50S ribosomal protein L14.

Mol	Chain	Residues		At	oms	AltConf	Trace		
15	1,	122	Total	С	N	О	S	0	0
10	K	122	939	587	180	166	6	0	U

• Molecule 16 is a protein called 50S ribosomal protein L15.

Mol	Chain	Residues		At	oms	AltConf	Trace		
16	1	143	Total 1045	C 649	N 206	O 189	S 1	0	0

• Molecule 17 is a protein called 50S ribosomal protein L16.

Mol	Chain	Residues		At	oms	AltConf	Trace		
17	m	136	Total	С	N	О	S	0	0
11	111	100	1074	686	205	177	6		

• Molecule 18 is a protein called 50S ribosomal protein L17.

Mol	Chain	Residues		At	oms	AltConf	Trace		
18	n	120	Total 961	C 593	N 196	O 167	S 5	0	0

• Molecule 19 is a protein called 50S ribosomal protein L18.



\mathbf{Mol}	Chain	Residues		Ato	ms		AltConf	Trace
19	О	116	Total 892	C 552	N 178	O 162	0	0

• Molecule 20 is a protein called 50S ribosomal protein L19.

Mol	Chain	Residues		At	oms	AltConf	Trace		
20		11/	Total	С	N	О	S	0	0
20	p	114	917	574	179	163	1	0	U

• Molecule 21 is a protein called 50S ribosomal protein L20.

Mol	Chain	Residues		Ato	ms	AltConf	Trace	
21	q	117	Total 947	C 604	N 192	O 151	0	0

• Molecule 22 is a protein called 50S ribosomal protein L21.

Mol	Chain	Residues		At	oms	AltConf	Trace		
22	r	103	Total	С	N	О	S	0	0
22	1	105	816	516	153	145	2	0	

• Molecule 23 is a protein called 50S ribosomal protein L22.

Mol	Chain	Residues		At	oms	AltConf	Trace		
23	s	110	Total 857	C 532	N 166	O 156	S 3	0	0

• Molecule 24 is a protein called 50S ribosomal protein L23.

Mol	Chain	Residues		At	oms	AltConf	Trace		
24	t	93	Total	С	N	0	S	0	0
			739	466	139	132	2		_

• Molecule 25 is a protein called 50S ribosomal protein L24.

Mol	Chain	Residues		Ato	ms	AltConf	Trace	
25	u	102	Total 780	C 492	N 146	O 142	0	0

• Molecule 26 is a protein called 50S ribosomal protein L25.



Mol	Chain	Residues		At	oms	AltConf	Trace		
26	**	0.4	Total	С	N	О	S	0	0
20	V	94	753	479	137	134	3	U	U

• Molecule 27 is a protein called 50S ribosomal protein L27.

Mol	Chain	Residues		At	AltConf	Trace			
27	W	75	Total	C 256	N 116	0	S	0	0
21	W	(9	575	356	116	102	1	U	

• Molecule 28 is a protein called 50S ribosomal protein L28.

Mol	Chain	Residues		At	oms	AltConf	Trace		
28	X	77	Total 625	C 388	N 129	O 106	S 2	0	0

• Molecule 29 is a protein called 50S ribosomal protein L29.

Mol	Chain	Residues		Ato	ms	AltConf	Trace		
29	**	62	Total	С	N	О	S	0	0
29	У	63	509	313	99	95	2	0	U

• Molecule 30 is a protein called 50S ribosomal protein L30.

Mol	Chain	Residues		Atoms					Trace
30	7.	58	Total	С	N	О	S	0	0
	2	90	449	281	87	79	2		U

• Molecule 31 is a protein called 50S ribosomal protein L31.

Mol	Chain	Residues		Ato	$\mathbf{m}\mathbf{s}$			AltConf	Trace
91	Λ	46	Total	С	N	О	S	0	0
31	A	40	355	221	62	66	6	0	U

• Molecule 32 is a protein called 50S ribosomal protein L32.

Mol	Chain	Residues		Ato	ms			AltConf	Trace
32	В	56	Total 444	C 269	N 94	O 80	S 1	0	0

• Molecule 33 is a protein called 50S ribosomal protein L33.



Mol	Chain	Residues		Aton	ns	AltConf	Trace	
33	С	50	Total 410	C 263	N 75	O 72	0	0

• Molecule 34 is a protein called 50S ribosomal protein L34.

Mol	Chain	Residues		Ato	$\overline{\mathrm{ms}}$			AltConf	Trace
34	D	46	Total	С	N	О	S	0	0
94	D	40	377	228	90	57	2		U

• Molecule 35 is a protein called 50S ribosomal protein L35.

Mol	Chain	Residues		\mathbf{At}	oms			AltConf	Trace
35	Е	64	Total 504	C 323	1	O 74	S 2	0	0

• Molecule 36 is a protein called 50S ribosomal protein L36.

Mol	Chain	Residues		Ato	ms			AltConf	Trace
36	E	20	Total	С	N	О	S	0	0
30	Г	38	302	185	65	48	4	0	U

• Molecule 37 is a RNA chain called mRNA.

Mol	Chain	Residues		At	oms	AltConf	Trace		
37	4	16	Total 350	C 156	N 70	O 108	P 16	0	0

• Molecule 38 is a RNA chain called tRNA Pro.

Mol	Chain	Residues		\mathbf{A}	toms	AltConf	Trace		
38	5	77	Total 1647	C 733	N 295	O 542	P 77	0	0

• Molecule 39 is a protein called 30S ribosomal protein S2.

Mol	Chain	Residues		Ato	AltConf	Trace			
39	G	218	Total 1705	C 1081	N 305	O 312	S 7	0	0

• Molecule 40 is a protein called 30S ribosomal protein S3.



Mol	Chain	Residues		Ato	oms			AltConf	Trace
40	Н	206	Total 1625	C 1028	N 305	O 289	S 3	0	0

• Molecule 41 is a protein called 30S ribosomal protein S4.

Mol	Chain	Residues		Ato	oms			AltConf	Trace
41	I	205	Total 1643	C 1026	N 315	O 298	S 4	0	0

• Molecule 42 is a protein called 30S ribosomal protein S5.

Mol	Chain	Residues		At	oms			AltConf	Trace
49	т	157	Total	С	N	О	S	0	0
42	J	157	1157	719	218	214	6	0	U

• Molecule 43 is a protein called 30S ribosomal protein S6.

Mol	Chain	Residues		At	oms			AltConf	Trace
49	I/	100	Total	С	N	О	S	0	0
45	K	100	818	515	148	149	6	0	0

• Molecule 44 is a protein called 30S ribosomal protein S7.

Mol	Chain	Residues		At	oms	AltConf	Trace		
44	L	151	Total 1182	C 735	N 227	O 216	S 4	0	0

• Molecule 45 is a protein called 30S ribosomal protein S8.

Mol	Chain	Residues		At	oms			AltConf	Trace
45	М	129	Total	С	N	О	S	0	0
45	M	129	979	616	173	184	6	0	U

• Molecule 46 is a protein called 30S ribosomal protein S9.

Mol	Chain	Residues		At	oms			AltConf	Trace
46	N	127	Total 1022	C 634	N 206	O 179	S 3	0	0

• Molecule 47 is a protein called 30S ribosomal protein S10.



Mol	Chain	Residues		At	oms	AltConf	Trace		
47	О	98	Total	C	N 150	0	S	0	0
			787	493	150	143	1		

• Molecule 48 is a protein called 30S ribosomal protein S11.

Mol	Chain	Residues		At	oms			AltConf	Trace
48	D	116	Total	С	N	О	S	0	0
40	1	110	870	535	173	159	3	0	U

• Molecule 49 is a protein called 30S ribosomal protein S12.

Mol	Chain	Residues		At	oms			AltConf	Trace
49	Q	123	Total 955	C 590	N 196	O 165	S	0	0

• Molecule 50 is a protein called 30S ribosomal protein S13.

Mol	Chain	Residues		At	oms			AltConf	Trace
50	D	11/	Total	С	N	О	S	0	0
30	n	114	884	546	178	157	3	U	U

• Molecule 51 is a protein called 30S ribosomal protein S14.

Mol	Chain	Residues		At	oms			AltConf	Trace
51	S	100	Total 805	C 499	N 164	O 139	S 3	0	0

• Molecule 52 is a protein called 30S ribosomal protein S15.

Mol	Chain	Residues		At	oms			AltConf	Trace
59	Т	88	Total	С	N	О	S	0	0
32	1	00	714	439	144	130	1	0	U

• Molecule 53 is a protein called 30S ribosomal protein S16.

Mol	Chain	Residues		At	oms			AltConf	Trace
53	II	99	Total	С	N	О	S	0	0
55	U	02	649	406	128	114	1		U

• Molecule 54 is a protein called 30S ribosomal protein S17.



\mathbf{Mol}	Chain	Residues		At	oms			AltConf	Trace
54	V	80	Total	С	N	О	S	0	0
54	V	0 0	649	411	121	114	3	U	U

• Molecule 55 is a protein called 30S ribosomal protein S18.

Mol	Chain	Residues	Atoms			AltConf	Trace		
55	7.7.7	65	Total	С	N	О	S	0	0
00	VV	0.5	536	339	100	96	1	U	U

• Molecule 56 is a protein called 30S ribosomal protein S19.

Mol	Chain	Residues	Atoms			AltConf	Trace		
56	X	79	Total 638	C 408	N 120	O 108	S 2	0	0

• Molecule 57 is a protein called 30S ribosomal protein S20.


Mol	Chain	Residues	Atoms			AltConf	Trace		
57	V	05	Total	С	N	О	S	0	0
37	1	85	665	411	137	114	3	0	U

• Molecule 58 is a protein called 30S ribosomal protein S21.

Mol	Chain	Residues	Atoms			AltConf	Trace		
58	Z	65	Total 545	C 335	N 117	O 92	S 1	0	0

• Molecule 59 is GUANOSINE-5'-DIPHOSPHATE (three-letter code: GDP) (formula: $C_{10}H_{15}N_5O_{11}P_2$) (labeled as "Ligand of Interest" by depositor).






Mol	Chain	Residues	Atoms			AltConf		
50	0	1	Total	С	N	О	Р	0
39	0	1	28	10	5	11	2	U




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: 16S rRNA








• Molecule 3: 5S RNA

Chain 2: 89% 11% • Molecule 4: Elongation factor G

Chain 8: 93%





22%



Chain 6:

• Molecule 5: tRNA fMet

• Molecule 6: 50S ribosomal protein L2

Chain b: 97%

78%

• Molecule 7: 50S ribosomal protein L3

Chain c: 99% .

M1 T133 H134 A209

• Molecule 8: 50S ribosomal protein L4

Chain d: 98% .

M1 R61 R79 G82 G82 V146

• Molecule 9: 50S ribosomal protein L5

Chain e: 96% ...

N20 N20 R29 K114 K114 L151 L151 L151 R177

• Molecule 10: 50S ribosomal protein L6


Chain f: 99%

A45 V131 A173 K176

• Molecule 11: 50S ribosomal protein L9

Chain g: 98%





• Molecule 18: 50S ribosomal protein L17



Chain n:	99%	•
M1 E120		
• Molecule 19: 50S	ribosomal protein L18	
Chain o:	99%	
753 F117		
• Molecule 20: 50S	ribosomal protein L19	
Chain p:	100%	
There are no outlier	residues recorded for this chain.	
• Molecule 21: 50S	ribosomal protein L20	
Chain q:	100%	
There are no outlier	residues recorded for this chain.	
• Molecule 22: 50S	ribosomal protein L21	
Chain r:	97%	
M1		
• Molecule 23: 50S	ribosomal protein L22	
Chain s:	100%	
There are no outlier	residues recorded for this chain.	
• Molecule 24: 50S	ribosomal protein L23	
Chain t:	99%	
183 H		
• Molecule 25: 50S	ribosomal protein L24	
Chain u:	100%	
There are no outlier	residues recorded for this chain.	

 \bullet Molecule 26: 50S ribosomal protein L25



Chain v:	100%	
There are no outlier residues recorded	for this chain.	
• Molecule 27: 50S ribosomal protein I	227	
Chain w:	100%	
There are no outlier residues recorded	for this chain.	
• Molecule 28: 50S ribosomal protein I	228	
Chain x:	100%	
There are no outlier residues recorded	for this chain.	
• Molecule 29: 50S ribosomal protein I	L29	
Chain y:	100%	
There are no outlier residues recorded	for this chain.	
• Molecule 30: 50S ribosomal protein I	1 30	
Chain z:	98%	•
A11		
• Molecule 31: 50S ribosomal protein I	. 31	
Chain A: 70%		30%
M1 C46 C1/S C1/S C1/S C1/S C1/S C1/S C1/S C1/S		
• Molecule 32: 50S ribosomal protein I	. 32	
Chain B:	100%	
There are no outlier residues recorded	for this chain.	
• Molecule 33: 50S ribosomal protein I	- 33	
Chain C:	100%	
There are no outlier residues recorded	for this chain.	
• Molecule 34: 50S ribosomal protein I	. 34	
Chain D:	96%	•





• Molecule 35: 50S ribosomal protein L35

Chain E: 94% 5% •

P1 L28 R29 H30 I31 L32


• Molecule 36: 50S ribosomal protein L36

Chain F:




• Molecule 37: mRNA

Chain 4: 36% . . 59%



• Molecule 38: tRNA Pro

Chain 5: 78% 22%



• Molecule 39: 30S ribosomal protein S2

Chain G: 89% 8% .



• Molecule 40: 30S ribosomal protein S3


Chain H: 99%



• Molecule 41: 30S ribosomal protein S4

Chain I:





• Molecule 42: 30S ribosomal protein S5

Chain J: 86% 13% •

E9 0111 0111 0111 V20 F47 G50 G60 G60 G60 W77 W88 W98 W98 W98 W98 W99 W122 W122 W122 W122 W122 W122 W146 W151 W151 W151 W151 W151

• Molecule 43: 30S ribosomal protein S6

Chain K: 96% ...

M1 151 N52 K53 L54

• Molecule 44: 30S ribosomal protein S7

Chain L:



• Molecule 45: 30S ribosomal protein S8

Chain M:

There are no outlier residues recorded for this chain.

• Molecule 46: 30S ribosomal protein S9

Chain N: 97%



• Molecule 47: 30S ribosomal protein S10

Chain O:




• Molecule 48: 30S ribosomal protein S11

Chain P: 97%



Chain W:





95%

5%



• Molecule 56: 30S ribosomal protein S19

Chain X: 96% ...



• Molecule 57: 30S ribosomal protein S20

Chain Y: 100%

There are no outlier residues recorded for this chain.

• Molecule 58: 30S ribosomal protein S21

Chain Z: 94% 6%





4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	3778	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE	Depositor
	CORRECTION	
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{Å}^2)$	47.6	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	27.032	Depositor
Minimum map value	-14.235	Depositor
Average map value	0.111	Depositor
Map value standard deviation	0.944	Depositor
Recommended contour level	0.7	Depositor
Map size (Å)	370.5408, 370.5408, 370.5408	wwPDB
Map dimensions	448, 448, 448	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.8271, 0.8271, 0.8271	Depositor



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: GDP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

N / L - 1	Clara in	Bond	lengths	В	Sond angles
Mol	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	3	0.33	0/36963	0.66	0/57662
2	1	0.36	0/69796	0.66	0/108888
3	2	0.33	0/2872	0.66	0/4479
4	8	0.34	0/5253	0.65	3/7106 (0.0%)
5	6	0.35	0/1832	0.66	0/2855
6	b	0.52	0/2122	0.70	$1/2852 \ (0.0\%)$
7	С	0.44	0/1586	0.66	0/2134
8	d	0.38	0/1571	0.63	0/2113
9	е	0.36	0/1435	0.67	2/1926 (0.1%)
10	f	0.33	0/1343	0.58	0/1816
11	g	0.33	0/1122	0.66	0/1515
12	a	0.30	0/1033	0.58	0/1387
13	i	0.31	0/1046	0.63	0/1410
14	j	0.43	0/1152	0.59	0/1551
15	k	0.45	0/948	0.75	1/1268 (0.1%)
16	1	0.40	0/1054	0.70	0/1403
17	m	0.46	0/1093	0.64	0/1460
18	n	0.44	0/974	0.71	1/1301 (0.1%)
19	О	0.34	0/902	0.56	0/1209
20	p	0.42	0/929	0.59	0/1242
21	q	0.46	0/960	0.53	0/1278
22	r	0.39	0/829	0.63	0/1107
23	s	0.40	0/864	0.60	0/1156
24	t	0.39	0/745	0.64	1/994 (0.1%)
25	u	0.38	0/788	0.58	0/1051
26	V	0.37	0/766	0.57	0/1025
27	W	0.41	0/582	0.61	0/769
28	X	0.45	0/635	0.62	0/848
29	у	0.32	0/510	0.58	0/677
30	Z	0.34	0/453	0.67	0/605
31	A	0.31	0/362	0.61	0/485
32	В	0.42	0/450	0.64	0/599



3.7.1	<i>α</i> ι :	Bond	lengths	Е	Bond angles
Mol	Chain	RMSZ	# Z >5	RMSZ	# Z >5
33	С	0.39	0/417	0.59	0/554
34	D	0.46	0/380	0.78	1/498~(0.2%)
35	Е	0.42	0/513	0.83	2/676~(0.3%)
36	F	0.45	0/303	0.73	0/397
37	4	0.66	0/392	0.73	0/608
38	5	0.38	0/1840	0.69	0/2868
39	G	0.35	0/1736	0.54	0/2338
40	Н	0.33	0/1652	0.61	0/2225
41	I	0.32	0/1665	0.64	$1/2227 \ (0.0\%)$
42	J	0.32	0/1170	0.55	0/1573
43	K	0.39	0/836	0.73	2/1128 (0.2%)
44	L	0.29	0/1196	0.58	0/1602
45	M	0.35	0/989	0.59	0/1326
46	N	0.34	0/1034	0.70	1/1375~(0.1%)
47	O	0.32	0/797	0.71	0/1077
48	Р	0.35	0/886	0.64	1/1195 (0.1%)
49	Q	0.35	0/969	0.71	0/1300
50	R	0.31	0/893	0.62	0/1193
51	S	0.31	0/817	0.56	0/1088
52	Τ	0.35	0/722	0.61	0/964
53	U	0.32	0/659	0.68	0/884
54	V	0.38	0/658	0.68	1/881 (0.1%)
55	W	0.34	0/545	0.63	1/731 (0.1%)
56	X	0.34	0/653	0.68	1/877 (0.1%)
57	Y	0.30	0/671	0.57	0/888
58	Z	0.36	0/551	0.71	0/728
All	All	0.36	0/165914	0.66	$20/247372 \ (0.0\%)$

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	1	0	3
4	8	0	5
7	c	0	1
8	d	0	2
9	е	0	1
10	f	0	1
11	g	0	1
16	1	0	1



Continued from previous page...

Mol	Chain	#Chirality outliers	#Planarity outliers
17	m	0	2
22	r	0	1
30	Z	0	1
35	Е	0	1
37	4	0	1
41	I	0	1
43	K	0	1
44	L	0	1
46	N	0	1
49	Q	0	4
50	R	0	2
54	V	0	2
55	W	0	1
56	X	0	1
58	Z	0	3
All	All	0	38

There are no bond length outliers.

The worst 5 of 20 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$Ideal(^{o})$
4	8	195	ASP	CB-CG-OD1	9.24	126.62	118.30
41	I	4	LEU	CA-CB-CG	6.86	131.07	115.30
35	Е	28	LEU	CA-CB-CG	6.63	130.56	115.30
35	Е	32	LEU	CA-CB-CG	6.50	130.25	115.30
18	n	38	LEU	CA-CB-CG	6.27	129.72	115.30

There are no chirality outliers.

5 of 38 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	1	1779	U	Sidechain
2	1	1964	G	Sidechain
2	1	511	U	Sidechain
4	8	193	TRP	Peptide
4	8	368	ALA	Peptide

5.2 Too-close contacts (i)

Due to software issues we are unable to calculate clashes - this section is therefore empty.



5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
4	8	662/697~(95%)	595 (90%)	65 (10%)	2 (0%)	37	66
6	b	269/271 (99%)	234 (87%)	33 (12%)	2 (1%)	19	50
7	c	207/209 (99%)	190 (92%)	16 (8%)	1 (0%)	25	56
8	d	199/201 (99%)	173 (87%)	25 (13%)	1 (0%)	25	56
9	e	175/177 (99%)	152 (87%)	23 (13%)	0	100	100
10	f	174/176 (99%)	152 (87%)	22 (13%)	0	100	100
11	g	147/149 (99%)	128 (87%)	19 (13%)	0	100	100
12	a	130/234 (56%)	118 (91%)	12 (9%)	0	100	100
13	i	139/142 (98%)	115 (83%)	24 (17%)	0	100	100
14	j	140/142 (99%)	128 (91%)	12 (9%)	0	100	100
15	k	120/122 (98%)	106 (88%)	13 (11%)	1 (1%)	16	46
16	1	141/143 (99%)	120 (85%)	20 (14%)	1 (1%)	19	50
17	m	134/136 (98%)	114 (85%)	18 (13%)	2 (2%)	8	33
18	n	118/120 (98%)	102 (86%)	16 (14%)	0	100	100
19	О	114/116 (98%)	111 (97%)	3 (3%)	0	100	100
20	р	112/114 (98%)	97 (87%)	15 (13%)	0	100	100
21	q	115/117 (98%)	113 (98%)	2 (2%)	0	100	100
22	r	101/103 (98%)	88 (87%)	13 (13%)	0	100	100
23	S	108/110 (98%)	102 (94%)	6 (6%)	0	100	100
24	t	91/93 (98%)	80 (88%)	11 (12%)	0	100	100
25	u	100/102 (98%)	89 (89%)	11 (11%)	0	100	100
26	v	92/94 (98%)	81 (88%)	11 (12%)	0	100	100
27	w	73/75 (97%)	64 (88%)	9 (12%)	0	100	100
28	X	75/77 (97%)	67 (89%)	8 (11%)	0	100	100
29	у	61/63 (97%)	60 (98%)	1 (2%)	0	100	100



Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
30	Z	56/58~(97%)	49 (88%)	7 (12%)	0	100	100
31	A	44/66~(67%)	35 (80%)	9 (20%)	0	100	100
32	В	54/56~(96%)	49 (91%)	5 (9%)	0	100	100
33	С	48/50 (96%)	47 (98%)	1 (2%)	0	100	100
34	D	44/46 (96%)	38 (86%)	6 (14%)	0	100	100
35	Е	62/64 (97%)	51 (82%)	9 (14%)	2 (3%)	3	20
36	F	36/38 (95%)	30 (83%)	6 (17%)	0	100	100
39	G	216/225 (96%)	179 (83%)	28 (13%)	9 (4%)	2	15
40	Н	204/206 (99%)	193 (95%)	11 (5%)	0	100	100
41	I	203/205 (99%)	180 (89%)	22 (11%)	1 (0%)	25	56
42	J	155/157 (99%)	108 (70%)	32 (21%)	15 (10%)	0	3
43	K	98/100 (98%)	79 (81%)	18 (18%)	1 (1%)	13	42
44	L	149/151 (99%)	132 (89%)	17 (11%)	0	100	100
45	M	127/129 (98%)	114 (90%)	13 (10%)	0	100	100
46	N	125/127 (98%)	107 (86%)	18 (14%)	0	100	100
47	О	96/98 (98%)	78 (81%)	18 (19%)	0	100	100
48	Р	114/116 (98%)	99 (87%)	14 (12%)	1 (1%)	14	44
49	Q	121/123 (98%)	94 (78%)	23 (19%)	4 (3%)	3	19
50	R	112/114 (98%)	101 (90%)	11 (10%)	0	100	100
51	S	98/100 (98%)	91 (93%)	7 (7%)	0	100	100
52	Т	86/88 (98%)	79 (92%)	7 (8%)	0	100	100
53	U	80/82 (98%)	64 (80%)	15 (19%)	1 (1%)	10	36
54	V	78/80 (98%)	68 (87%)	10 (13%)	0	100	100
55	W	63/65 (97%)	59 (94%)	4 (6%)	0	100	100
56	X	77/79 (98%)	64 (83%)	13 (17%)	0	100	100
57	Y	83/85 (98%)	77 (93%)	6 (7%)	0	100	100
58	Z	63/65 (97%)	48 (76%)	14 (22%)	1 (2%)	8	32
All	All	6489/6756 (96%)	5692 (88%)	752 (12%)	45 (1%)	21	50

5 of 45 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
35	Ε	31	ILE



Continued from previous page...

Mol	Chain	Res	Type
39	G	18	GLN
39	G	34	ARG
42	J	87	VAL
42	J	160	VAL

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
4	8	548/574~(96%)	542 (99%)	6 (1%)	70	82
6	b	$216/216\ (100\%)$	212 (98%)	4 (2%)	52	72
7	c	164/164 (100%)	164 (100%)	0	100	100
8	d	165/165~(100%)	164 (99%)	1 (1%)	84	90
9	е	148/148 (100%)	143 (97%)	5 (3%)	32	59
10	f	137/137 (100%)	136 (99%)	1 (1%)	81	88
11	g	114/114 (100%)	112 (98%)	2 (2%)	54	74
12	a	110/181 (61%)	107 (97%)	3 (3%)	40	65
13	i	109/110 (99%)	107 (98%)	2 (2%)	54	74
14	j	116/116 (100%)	116 (100%)	0	100	100
15	k	103/103 (100%)	103 (100%)	0	100	100
16	1	102/102 (100%)	101 (99%)	1 (1%)	73	84
17	m	109/109 (100%)	107 (98%)	2 (2%)	54	74
18	n	100/100 (100%)	99 (99%)	1 (1%)	73	84
19	О	86/86 (100%)	85 (99%)	1 (1%)	67	80
20	р	99/99 (100%)	99 (100%)	0	100	100
21	q	89/89 (100%)	89 (100%)	0	100	100
22	r	84/84 (100%)	82 (98%)	2 (2%)	44	68
23	S	93/93 (100%)	93 (100%)	0	100	100
24	t	80/80 (100%)	80 (100%)	0	100	100



 $Continued\ from\ previous\ page...$

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
25	u	83/83 (100%)	83 (100%)	0	100	100
26	v	78/78~(100%)	78 (100%)	0	100	100
27	W	57/57~(100%)	57 (100%)	0	100	100
28	X	$67/67 \; (100\%)$	67 (100%)	0	100	100
29	у	55/55~(100%)	55 (100%)	0	100	100
30	Z	48/48 (100%)	48 (100%)	0	100	100
31	A	42/59~(71%)	42 (100%)	0	100	100
32	В	$47/47 \; (100\%)$	47 (100%)	0	100	100
33	С	45/45 (100%)	45 (100%)	0	100	100
34	D	38/38 (100%)	37 (97%)	1 (3%)	41	66
35	Е	51/51 (100%)	51 (100%)	0	100	100
36	F	34/34~(100%)	34 (100%)	0	100	100
39	G	180/186~(97%)	171 (95%)	9 (5%)	20	48
40	Н	170/170 (100%)	168 (99%)	2 (1%)	67	80
41	I	172/172 (100%)	171 (99%)	1 (1%)	84	90
42	J	119/119 (100%)	111 (93%)	8 (7%)	13	39
43	K	87/87 (100%)	86 (99%)	1 (1%)	70	82
44	L	124/124~(100%)	121 (98%)	3 (2%)	44	68
45	M	104/104~(100%)	104 (100%)	0	100	100
46	N	105/105~(100%)	103 (98%)	2 (2%)	52	72
47	O	86/86~(100%)	86 (100%)	0	100	100
48	Р	89/89 (100%)	87 (98%)	2 (2%)	47	69
49	Q	103/103 (100%)	101 (98%)	2 (2%)	52	72
50	R	$92/92\ (100\%)$	91 (99%)	1 (1%)	70	82
51	S	83/83 (100%)	81 (98%)	2 (2%)	44	68
52	Т	76/76 (100%)	76 (100%)	0	100	100
53	U	$65/65 \; (100\%)$	64 (98%)	1 (2%)	60	77
54	V	74/74 (100%)	73 (99%)	1 (1%)	62	78
55	W	56/56 (100%)	55 (98%)	1 (2%)	54	74
56	X	70/70 (100%)	68 (97%)	2 (3%)	37	63
57	Y	$65/65 \; (100\%)$	65 (100%)	Continued of	100	100



Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
58	Z	55/55~(100%)	55 (100%)	0	100	100
All	All	5392/5513 (98%)	5322 (99%)	70 (1%)	64	79

5 of 70 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
46	N	65	THR
48	Р	127	ARG
53	U	8	ARG
16	1	94	THR
13	i	81	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 84 such sidechains are listed below:

Mol	Chain	Res	Type
41	I	115	GLN
47	О	64	GLN
42	J	69	ASN
43	K	81	ASN
52	Т	36	ASN

5.3.3 RNA (i)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	3	1538/1539 (99%)	201 (13%)	2 (0%)
2	1	2902/2903 (99%)	383 (13%)	9 (0%)
3	2	119/120 (99%)	12 (10%)	1 (0%)
37	4	14/39 (35%)	2 (14%)	0
38	5	76/77 (98%)	17 (22%)	0
5	6	76/77~(98%)	17 (22%)	0
All	All	4725/4755 (99%)	632 (13%)	12 (0%)

5 of 632 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	3	5	U
1	3	9	G
1	3	22	G
1	3	30	U
1	3	31	G



5 of 12 RNA pucker outliers are listed below:

Mol	Chain	Res	Type
2	1	1930	G
2	1	2326	С
3	2	88	С
2	1	2391	G
2	1	784	G

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Γ.	Mol	Type	Chain Res Link		Link	Bond lengths			Bond angles		
-	MOI	Туре	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
	59	GDP	8	801	-	25,30,30	1.01	1 (4%)	30,47,47	1.14	2 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

\mathbf{Mol}	Type	Chain	Res	Link	Chirals	Torsions	Rings
59	GDP	8	801	ı	-	4/12/32/32	0/3/3/3

All (1) bond length outliers are listed below:



Mol	Chain	Res	Type	Atoms	\mathbf{Z}	Observed(A)	$\operatorname{Ideal}(ext{\AA})$
59	8	801	GDP	C6-N1	-2.71	1.33	1.37

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
59	8	801	GDP	C8-N7-C5	2.78	107.28	102.55
59	8	801	GDP	C5-C6-N1	2.09	118.05	114.07

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
59	8	801	GDP	C5'-O5'-PA-O1A
59	8	801	GDP	C5'-O5'-PA-O3A
59	8	801	GDP	C5'-O5'-PA-O2A
59	8	801	GDP	O4'-C4'-C5'-O5'

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.




6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-25407. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections (i)

6.1.1 Primary map



6.1.2 Raw map





The images above show the map projected in three orthogonal directions.




6.2 Central slices (i)

6.2.1 Primary map









X Index: 224


Y Index: 224

Z Index: 224

6.2.2 Raw map







X Index: 224

Y Index: 224


Z Index: 224

The images above show central slices of the map in three orthogonal directions.




6.3 Largest variance slices (i)

6.3.1 Primary map



6.3.2 Raw map




The images above show the largest variance slices of the map in three orthogonal directions.




6.4 Orthogonal standard-deviation projections (False-color) (i)

6.4.1 Primary map



6.4.2 Raw map




The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.




6.5 Orthogonal surface views (i)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.7. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.

6.6 Mask visualisation (i)


This section was not generated. No masks/segmentation were deposited.



7 Map analysis (i)

This section contains the results of statistical analysis of the map.


7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



7.2 Volume estimate (i)




The volume at the recommended contour level is $2300~\mathrm{nm}^3$; this corresponds to an approximate mass of $2078~\mathrm{kDa}$.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



7.3 Rotationally averaged power spectrum (i)




*Reported resolution corresponds to spatial frequency of 0.303 $\rm \mathring{A}^{-1}$



8 Fourier-Shell correlation (i)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC (i)



*Reported resolution corresponds to spatial frequency of 0.303 $\rm \mathring{A}^{-1}$



8.2 Resolution estimates (i)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
rtesolution estimate (A)	0.143	0.5	Half-bit
Reported by author	3.30	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	4.21	6.76	4.38


^{*}Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 4.21 differs from the reported value 3.3 by more than 10 %



9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-25407 and PDB model 7SSD. Per-residue inclusion information can be found in section 3 on page 15.


9.1 Map-model overlay (i)



The images above show the 3D surface view of the map at the recommended contour level 0.7 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.




9.2 Q-score mapped to coordinate model (i)



The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


9.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.7).



9.4 Atom inclusion (i)



At the recommended contour level, 96% of all backbone atoms, 95% of all non-hydrogen atoms, are inside the map.



9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (0.7) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.9460	0.3850
1	0.9740	0.4280
2	0.9770	0.3950
3	0.9420	0.3230
4	0.8170	0.1920
5	0.9450	0.3160
6	0.9450	0.3080
8	0.8230	0.2730
A	0.9260	0.3310
В	0.9740	0.4740
С	0.8860	0.3930
D	0.9580	0.5160
Е	0.9720	0.5020
F	0.9450	0.4570
G	0.8510	0.2790
Н	0.8920	0.3170
I	0.9040	0.2880
J	0.9370	0.3710
K	0.9490	0.3490
L	0.8520	0.2810
M	0.9500	0.4130
N	0.8800	0.3240
O	0.8400	0.2630
P	0.9490	0.3980
Q	0.9030	0.3800
R	0.8620	0.2820
S	0.9100	0.3160
Т	0.9540	0.3960
U	0.9250	0.3660
V	0.9380	0.3650
W	0.9610	0.4030
X	0.8670	0.2880
Y	0.9420	0.3570
Z	0.9130	0.3540
a	0.8540	0.2550



Continued on next page...



$Continued\ from\ previous\ page...$

Chain	Atom inclusion	Q-score
b	0.9670	0.4920
С	0.9620	0.4710
d	0.9590	0.4490
e	0.9190	0.3470
f	0.9540	0.4090
g	0.7910	0.2970
i	0.7430	0.2190
j	0.9650	0.4840
k	0.9640	0.4720
1	0.9710	0.4640
m	0.9590	0.4640
n	0.9580	0.4670
О	0.9570	0.4220
p	0.9610	0.4690
q	0.9690	0.4820
r	0.9610	0.4440
S	0.9810	0.4840
t	0.9650	0.4670
u	0.9580	0.4140
v	0.9620	0.4350
W	0.9590	0.4860
X	0.9670	0.4770
У	0.9380	0.4040
Z	0.9680	0.4630

